

How effective are brief interventions in reducing alcohol consumption: does setting, practitioner group and content matter? Findings from a systematic review and meta-regression analysis

Lucy Platt, G.J. Melendez-Torres, Amy O' Donnell, Jennifer Bradley, Dorothy Newbury-Birch, Eileen Kaner, Charlotte Ashton

Corresponding Author: Lucy Platt, Department of Social and Environmental Health Research, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, 15-17 Tavistock Place, London, WC1H 9SH Tel: +44 (0) 207 958 8156 Fax: +44 (0) 207 580 4524 email: lucy.platt@lshtm.ac.uk

Co-authors:

G.J. Melendez-Torres, Departmental Lecturer in Evidence-Based Social Intervention, University of Oxford, Oxford, UK

Amy O'Donnell, Research Associate, Institute of Health and Society, Newcastle University, Newcastle, UK

Jennifer Bradley, Research Assistant, Institute of Health and Society, Newcastle University, Newcastle, UK.

Dorothy Newbury-Birch, Professor of Alcohol and Public Health Research, University of Teesside, Teesside, UK.

Eileen Kaner, Institute Director and Professor of Public Health Research, Institute of Health and Society, Newcastle University, Newcastle, UK

Charlotte Ashton, Consultant in Public Health, Camden & Islington Public Health, London, UK

Key words: alcohol, systematic review, meta-analysis, multi-level model, brief interventions

Word count: 4544 words

Abstract

Background: While the efficacy and effectiveness of brief interventions for alcohol (ABI) have been demonstrated in primary care, there is weaker evidence in other settings and reviews do not consider differences in content. We conducted a systematic review to measure the effect of ABIs on alcohol consumption and how it differs by setting, practitioner group and content of intervention.

Methods: We searched MEDLINE, Embase, PsycINFO; CINAHL, Social Science Citation Index, Cochrane Library and Global Health up to January 2015 for randomised controlled trials that measured effectiveness of ABIs on alcohol consumption. We grouped outcomes into measures of quantity and frequency indices. We used multilevel meta-analysis to estimate pooled effect sizes and tested for the effect of moderators through a multiparameter Wald test. Stratified analysis of a sub-set of quantity and frequency outcomes was conducted as a sensitivity check.

Results: 52 trials were included contributing data on 29,891 individuals. ABIs reduced the quantity of alcohol consumed by 0.15 standard deviations. While neither setting nor content appeared to significantly moderate intervention effectiveness, provider did in some analyses. Interventions delivered by nurses had the most effect in reducing quantity ($d=-0.23$, 95% CI [-0.33, -0.13]) but not frequency of alcohol consumption. All content groups had statistically significant mean effects, brief advice was the most effective in reducing quantity consumed ($d=-0.20$, 95% CI [-0.30, -0.09]). Effects were maintained in the stratified sensitivity analysis at first and last assessment time.

Conclusion ABIs play a small but significant role in reducing alcohol consumption. Findings show the positive role of nurses in delivering interventions. The lack of evidence on impact of content of intervention reinforce advice that services should select the ABI tool that best suits their needs.

Article Summary

Strengths and Limitations of the study

A key strength of this review is the methodologically innovative approach to the meta-analysis through the use of a multilevel meta-analysis.

As a second sensitivity analysis we compared findings from the multi-level model with a stratified analysis focussing on a sub-set of outcome variables. Findings from the two analyses were comparable.

Quality assessment criteria were used to assess risk of bias and the majority of studies were at low risk in relation to the randomisation procedure and monitoring of loss to follow-up.

A large proportion of studies did not provide information on other aspects of the study design including blinding of participants to the intervention, intention to treat analysis and blinding to outcome measurement.

Our review suggested limited effect for interventions delivered in community settings, but relied on a small number of studies across a wide variety of settings.

What we already know on the topic

Screening to detect individuals drinking alcohol at hazardous or harmful levels and the delivery of a brief intervention on alcohol (ABI) to reduce their consumption have been implemented in primary care settings where their efficacy and effectiveness have been demonstrated.

There is weaker evidence for effectiveness beyond primary care, with moderate or no effect found in accident and emergency departments, college, community and general hospital settings.

Content of ABI is varied but usually focuses on structured advice involving an assessment of individual risk with feedback and advice, or brief motivational interviewing that takes a more patient-centred approach or a combination of both approaches. Existing evidence has not found much variability in effect by duration of intervention but this has not taken account differences in content.

What this study adds

Provider of the intervention does appear to matter in some outcomes, and in multilevel models interventions delivered by nurses had the greatest effect in reducing quantity of alcohol consumed ($d=-0.23$, 95% CI [-0.33, -0.13]).

Little evidence on the effectiveness of brief interventions in community settings or accident and emergency were found. University settings were associated with the greatest reduction in alcohol consumption then primary care.

Brief advice was associated with the greatest reduction in alcohol consumption ($d=-0.20$, 95% CI [-0.30, -0.09]) in the multi-level model and stratified analysis, but not in reducing frequency of drinking. However, overall neither setting nor content appeared to significantly moderate intervention effectiveness.

Introduction

Excessive alcohol consumption is a major public health concern, contributing to almost 4% of deaths worldwide(1), ranging from as high as 8% of deaths among men and women in the USA and Norway to 1.4% in the UK.(2, 3) It is estimated that over ten million people in the UK alone drink more than the recommended daily units.(4) Screening to detect individuals drinking alcohol at hazardous or harmful levels and the delivery of a brief intervention on alcohol (ABI) to reduce their consumption have been implemented in primary care settings where their efficacy and effectiveness have been demonstrated.(1) The content of ABIs is varied, but usually focuses on the provision of structured advice, involving an assessment of individual risk with feedback and advice, or brief motivational interviewing that takes a more patient-centred approach, or a combination of both.(5) Existing systematic reviews have found variability in effect by duration of intervention or number of visits, but this has not taken into account differences in content or provider.(6-10) Although there is some emerging evidence that motivational interviewing can be more effective than ‘traditional’ advice (based on a provider-centred definition of a problem) across a range of health behaviours(11), this is not conclusive.(12) Further, while the efficacy and effectiveness of ABIs have been demonstrated in primary care settings,(13-15) the evidence-base in health settings beyond primary care is weaker with moderate or no effect found in college(16, 17) and community settings.(18) Some benefits have been observed from a small number of studies in accident and emergency (A&E) departments(19, 20), as well as in general hospital settings but among mainly male patients.(21, 22) Implementation research has shown that contextual factors affecting the routine delivery of ABIs in primary health care settings are closely linked to practitioners. However, there has been little research looking at the impact of practitioners on intervention effectiveness outside primary health care settings.(23, 24)

In England, the Government’s Alcohol Strategy calls for the increased implementation of ABIs in Primary Care and A&E settings, while targets for implementing ABIs in these settings as well as antenatal clinics have been set by NHS Scotland.(25, 26) NICE guidance recommends that ABI should be offered opportunistically by a range of relevant practitioners and front-line staff, while also acknowledging that the strength of evidence was clearer in some health settings compared to others. Nevertheless, this guidance flagged the relevance of social care, criminal justice, community and voluntary sector professionals to supporting alcohol risk-reduction work.(27) This recommendation has been implemented by some Public Health Authorities, rolling out interventions in sexual health clinics and community settings such as criminal justice services, and has also been advocated by global health agencies including the WHO.(28) Given the international, national and local level support for the expansion of ABIs beyond primary care settings, there is an urgent need to understand how brief intervention process (including setting, provider and content) moderates their effectiveness in order to inform their implementation.(5) We therefore undertook a systematic review and meta-regression to measure the effect of ABIs on alcohol consumption and how effect differs by setting, provider group and content of intervention.

Methods

Search strategy and selection criteria

We followed the PRISMA guidelines on reporting of systematic reviews.(29) Studies eligible for this review were peer reviewed randomised controlled trials of ABIs published in English. We included all populations aged 16 years or older but excluded populations with complex health problems, for example studies of people living with HIV, TB, HCV or homeless populations where it is difficult to generalize findings to the general population. Similarly we excluded populations seeking help at specialist addiction, mental health services or antenatal clinics. We included studies with control groups comprising: treatment as usual; information-only; assessment only; no assessment; or non-intervention, and excluded control groups consisting of other interventions, including other brief

interventions such as advice and extended psychological treatments. Brief interventions were defined as *person-to-person* discussions on alcohol between one and four sessions and not more than two hours total intervention time. Computerized interventions tested alone, group interventions and those that target multiple behaviours were excluded. We also excluded studies where no measure of alcohol consumption was reported.

The primary outcome of interest was a quantitative continuous measure of total alcohol consumption within a specified time-frame (standard drinks, grams of ethanol, or days of drinking) where the standardized mean difference between brief intervention and control group was measured at time of follow up.

We searched: MEDLINE; Embase; PsycINFO; CINAHL; Social Science Citation Index and Science Citation Index through Web of Science; Cochrane Effective Practice and Organisation of Care Group specialised register; and Global Health between 1966 and 2015. The search was conducted in January 2015. We also scanned citations and contacted experts in the field to minimise selection bias. The search terms used were: 'Brief intervention' OR 'minimal intervention' OR 'early intervention' OR 'cognitive behavioural' OR 'screening' OR 'counselling' OR 'brief advice' OR 'identification' OR 'managed care' or 'motivational interview' AND 'Alcohol drinking' or 'binge drinking' OR 'alcohol consumption' OR 'alcohol units' OR 'alcohol use and misuse' OR 'alcohol intake' OR 'alcohol rate binge drinking' OR 'beer or wine or lager or spirit drinking' AND 'randomized controlled trial' OR 'random allocation' OR 'double blind methods' OR 'clinical trial' OR 'controlled clinical trial' OR 'multi centre studies'. Searches were tailored to the search functionality of each database (see Web Appendix).

Eligibility assessment was conducted independently by two reviewers. Disagreements between reviewers were resolved by consensus. We selected a list of risk of bias criteria from recommendations in the Cochrane Collaboration Reviewers' Handbook to assess the quality of the trials.⁽³⁰⁾ Criteria included: methods used to generate the allocation sequence to produce comparable groups and concealment of allocation to determine whether intervention allocations could have been foreseen before or during enrolment; blinding of participants and providers to intervention groups; blinding of outcome assessment; incomplete outcome data (including intention to treat analysis); and measurement of attrition rate.

Data were extracted from each publication into a database piloted on five studies, independently by GJM, LP, AO and JB without masking of authors' names, study site, intervention, or trial results. These researchers jointly reviewed the extracted data and 10% of studies were double extracted. Data were extracted on characteristics of trial participants, type of interventions (including content, duration, frequency, provider, setting), type of outcome measure, time of assessment and effect estimates.

We extracted continuous outcomes in the units in which they were presented and then converted to Cohen's *d* for comparability. When extracting continuous outcomes, we preferred estimates that were ANCOVA-adjusted for baseline score, followed by unadjusted post-test scores, and finally repeated measures or 'change score' models. Change score models were reparametrized into a raw-score metric using $r=0.5$, with sensitivity analysis at $r=0.1$ and $r=0.9$. Though past reviews have attempted to convert all measures to 'natural units' such as grams of ethanol, we decided that this was inadvisable because of the large number of trials in this review and because of our goal to include all relevant information, a key benefit of multilevel meta-analysis models.

Data synthesis

We grouped intervention content into three categories (Figure 1). The first was motivational interviewing, including motivational interviewing-style, advice approaches such as FRAMES, motivational enhancement therapy as adapted for Project MATCH (Project MATCH Research Group,

1998) or brief motivational interviewing. We also identified a second subset of trials that tested specific enhanced interventional protocols for motivational interviewing (e.g. Drink-less) or additions to motivational interviewing (e.g. cognitive behavioural approaches) from other therapeutic modalities and labelled this category motivational interviewing 'plus'. A third subset included brief advice approaches, often labelled as such without any additional information.

Intervention providers were grouped into: counsellors (defined as any mental health providers including clinical and research psychologists or clinical social workers); GPs (including primary care providers and general physicians); nurses (including research or clinical nurses on secondment); peer-delivered; and different providers (but with no fixed provider). Setting of intervention delivery was categorised as: accident and emergency services; community-based delivery that included a range of non-clinical settings; primary or ambulatory care delivered in clinical settings as outpatient services; hospital inpatient services; and university services.

The systematic review protocol was registered on PROSPERO at the University of York (CRD42014014799).

Statistical analyses

We grouped outcomes hierarchically. We identified an overarching set of outcomes addressing quantity of alcohol consumption, from which we created two subsets of outcomes: (i) amount of alcohol consumed per unit of time; and (ii) amount of alcohol consumed per drinking occasion. We also identified an overarching set of outcomes addressing frequency of alcohol consumption, from which we created a subset of outcomes including: (i) frequency of any drinking occasion; and (ii) frequency of binge drinking occasions.

For each overarching set and subset of outcomes, we specified five models: 1) an unconditional model that included all eligible continuous outcomes; 2) a model that included a grand mean-centred covariate for time of follow-up post-baseline, to address differences in follow up; 3) a model including where the intervention was initially delivered and time of follow-up; 4) a model including the provider of the intervention and time of follow-up; and 5) a model including the content of the intervention and time of follow-up. To estimate mean effects for all groups simultaneously, we refit models with no intercept.⁽³¹⁾ We used the statistical package metafor,⁽³²⁾ which implements advanced meta-analysis models, in the R environment for all multilevel analyses.

For our main analysis, we used a multilevel meta-analysis method to estimate pooled effect sizes.⁽³³⁾ Models included random effects on the effect size and study levels because of anticipated heterogeneity both within and across studies. Several trials tested different intervention or provider types in the same experiment, but insufficient trials did this to treat intervention as a 'within-trial' covariate. In order to adequately model these two moderators, we split the control groups in two for these trials and treated each intervention-control comparison as a separate trial. This avoided double-counting participants across intervention-control comparisons. Moreover, several studies presented results stratified by group. In our multilevel meta-analyses, we included these in the same cluster. We tested for the effect of our hypothesised moderators by conducting a multiparameter Wald test on provider, setting or content coefficients as appropriate. We additionally examined the residual heterogeneity, measured as I^2 , between the time-adjusted model and the models including each of the three sets of covariates. We regarded a p-value of <0.05 as statistically significant and a p-value of <0.10 as marginal, but not significant.

Sensitivity check

In addition to sensitivity analysis on the correlation used for repeated measures conversion, we estimated a set of meta-regressions for each subset of outcomes including one effect size per relevant comparison for each of first and last follow-up in the included trials. We did this by combining

intervention and control groups where appropriate, and by selecting effect sizes within studies that used shorter time periods for measurement and timeline follow-back procedures over general frequency/quantity questionnaires. We also treated non-overlapping subgroups from the same study as separate data points as suggested by Borenstein et al.(34) Sensitivity analyses were estimated in both Stata v 13.1 (Stata Corp. 2013) and R .(35) We did not undertake meta-analysis of effect sizes from common time points because these models would have been poorly powered.

Results

We identified a total of 4551 records from the search of electronic databases and 41 records from key experts. A total of 52 studies met our inclusion criteria, with three studies presenting different outcomes for the same data and therefore considered as one. (36-38) One study was dropped as it only contained biological outcomes which were not included in the main analyses.(39) The review and selection process is summarised in Figure 2.

Included studies contributed data for 29,891 individuals. Table 1 presents a summary of study characteristics (country, age, sex and sample size) as well as type of intervention (setting, provider and content), key outcomes and time of assessment. Most studies originated from Europe or North America with the exception of three studies from Australia, Taiwan and Thailand.(40-42) Almost half (45%) of the studies were conducted in the USA and 22% in the UK.

Table 1 Characteristics of Included studies

Author	Country	Sample n [†] % F, age (yrs)	Intervention					Outcomes	
			Setting	Provider	Ar m	Content	Total mins (sessions)	Definition (Q=Quantity, F= Frequency)	Time (mths)
Aalto 2000(43)	Finland	118, 100%, 41	GP	GP/nurse	1	MI	70-130 (7)	Q: Amount per week; usual amount per occasion (Grams); F: Drinking times per week	36
				GP only	2	MI	30-60 (7)		
				N/A	C	TAU			
Aalto 2001(44)	Finland	296, 0%, 41	GP	GP/ nurse	1	MI	70-130 (7)	Q: Grams per week/ per occasion F: Drinking times per week	36
				GP only	2	MI	30-60 (7)		
				N/A	C	TAU			
Anderson 1992(45)	UK	154, 0%, 44	GP	GP	1	Brief advice	10 (1)	Q: Breath alcohol (mg/100 ml); HSQ quant/freq and interview (grams/week)	12
					2	TAU			
Antti-Poika 1988(46)	Finland	120, 0%, 39	A&E	Nurse	1	Brief advice	NR (1)	Q: Grams of absolute alcohol during 1 week period	6 (P-I)
Baer 2001(47)	USA	508, 55%, NR	College	Counsellor	1	MI	Unclear (NR)	Q: Mean drinks per drinking day; F: Drinking days per average week	24; 36
					C	Screening			
Beich, 2007(48)	Denmark	6897, 62%, 36	GP	GP	1	MI Plus	10 (1)	Q: Usual weekly consumption of beer, wine and spirits (units/week)	12
					C	Screening			
Bernstein 2010(49)	USA	835, 56%, 88%>18	A&E	Peer	1	MI	Unclear (1)	Q: Max drinks per day; Mean drinks per drinking day; Mean drinks per week F: Drinking days per month	3, 12
					C	Screening			
Butler 2009(50)	USA	114, 65%, 20	College	Media	1	Brief advice	11 (1)	Q: Standard drinks per week; F: Binge episodes; drinking occasions; drinking occasions	1 (P-I)
				Counsellor	2	MI	41 (1)		
				N/A	C	Screening			
Carey 2006(51)	USA	509, 65%, 19	College	Counsellor	1	MI	65 (1)	Q: Drinks per drinking day; F: Drinks per week; Heavy drinking frequency	6 or 12
					2	MI Plus	70 (1)		
					C	Screening			
Cherpitel 2010(52)	Poland	446, 17%, 54% >30	A&E	Nurse	1	MI plus	15-20 (3)	Q: Drinks per drinking day; Maximum drinks per occasion last month; F: Drinking days per week	12
					C	Screening			
					C	Assessment			
Chick 1985(53)	UK	156, 0%, 18-65	A&E	Nurse	1	Brief advice	60 (1)	Q: Consumption on past week (units)	12
					2	Screening			
Cordoba 1998(54)	Spain	229, 0%, 36.5	GP	GP	1	Branded	15 (1)	Q: Alcohol consumption units/week	12
					C	Simple advice			

Author	Country	Sample n [†] % F, age (yrs)	Intervention					Outcomes	
			Setting	Provider	Ar m	Content	Total mins (sessions)	Definition (Q=Quantity, F= Frequency)	Time (mths)
Crawford 2004(55)	UK	599, 21%, 44	A&E	Nurse	1	MI	30 (3)	Q: Mean units per drinking day; Mean weekly units	6 or 12
					C	Information			
Crawford 2014(56)	UK	802, 54%, 27	GP	Nurse	1	Brief Advice	2-3 (1)	Q: Mean units on drinking days; Weekly alcohol consumption in units	6
					C	Information			
Curry 2003(57)	USA	333, 35%, 47	GP	GP and counsellor	1	MI Plus	47 (1)	Q: Drinks per week	12
					C	TAU			
Daeppen 2007(58)	Switzerland	987, 22%, 36.7	A&E	Counsellor	1	MI	17 (1)	Q: Number of drinks per occasion/last week (last year) F: Number of binge drinking occasions per month/per week (last year)	12
					C	Assessment			
					C	Nothing			
Daeppen 2011(59)	Switzerland	2831, 0%, 19.9	Community (Military)	Counsellor	1	MI	15.8 (2)	Q: Change in drinks per week F: Change in binge drinking occasions per month	6
					C	Assessment			
Drummond 2014(60)	UK	1204, 35%, 34.6	A&E	Counsellor	1	Branded	20 (1)	Q: Average daily drinks	6, 12 (P-I)
					2	MI			
					C	Information			
Field 2010(61)	USA	1439, 18%, 33	A&E	Counsellor	1	MI	Unclear (1)	Q: Change in: alcohol per week; max. amount in a day in past 6 mths; F: Change in percent days heavy drinking;	6, 12
					C	TAU + Assess			
Fleming 1997; Manwell 2000, Grossberg 2004(36-38)	USA	774, 38%, 29% 18-30	GP	GP and nurse	1	Branded	30 (2)	Q: No. drinks in past 7 days F: No. binge drinking episodes in last 30 days [binge drinking defined as having more than 4 drinks per occasion]	6, 12, 24, 36, 48 (P-I)
					C	Information			
Fleming 1999(62)	USA	158, 34%, 65- 75	GP	GP / nurse	1	Branded	30 (2)	Q: Number of drinks in last week; F: Number of binge drinking occasions in last month;	6, 12
					C	Information			
Fleming 2010(63)	USA	986, 51%, 21	GP	GP	1	Branded	30 (2)	Q: Mean number of drinks; F: Mean number of drinking days; Mean number of heavy drinking days (last 28 days)	6
					C	Information			
Freyer-Adam 2008(64)	Germany	595, 6%, 41	Hospital	Different providers*	1	MI	unclear (1)	Q: Average daily alcohol intake (grams); Total alcohol intake in past week (grams)	12
					2	MI			
					C	TAU			
Gaume 2011(65)	Switzerland	572, 0%, 19.9	Community (Military)	Counsellor	1	MI [^]	21.8 (1)	Q: Mean change in number of standard (~10 g of alcohol) drinks per week; F: Mean change in heavy episodes (6 drinks or more) per month	6
					2	MI [^]			
					C	Assessment			

Author	Country	Sample n [†] % F, age (yrs)	Intervention					Outcomes	
			Setting	Provider	Ar m	Content	Total mins (sessions)	Definition (Q=Quantity, F= Frequency)	Time (mths)
Gaume 2014(66)	Switzerland	431, 0%, 19	Community (Military)	Different providers	1	MI	20-30 (1)	Q: Number of drinks/day F: Number of drinking days/week	3
Gentilello 1999(67)	USA	762, 18%, 35.4	A&E	Counsellor	1	MI	30 (1)	Q: Changes in the no. of drinks consumed per week	6, 12
					C	Assessment			
Gottlieb- Hansen 2012(68)	Denmark	772, 49%, 60	Community (Research)	Different providers	1	MI	15 (2)	Q: Number of drinks per week	6, 12
					C	Information			
Heather 1987(69)	UK	104, 25%, 36.4	GP	GP	1	Branded	NR	Q: Heaviest months consumption in last 6 months(units); Last month's consumption (units)	6
					2	Brief advice			
					C	Assessment			
Holloway 2007(70)	UK	215, 15%, 44	Hospital	Nurse	1	MI	20 (1)	Q: Change from baseline in alcohol units in the past 7 days; Change in maximum units in 1 day F: Change in drink days in last week;	6
				Media	2	Media	NR		
				N/A	C	TAU			
Ingersoll 2013(71)	USA	217, 100%, 27.9	Community (Research)	Counsellor	1	MI, Branded	60 (1)	Q: Drinks per drinking day	3, 6
				N/A	C	Information			
Juarez 2006(72)	USA	122, 53%, 19.4	College	Counsellor	1	BA, MI	60-80 (1)	Q: Drinks per day; Maximum BAC	2
				Counsellor	2	BA, MI	40-60 (1)		
				Media	3	BA, MI	Unclear (1)		
				Counsellor	4	BA, MI	40-60 (1)		
				N/A	C	Assessment			
Kulesza 2010(73)	USA	114, 72%, 20	College	Counsellor	1	MI Plus	10 (1)	Q: The Daily Drinking Questionnaire;	6 wks
					2	MI Plus	50 (1)		
					C	Waiting List			
Kulesza 2013(74)	USA	268, 71%, 20	College	Counsellor	1	MI Plus	10 (1)	Q: Average no. drinks/week	4 wks (P- I)
					2	MI Plus	50 (1)		
					C	Discussion			
Larimer 2001(75)	USA	159, NR< 18.8	College	Peer	1	MI Plus	60 (1)	Q: BAC (based on quantity & rate of consumption peak; Number of drinks over past month; Total average use; F: Frequency of use	12
					2	MI Plus	60 (1)		
					C	TAU			
Liu 2011(40)	Taiwan	616, 0%, 41	A&E	Counsellor	1	MI	60 (2)	Q: No. drinks in last 3 months (QDS); F: No. days heavy drinking (≥5 drinks) in last 3 mths (QDS); No. days heavy drinking in the previous week (TLFB)	4
					C	TAU			

Author	Country	Sample n [†] % F, age (yrs)	Intervention					Outcomes	
			Setting	Provider	Ar m	Content	Total mins (sessions)	Definition (Q=Quantity, F= Frequency)	Time (mths)
Lock 2006(76)	UK	127, 100%, 44.1	GP	Nurse	1 C	Branded TAU	5-10	Q: Units per week	12 (P-I)
Maisto 2001(77)	USA	301, 31%, 45.6	GP	Researcher Counsellor N/A	1 2 3	Brief advice MI Control	10-15 (1) 60-85 (1)	Q: No. of drinks in last 30 days F: No. of days of 1-6 drinks in last 30 days;	6, 12
Murphy 2001(78)	USA	99, 54%, 19.6	College	Counsellor Counsellor N/A	1 2 C	MI Brief advice Assessment	45 (1) 50 (1)	Q: Drinks per week; F: Binge drinking days per week [4+ drinks for women; 5+ drinks for men]; Drinking days per week;	9
Noknoy 2010(41)	Thailand	59, 9%, 37	GP	Nurse	1 C	MI Assessment	45 (3)	Q: Average drinking per drinking day during the previous week (drinks/drinking day)	6
Richmond 1995(42)	Australia	378, 43%, 37.7	GP	GP	1 2 C	Branded Brief advice Nothing	30-55 (1) 5 (1)	Q: No units of ethanol in the last 7 days	6, 12
Rubio 2010(79)	Spain	752, 35%, 18- 65	GP	GP	1 C	Branded Information	20-30 (2)	Q: No. of drinks in last 7 days [mean/SD]; F: No. of binge episodes (last 30 days) [mean/SD] (> 4 drinks for women and 5 for men in a single occasion)	12
Rubio 2014(80)	USA	330, 100%, 24	GP	Different providers	1 2	MI Control	70 (1)	Q: Drinks per day	6 wks, 6, 12 PP
Saitz 2007(81)	USA	341, 29%, 45	Hospital	Counsellor	1 C	MI TAU	30 (1)	Q: Change decrease in number drinks/day F: Change decrease in heavy drinking episodes	12
Schaus 2009(82)	USA	363, 52%, 20.6	GP	GP	1 C	MI Plus Information	40 (2)	Q: Average drinks per sitting/week; Typical BAC; Peak BAC; Peak no. drinks in sitting; F: No. days drinking 4+ drinks in month; No. times drunk in typical week.	6, 9
Senft 1997(83)	USA	516, 30%. 41.9	GP	GP/ counsellor	1 C C	MI TAU Referral to GP	15 (1)	Q: Drinks/drinking day over past 6 months; Total SECs past 3 months; F: Drinking days/week over past 6 months;	6, 12
Shiles 2014(84)	UK	154, NR, 51	Hospital	Nurse	1 C	Brief advice TAU	10 (1)	Q: Daily units of alcohol in last week	3, 12
Smith 2003(85)	UK	151, 0%, 24	Hospital	Nurse	1 C	MI TAU	NR	Q: 84-day alcohol consumption; Alcohol consumption in a typical week	3, 12
	USA	152, 45%, 20.9	College	Media	1	MI	45 (1)		10 wks

Author	Country	Sample n† % F, age (yrs)	Intervention					Outcomes		
			Setting	Provider	Ar m	Content	Total mins (sessions)	Definition (Q=Quantity, F= Frequency)	Time (mths)	
Wagener 2012(86)				Counsellor	2	MI	105-135 (1)	Q: Peak BAC; Typical BAC) Weekly alcohol consumption using DDQ		
				Counsellor	3	MI	NR (1)			
				N/A	C	Assessment				
Walters 2009(87)	USA	279	College	Counsellor	1	MI (no feedback)	40 (1)	Q: No. of drinks per week; Peak BAC	3, 6	
				Counsellor	2	MI (feedback)				
				Media	3					
					C	Assessment				
Watt 2008(88)	UK	269	Community (CJS)	Different providers	1	MI	15-20	Q: No. of units consumed per week; F: Number of drinking days in the past 3 months	3, 12 (PI)	
					C	NR				

NR= not reported

Sample: † n denotes eligible sample randomised at baseline; F=Female

Setting: CJS=Criminal Justice Service GP=General Practice; A&E=Accident and Emergency;

Providers: *Different providers defined as (Psychologist, Social Worker or Research nurse)

Content: ^ Stratified by heavy episodic and non-heavy episodic users. TAU= Treatment as usual; BA Brief advice; MI= Motivational Interviewing

Arm: C=Control group

Outcome: QDS=Quick Drinking Screen; TFLB=Alcohol Timeline Follow-Back; DDQ=Daily Drinking Questionnaire

Outcome time = All outcomes measured in months post baseline, unless specified: PI = Post Intervention; wks=weeks, PP= Post partum

In total, 68% of trials were delivered in primary or healthcare settings (hospital or A&E). Only six studies were conducted in community settings defined as: military(59, 65, 66); research sites, recruiting a sample through a household survey(68); women at risk of alcohol exposed pregnancy (defined as aged 18-44 years, with ineffective or no use of contraceptives, sexually active in the last 6 months, but not currently pregnant or planning a pregnancy) recruited via the media, in a prison, community health centre and a gynaecology centre (71); and one criminal justice setting.(88) The most common providers included counsellors, who were the sole providers of interventions in 43% of trials, and physicians who accounted for 24% of trials. A minority category of different providers (8%) included a combination of psychologists, social workers or research nurses. Intervention categories were well-distributed, though a majority of trials (47%) included motivational interviewing alone and 39% included motivational interviewing 'plus'. A total of 50 trials reported 275 eligible effect sizes on outcomes measuring quantity of alcohol consumed with a mean follow up of nine months. This is summarised in Table 2.

Table 2: Summary of study characteristics

TRIALS		
Setting of intervention		
A&E	20%	10
Non-health settings	12%	6
Ambulatory or primary care	38%	19
Hospital inpatient services	10%	5
University	20%	10
Provider		
Counsellor/mental health clinician	44%	22
Different providers	8%	4
GP	22%	11
Nurse	18%	9
Peer intervention	4%	2
Combination	12%	6
GP and nurse	8%	4
GP and counsellor	4%	2
Content		
Brief advice	24%	12
Motivational interviewing	48%	24
Motivational interviewing 'plus'	40%	20
OUTCOMES		
Quantity		50
Mean follow-up in months (SD)	9.0 (8.3)	
Quantity per unit time	94%	47
Quantity per drinking occasion	30%	15
Frequency		26
Mean follow-up in months (SD)	11.1 (10.5)	
Frequency of any drinking occasion per unit time	32%	16
Frequency of binge drinking occasions per unit time	30%	15

A & E= Accident and Emergency GP=General Practice SD=Standard Deviation

The majority (71%) of studies were categorised as low risk of bias in relation to randomisation and allocation concealment strategies. In the majority of studies the process used to assess blinding of participants and providers as well as outcome assessment was unclear. Intention to treat analysis was conducted in 47% of studies and loss to follow-up assessed in the majority (80%) of studies. This is summarised in Table 3 and risk of bias assessment for all trials is included in the Web Appendix (Online Table 1)

Table 3: Summary risk of bias assessment

Risk of bias indicator	Score - Proportion (number of estimates)		
	High risk % (k)	Low risk % (k)	Unclear risk % (k)
Allocation concealment	2% (1)	72% (36)	26% (13)
Blinding of participants and providers	12% (6)	30% (15)	58% (29)
Blinding of outcome assessment	10% (5)	42% (21)	48% (24)
Intention to treat analysis	6% (3)	48% (24)	46% (23)
Loss to follow up	20% (10)	80% (40)	0% (0)

Meta regression on combined quantity and frequency outcomes

Interventions produced a beneficial effect at reducing the quantity of alcohol consumed by 0.15 standard deviations— a small but statistically significant effect (see Table 4). This effect persisted after controlling for time to follow-up and when examining the sub-set of outcomes. In both unconditional models and models controlling for time of follow-up, study-level heterogeneity as measured by I^2 (that is, the percentage of variation between effect sizes due to heterogeneity rather than chance) was in the small to moderate range (0-40%) as defined by the Cochrane Handbook.(30) Findings were robust to sensitivity analysis on the pre-post correlation in change score models. The mean time-adjusted effect of brief alcohol interventions on frequency of alcohol consumption outcomes was similar in magnitude ($d=-0.15$, 95% CI [-0.20, -0.11]), but lower in heterogeneity ($I^2=23%$), compared with the effect on quantity of alcohol consumption (see Table 5). The time-adjusted effect remained statistically significant when limited to the sub-set of outcomes (frequency of drinking occasions $d=-0.12$, 95% CI [-0.19, -0.06] and frequency of binge drinking $d=-0.17$, 95% CI [-0.23, -0.11]).

Table 4 Results of multi-level meta-regression for quantity outcomes

Outcomes	Group name	All quantity outcomes				Quantity of alcohol per unit time				Quantity of alcohol per drinking occasion			
		ES (95% CI)	k (n)	I ² (%)	p	ES (95% CI)	k (n)	I ² (%)	p	ES (95% CI)	k (n)	I ² (%)	p
Overall	Mean effect	-0.15 (-0.20, -0.10)	50 (268)	37%		-0.17 (-0.22, -0.12)	47 (144)	38%		-0.10 (-0.18, -0.01)	15 (59)	36%	
Overall, time-adjusted	Mean effect	-0.15 (-0.20, -0.11)	50 (268)	36%	0.03	-0.17 (-0.22, -0.12)	47 (144)	38%	0.21	-0.11 (-0.19, -0.03)	15 (59)	34%	0.09
	Time (month)	0.003 (0.0003, 0.006)				0.002 (-0.001, 0.006)				0.005 (-0.001, 0.01)			
Setting of intervention	A&E	-0.10 (-0.19, -0.002)	10 (44)	34%	0.12	-0.12 (-0.22, -0.01)	9 (26)	37%	0.17	-0.001 (-0.14, 0.13)	4 (8)	28%	0.17
	Ambulatory or primary care	-0.20 (-0.27, -0.13)	19 (84)			-0.22 (-0.29, -0.14)	19 (51)			-0.14 (-0.25, -0.03)	7 (18)		
	Hospital inpatient services	-0.14 (-0.29, 0.01)	5 (13)			-0.15 (-0.31, 0.006)	5 (12)			N/A	N/A		
	Non-health settings	-0.03 (-0.16, 0.10)	6 (15)			-0.04 (-0.18, 0.11)	5 (11)			-0.01 (-0.30, 0.29)	1 (4)		
	University	-0.20 (-0.39, -0.09)	10 (112)			-0.21 (-0.23, -0.09)	9 (44)			-0.22 (-0.39, -0.06)	3 (29)		
Provider	Counsellor/mental health clinician	-0.11 (-0.17, -0.05)	24 (163)	34%	0.09	-0.10 (-0.17, -0.04)	22 (79)	32%	0.01	-0.11 (-0.23, 0.01)	8 (41)	43%	0.67
	Different providers	-0.12 (-0.27, 0.03)	4 (10)			-0.12 (-0.25, 0.02)	4 (10)			N/A			

	Physician	-0.12 (-0.20, -0.04)	17 (65)			-0.14 (-0.22, -0.06)	17 (40)			0.02 (-0.16, 0.21)	6 (10)		
	Nurse	-0.23 (-0.33, -0.13)	13 (41)			-0.28 (-0.38, -0.18)	12 (29)			-0.18 (-0.37, -0.003)	5 (9)		
	Peer intervention	-0.08 (-0.29, 0.13)	2 (10)			-0.05 (-0.28, 0.17)	2 (3)			-0.004 (-0.28, 0.27)	2 (3)		
Content	Brief advice	-0.20 (-0.31, -0.09)	12 (26)	39%	0.5 4	-0.22 (-0.34, -0.11)	11 (18)	59%	0.3 1	-0.16 (-0.37, 0.05)	3 (6)	43%	0.8 9
	Motivational interviewing	-0.13 (-0.19, -0.07)	24 (132)			-0.13 (-0.20, -0.07)	24 (73)			-0.11 (-0.22, 0.004)	9 (28)		
	Motivational interviewing plus	-0.16 (-0.23, -0.09)	20 (110)			-0.19 (-0.27, -0.11)	17 (53)			-0.10 (-0.24, 0.03)	6 (25)		

k =number of studies, n =number of effect sizes, p is the value from a multiparameter Wald test of coefficients. Models for setting, provider and content include mean-centred time as a covariate, but not in the multiparameter Wald

Setting

For all quantity outcomes setting of intervention did not appear to fully explain heterogeneity between studies, with residual heterogeneity at 34% and a statistically marginal but non-significant joint test of moderators ($p=0.09$). Interventions conducted in university settings ($d=-0.20$, 95% CI [-0.39, -0.09]) and in primary or ambulatory care (-0.20, [-0.27, -0.13]) appeared to be most effective, with a small but statistically significant effect of the intervention. Interventions delivered in community settings (military, criminal justice, research sites and targeted recruitment) did not appear to be effective (-0.03, [-0.16, 0.10]). (Table 4)

For all frequency outcomes, setting of intervention did not explain heterogeneity (residual $I^2=25%$, Wald $p=0.54$). Of subgroups with statistically significant pooled effect sizes, interventions delivered in university contexts appeared to be most effective for frequency outcomes (-0.21, [-0.33, -0.08]). Analysis was hampered by small numbers of studies in several categories. (Table 5)

Table 5 Results of multi-level meta-regression for frequency outcomes

Outcomes	Group name	All quantity outcomes				Quantity of alcohol per unit time				Quantity of alcohol per drinking occasion			
		ES (95% CI)	k (n)	I ² (%)	p	ES (95% CI)	k (n)	I ² (%)	p	ES (95% CI)	k (n)	I ² (%)	p
Overall	Mean effect	-0.15(-0.20, -0.11)	26 (114)	23%		-0.12 (-0.19, -0.06)	16 (38)	23%		-0.17 (-0.23, -0.11)	15 (76)	20%	
Overall, time-adjusted	Mean effect	-0.16 (-0.20, -0.11)	26 (114)	23%	0.36	-0.12 (-0.19, -0.06)	16 (38)	24%	0.55	-0.18 (-0.24, -0.11)	15 (76)	20%	0.56
	Time (month)	0.002 (-0.002, 0.005)				0.002 (-0.004, 0.007)				0.001 (-0.003, 0.006)			
Setting of intervention	A&E	-0.11 (-0.21, -0.005)	5 (26)	25%	0.54	-0.13 (-0.26, -0.0002)	4 (12)	28%	0.41	-0.11 (-0.22, 0.01)	3 (14)	20%	0.25
	Ambulatory or primary care	-0.18 (-0.26, -0.10)	10 (40)			-0.07 (-0.19, 0.06)	5 (12)			-0.24 (-0.33, -0.15)	6 (28)		
	Hospital inpatient services	-0.21 (-0.47, 0.04)	2 (2)			-0.50 (-0.94, -0.06)	1 (1)			-0.07 (-0.37, 0.23)	1 (1)		
	Non-health settings	-0.08 (-0.22, 0.06)	4 (7)			-0.11 (-0.32, 0.11)	2 (3)			-0.06 (-0.24, 0.13)	2 (4)		
	University	-0.21 (-0.33, -0.08)	5 (39)			-0.18 (-0.36, -0.003)	4 (10)			-0.21 (-0.37, -0.05)	3 (29)		
Provider	Counsellor/mental health clinician	-0.11 (-0.17, -0.04)	14 (73)	23%	0.17	-0.12 (-0.22, -0.02)	9 (25)	32%	0.73	-0.12 (-0.20, -0.05)	9 (48)	18%	0.07
	Different providers	-0.24 (-0.52, 0.03)	1 (1)			-0.25 (-0.56, 0.07)	1 (1)			N/A			
	Physician	-0.13 (-0.22, -0.04)	10 (30)			-0.03 (-0.19, 0.13)	6 (8)			-0.18 (-0.28, -0.07)	5 (22)		
	Nurse	-0.19 (-0.31, -0.07)	7 (22)			-0.20 (-0.31, 0.01)	4 (5)			-0.17 (-0.31, -0.02)	3 (17)		
	Peer intervention	-0.06 (-0.27, 0.13)	2 (3)			-0.08 (-0.31, 0.16)	2 (3)			N/A			
Content	Brief advice	-0.08 (-0.26, 0.09)	3 (7)	29%	0.48	0.17 (-0.11, 0.44)	2 (4)	26%	0.10	-0.23 (-0.44, -0.02)	2 (3)	26%	0.52

	Motivational interviewing	-0.15 (-0.21, -0.08)	15 (58)			-0.15 (-0.23, -0.06)	9 (20)			-0.14 (-0.23, -0.06)	9 (38)		
	Motivational interviewing plus	-0.19 (-0.27, -0.11)	11 (49)			-0.13 (-0.24, -0.03)	7 (14)			-0.21 (-0.31, -0.11)	6 (35)		

k=number of studies, *n*=number of effect sizes, *p* is the value from a multiparameter Wald test of coefficients. Models for setting, provider and content include mean-centred time as a covariate, but not in the multiparameter Wald test.

When limiting the analysis to the sub-set of either quantity or frequency outcomes setting of intervention did not explain heterogeneity (all joint tests of moderators $p > 0.10$).

Provider

In the model including all quantity outcomes, provider of intervention did not meaningfully explain heterogeneity, based on I^2 for this model (34%). Interventions delivered at least in part by nurses appeared to have the largest effect by magnitude ($d = -0.23$, 95% CI [-0.33, -0.13]), though this difference was not supported by a significant joint test of moderators (Wald $p = 0.09$).

Analyses with more specific sets of outcomes revealed a similar picture. Examination of effects at first time point for amount of alcohol per unit time showed that interventions delivered at least in part by nurses ($d = -0.30$, 95% CI [-0.47, -0.12]) were the most effective, with a significant joint test of moderators (Wald $p = 0.048$) (Online Table 2). Interventions delivered by a range of different providers were least effective and did not yield a statistically significant effect. However, few studies were included in this category of providers. Provider of intervention explained some heterogeneity when the analysis was limited to amount of alcohol per unit time (residual $I^2 = 32\%$, Wald $p = 0.01$) but not per drinking occasion.

For frequency outcomes, provider of intervention did not explain heterogeneity either combined (Wald $p = 0.17$) or for drinking occasion per unit time (Wald $p = 0.73$) but the effect was marginal, but non-significant, for bingeing occasions (Wald $p = 0.07$).

Content

For quantity outcomes, content of intervention did not explain a statistically significant amount of heterogeneity (residual $I^2 = 39\%$, Wald $p = 0.54$), with little apparent reduction in I^2 . While all content groups had statistically significant mean effects, brief advice appeared to be most effective ($d = -0.20$, 95% CI [-0.30, -0.09]) with the impact of motivational interviewing ($d = -0.13$) and motivational interviewing plus ($d = -0.16$) also statistically significant.

For frequency outcomes, content of intervention did not explain a significant amount of heterogeneity (residual $I^2 = 29\%$, Wald $p = 0.48$). Effects by content group for motivational interviewing were similar to those in the analysis of quantity outcomes, though brief advice did not have a statistically significant effect on frequency of alcohol use ($d = -0.08$, [-0.26, 0.09]).

Estimates of heterogeneity remained the same when limiting the analysis to the sub-set of either quantity or frequency outcomes.

Sensitivity check: meta-regression on subset of outcomes by first and last time point

Overall effect estimates based on first and last time point were similar to the corresponding value reported in the main analysis, but estimates of heterogeneity (measured through I^2) tend to be higher. Setting of intervention explained some heterogeneity for the alcohol per unit time outcome at first time of marginal significance (residual $I^2 = 49\%$, Wald $p = 0.08$). Findings also suggest that provider explained some heterogeneity (residual $I^2 = 43\%$, Wald $p = 0.05$) with nurses having the biggest effect ($d = -0.30$, 95% CI [-0.41, -0.20]) and interventions delivered by different providers the least effect ($d = -0.07$, 95% CI [-0.12, -0.03]). Content of intervention explained some heterogeneity (residual $I^2 = 43\%$, Wald $p = 0.04$), brief advice was the most effective ($d = -0.25$, 95% CI [-0.42, -0.07]) and motivational interviewing least effective ($d = -0.09$, 95% CI [-0.15, -0.04]). (Figures 3-5) With the exception of content, evidence of heterogeneity did not remain significant at the last time point. There was no evidence of heterogeneity for alcohol consumed per drinking occasion or for either subset of frequency outcomes. All findings are summarised in the Online Tables 2-5.

Discussion

Our findings provide important new evidence on how the effectiveness of brief alcohol interventions differs by setting, provider and content, informing us of optimum modality. Our findings show that provider of intervention may matter. We observed some reductions in heterogeneity in the multi-level analysis of amount of alcohol consumed per unit time, and interventions delivered by nurses having the most effect in reducing quantity of alcohol consumed, but not frequency of consumption. This finding builds on other evidence showing a modest effect of brief interventions delivered by non-physicians (nurses and health care workers) in primary care settings.(24) We found that neither setting nor content appeared to significantly moderate intervention effectiveness. We found little evidence on the effectiveness of brief interventions in community settings or accident and emergency; brief advice was the most effective content in reducing quantity of alcohol consumed but not frequency of drinking; and there seemed to be little difference in the effect of motivational interviewing or MI Plus on either quantity or frequency outcomes.

While setting did not explain heterogeneity, findings show that university and ambulatory/ primary care settings were the most effective in terms of magnitude of effect size, which is supported by previous reviews in this area.(14, 15, 17) Prior research has suggested that while ABIs delivered in A&E settings may be effective in reducing alcohol consumption among hazardous and harmful drinkers,(19) it may not provide the most appropriate context for discussion on alcohol use.(89) The brevity of visits, lack of privacy for the delivery of the intervention and severity of injury may hinder the interaction between patient and practitioner reducing effectiveness.(89-92). Other evidence shows that discussion of drinking behaviours is facilitated by a good relationship between practitioner and client.(76) Our finding of increased reduction in alcohol consumption when the intervention is delivered by a nurse is important. The majority of previous research has focussed on physician-led interventions, but there is growing evidence to support the effectiveness of nurse-led interventions in both primary care and other settings.(24, 93, 94) As the largest group of health care workers with repeated patient contact and with a health promotion remit as part of their role, they are well placed to deliver ABIs.(93, 95) Barriers to nurses delivering the interventions include lack of time, worry about losing trust of the patient and inadequate training.(96, 97) Resources and training should be provided to support nurses to undertake this role and embed it within services. The provision of ABIs under the category of different providers was not associated with a reduction in consumption in alcohol. This may be related to problems with training of different providers, but the category was small and included a diverse range of providers, making the finding difficult to interpret. Similarly only a moderate effect was associated with counsellors, but again this definition encompasses a diverse group of practitioners ranging from clinical psychology students(78) to alcohol workers with specialist training in alcohol counselling.(60)

While our categories of intervention content did not meaningfully or statistically explain heterogeneity in either quantity or frequency outcomes in the multi-level analysis, they did in the stratified analysis for both first and last assessment time points. Effect sizes for quantity outcomes for all three classes of content were statistically significant, with brief advice yielding the largest effect. This provides important empirical evidence that brief advice can reduce alcohol intake, where evidence was lacking, and corroborates previous research that demonstrated no difference in effect between brief advice and longer motivational interviewing in reducing harmful levels of drinking in A&E, primary care and criminal justice settings.(12, 60, 98, 99)

Strengths and Limitations

A key strength of this review is the use of a multilevel meta-analysis method to integrate all relevant effect sizes from included studies. This circumvented problems in other systematic reviews around selection of specific effect sizes for meta-analysis. However, we were unable to explicitly model

correlation between outcomes within studies, though simulation evidence suggests that this may not have a large impact on estimation of intervention effects.(100) We used Cohen's *d* to standardise outcomes. While this is common across many systematic reviews addressing continuous outcomes, it is uncommon for systematic reviews of alcohol outcomes, where standardisation is often in terms of standard drinks or grams of ethanol consumed. This may somewhat limit comparability between reviews, but it was a critical step in employing the multilevel meta-analysis model we used. As a second sensitivity analysis we compared findings from the multi-level model with a stratified analysis focussing on a sub-set of outcome variables. Findings from the two analyses were comparable. The stratified analysis of quantity of alcohol consumed per unit time suggested stronger effects of setting, provider and content of intervention at first time-point of assessment than indicated in the multi-level models but with comparable effect estimates within each category. Tests for publication bias do not yet exist for multilevel meta-analyses. While our tests using all available effect sizes did not reveal significant publication bias on either quantity or frequency outcomes, it is unlikely that this is the best way to test publication bias in the context of dependent effect sizes. While we used the broadest categories appropriate for setting and provider of interventions, the number of studies included in meta-analysis examining frequency outcomes meant that meta-regressions were likely underpowered. We did not examine the effect of sex, ethnicity or age as a covariate since the sample size would have been too small to conduct a multivariate meta-regression analysis. As the number of trials grows, this meta-analysis should be repeated in order to better estimate differences between categories and examine the effect of other factors.

These findings should also be viewed in context of study-level heterogeneity. In our multilevel meta-analyses, heterogeneity was surprisingly low considering the diversity of settings, providers and modalities included in this body of evidence. One possible reason for this is that because we included all relevant outcomes, we avoided some of the 'random error' that may arise when only selecting one outcome per study. That is, including more information from each study will provide an estimate of statistical heterogeneity that more meaningfully accounts for study-level differences. This is not to say that it was inappropriate to explore this heterogeneity through structured and pre-hypothesised subgroup analyses, as was done here. Rather, the magnitude of difference in effects between studies may not be as pronounced as would be expected in a systematic review with such diverse interventions. While there was a low risk of bias in relation to some aspects of the study design (randomisation, loss to follow-up), there was a high percentage of unclear risk for many criteria, limiting our ability to fully assess the risk of bias. Because of the substantial number of categories for many of our meta-regressions, we were unable to conduct a sensitivity analysis on risk of bias as that would have resulted in underpowered models.

Further research is needed to examine the effectiveness of ABIs in community settings. Our review suggested limited effect but relied on a small number of studies across a wide variety of settings. Our review excluded the use of computer-based interventions, which may be an important approach to reaching populations who do not consider themselves at risk. Some evidence shows that computer-delivered interventions with personalised feedback can effectively reduce alcohol consumption at short-term and long-term follow-up, however the evidence is weaker when comparing direct feedback between face-to-face and computerised feedback.(86) Our findings clearly show the importance of provider in effective delivery of ABIs and it will be important for future research to measure effectiveness of computerised feedback against different providers. Subsequent trials should also comprehensively describe intervention components to enable finer-grained analysis of the relationship between specific aspects of intervention modalities and their effectiveness.

Findings of this review contribute significantly to the understanding of the key processes involved in the delivery of effective ABIs, and have important policy implications for the design of preventative alcohol strategies both in the UK and internationally. The review provides important new evidence on

the effectiveness of brief advice in reducing quantity of alcohol consumed and the role that nurses play in moderating the effectiveness of interventions. Resources should be prioritised to provide further support and training for nurses to deliver ABIs, as well as to undertake research to understand why nurse-led interventions are more effective so appropriate training can be provided to other practitioners.

Data sharing agreement

No additional data are available

Role of Funders

The study was funded by Camden and Islington Public Health, who commented on the study protocol, analysis and interpretation of findings.

Contributors

LP and CA developed the study protocol with advice from EK. LP conducted the search with assistance from DNB. LP and DNB checked the eligibility criteria of all manuscripts with help from AO. AO and JB conducted the data extraction and validation of extraction. GJM developed the statistical approach and conducted all statistical analyses in collaboration with LP. All authors commented on the manuscript.

Acknowledgements

The authors would like to acknowledge Antonio Gasparinni and Colin Muirhead who provided statistical advice on the multi-level meta-analysis and sensitivity check applied in the review.

Licence for Publication

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in JECH and any other BMJPGJL products and sublicences such use and exploit all subsidiary rights, as set out in our licence (<http://group.bmj.com/products/journals/instructions-for-authors/licence-forms>).

Competing Interest: None declared.

List of Figures

Figure 1: Categories and definitions of Interventions by Content, Provider and Setting

Figure 2: Flow chart of systematic review and study selection

Figure 3: Meta-regression analysis on alcohol consumed per unit time at first-follow up by setting of intervention

Figure 4: Meta-regression analysis on alcohol consumed per unit time at first-follow up by provider of intervention

Figures 5: Meta-regression analysis on alcohol consumed per unit time at first-follow up by content of intervention

List of Tables

Table 1: Characteristics of Included studies

Table 2: Summary of study characteristics

Table 3: Summary risk of bias assessment

Table 4 Results of multi-level meta-regression for quantity outcomes

Table 5 Results of multi-level meta-regression for frequency outcomes

Web appendix

Medline search

Online Table 1: Study Level risk of bias assessment

Online table 2: Amount of alcohol per unit time

Online table 3: Amount of alcohol per occasion

Online table 4: Drinking occasions per unit time

Online table 5: Binge drinking occasions per unit time

References

1. World Health Organisation. Global Status Report on Alcohol and Health 2014 Luxembourg: World Health Organisation 2014.
2. Global Burden of Diseases Mortality Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;**385**.
3. Office of National Statistics. Alcohol-related deaths in the United Kingdom, registered in 2012 United Kingdom: Office of National Statistics 2014.
4. National Audit Office. Reducing Alcohol Harm: health services in England for alcohol misuse The Stationery Office, London, United Kingdom: Department of Health 2008.
5. Heather N. Breaking new ground in the study and practice of alcohol brief interventions. *Drug Alcohol Rev* 2010;**29**:584-8.
6. Kaner EF, Beyer F, Dickinson HO, et al. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev* 2007:CD004148.
7. McCambridge J, Kypri K. Can simply answering research questions change behaviour? Systematic review and meta analyses of brief alcohol intervention trials. *PLoS One* 2011;**6**:e23748.
8. McCambridge J, Rollnick S. Should brief interventions in primary care address alcohol problems more strongly? *Addiction* 2014;**109**:1054-8.
9. Alvarez-Bueno C, Rodriguez-Martin B, Garcia-Ortiz L, et al. Effectiveness of brief interventions in primary health care settings to decrease alcohol consumption by adult non-dependent drinkers: a systematic review of systematic reviews. *Prev Med* 2015;**76 Suppl**:S33-8.
10. Poikolainen K. Effectiveness of brief interventions to reduce alcohol intake in primary health care populations: a meta-analysis. *Prev Med* 1999;**28**:503-9.
11. Rubak SLM, Sandbaek A, Lauritzen T, et al. Motivational Interviewing: A systematic review and meta-analysis. *Brit J Gen Pract* 2005;**55**:305-12.
12. Gaume J, McCambridge J, Bertholet N, et al. Mechanisms of action of brief alcohol interventions remain largely unknown - a narrative review. *Front Psychiatry* 2014;**5**:108.
13. Ballesteros J, Gonzalez-Pinto A, Querejeta I, et al. Brief interventions for hazardous drinkers delivered in primary care are equally effective in men and women. *Addiction* 2004;**99**:103-8.
14. Kaner EFS, Dickinson HO, Beyer F, et al. The effectiveness of brief alcohol interventions in primary care settings: A systematic review. *Drug and Alcohol Rev* 2009;**28**:301-23.
15. O'Donnell A, Anderson P, Newbury-Birch D, et al. The impact of brief alcohol interventions in primary healthcare: a systematic review of reviews. *Alcohol Alcohol* 2014;**49**:66-78.
16. Marlatt GA, Baer JS, Kivlahan DR, et al. Screening and brief intervention for high-risk college student drinkers: results from a 2-year follow-up assessment. *J Consult Clin Psychol* 1998;**66**:604-15.
17. Carey KB, Scott-Sheldon LAJ, Carey MP, et al. Individual-level interventions to reduce college student drinking: A meta-analytic review. *Addict Behav* 2007;**32**:2469-94.
18. O'Connor MJ, Whaley SE. Brief intervention for alcohol use by pregnant women. *Am J Pub Health* 2007;**97**:252-8.
19. Elzerbi C, Donoghue K, Drummond C. A comparison of the efficacy of brief interventions to reduce hazardous and harmful alcohol consumption between European and non-European countries: a systematic review and meta-analysis of randomized controlled trials. *Addiction* 2015;**110**:1082-91.
20. Havard A, Shakeshaft A, Sanson-Fisher R. Systematic review and meta-analyses of strategies targeting alcohol problems in emergency departments: interventions reduce alcohol-related injuries. *Addiction* 2008;**103**:368-76; discussion 77-8.
21. Emmen MJ, Schippers GM, Bleijenberg G, et al. Effectiveness of opportunistic brief interventions for problem drinking in a general hospital setting: systematic review. *Bmj* 2004;**328**:318.

22. McQueen J, Howe TE, Allan L, et al. Brief interventions for heavy alcohol users admitted to general hospital wards. *Cochrane Database Syst Rev* 2011;Cd005191.
23. Nilsen P. Brief alcohol intervention--where to from here? Challenges remain for research and practice. *Addiction* 2010;**105**:954-9.
24. Sullivan LE, Tetrault JM, Braithwaite RS, et al. A meta-analysis of the efficacy of nonphysician brief interventions for unhealthy alcohol use: implications for the patient-centered medical home. *Am J Addict* 2011;**20**:343-56.
25. HM Government. The Government's Alcohol Strategy London: 2012.
26. Scottish Government. Chief executive's letter, 2007. Guidance on HEAT Target for NHS Boards Edinburgh: Scottish Government 2007.
27. National Institute for Health and Care Excellence. Alcohol use disorders: preventing harmful drinking London: National Institute for Health and Care Excellence, 2010.
28. World Health Organisation. Global strategy to reduce the harmful use of alcohol. Italy: World Health Organisation 2010.
29. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration 2009 2009-07-21 10:46:49.
30. Higgins JPT, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions. Version 5.1.0: The Cochrane Collaboration; 2011.
31. Van den Noortgate W, López-López J, Marín-Martínez F, et al. Meta-analysis of multiple outcomes: a multilevel approach. *Behav Res* 2014:1-21.
32. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;**36**:1-48.
33. Cheung MW. Modeling dependent effect sizes with three-level meta-analyses: a structural equation modeling approach. *Psychol Methods* 2014;**19**:211-29.
34. Borenstein M, Hedges LV, Higgins JPT, et al. Introduction to Meta-Analysis. Chichester, West Sussex: John Wiley and Sons 2009.
35. R Core Team. A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2016.
36. Fleming MF, Barry KL, Manwell LB, et al. Brief physician advice for problem alcohol drinkers. A randomized controlled trial in community-based primary care practices. *JAMA* 1997;**277**:1039-45.
37. Grossberg PM, Brown DD, Fleming MF. Brief physician advice for high-risk drinking among young adults. *Annals of family medicine* 2004;**2**:474-80.
38. Manwell LB, Fleming MF, Mundt MP, et al. Treatment of problem alcohol use in women of childbearing age: results of a brief intervention trial. *Alcoholism: Clinical & Experimental Research* 2000;**24**:1517-24.
39. Seppä K. Intervention in alcohol abuse among macrocytic patients in general practice. *Scand J Prim Health Care* 1992;**10**:217-22.
40. Liu SI, Wu SI, Chen SC, et al. Randomized controlled trial of a brief intervention for unhealthy alcohol use in hospitalized Taiwanese men. *Addiction* 2011;**106**:928-40.
41. Noknoy S, Rangsin R, Saengcharnchai P, et al. RCT of effectiveness of motivational enhancement therapy delivered by nurses for hazardous drinkers in primary care units in Thailand. *Alcohol Alcohol* 2010;**45**:263-70.
42. Richmond R, Heather N, Wodak A, et al. Controlled evaluation of a general practice-based brief intervention for excessive drinking. *Addiction* 1995;**90**:119-32.
43. Aalto M, Saksanen R, Laine P, et al. Brief intervention for female heavy drinkers in routine general practice: a 3-year randomized, controlled study. *Alcohol Clin Exp Res* 2000;**24**:1680-6.
44. Aalto M, Seppä K, Mattila P, et al. Brief intervention for male heavy drinkers in routine general practice: a three-year randomized controlled study. *Alcohol Alcohol* 2001;**36**:224-30.
45. Anderson P, Scott E. The effect of general practitioners' advice to heavy drinking men. *Br J Addict* 1992;**87**:891-900.

46. Antti-Poika I, Karaharju E, Roine R, et al. Intervention of heavy drinking--a prospective and controlled study of 438 consecutive injured male patients. *Alcohol Alcohol* 1988;**23**:115-21.
47. Baer JS, Kivlahan DR, Blume AW, et al. Brief intervention for heavy-drinking college students: 4-year follow-up and natural history. *Am J Pub Health* 2001;**91**:1310-6.
48. Beich A, Gannik D, Saelan H, et al. Screening and brief intervention targeting risky drinkers in Danish general practice--a pragmatic controlled trial. *Alcohol Alcohol* 2007;**42**:593-603.
49. Bernstein J, Heeren T, Edward E, et al. A brief motivational interview in a pediatric emergency department, plus 10-day telephone follow-up, increases attempts to quit drinking among youth and young adults who screen positive for problematic drinking. *Acad Emerg Med* 2010;**17**:890-902.
50. Butler LH, Correia CJ. Brief Alcohol Intervention With College Student Drinkers: Face-to-Face Versus Computerized Feedback. *Psychol Addict Behav* 2009;**23**:163-7.
51. Carey KB, Carey MP, Maisto SA, et al. Brief motivational interventions for heavy college drinkers: A randomized controlled trial. *J Consult Clin Psychol* 2006;**74**:943-54.
52. Cherpitel CJ, Korcha RA, Moskalewicz J, et al. Screening, brief intervention, and referral to treatment (SBIRT): 12-month outcomes of a randomized controlled clinical trial in a Polish emergency department. *Alcohol Clin Exp Res* 2010;**34**:1922-8.
53. Chick J, Lloyd G, Crombie E. Counselling problem drinkers in medical wards: a controlled study. *Br Med J (Clin Res Ed)* 1985;**290**:965-7.
54. Cordoba R, Delgado M, Pico V, et al. Effectiveness of brief intervention on non-dependent alcohol drinkers (EBIAL): A Spanish multi-centre study. *Fam Pract* 1998;**15**:562-8.
55. Crawford MJ, Patton R, Touquet R, et al. Screening and referral for brief intervention of alcohol-misusing patients in an emergency department: a pragmatic randomised controlled trial. *Lancet* 2004;**364**:1334-9.
56. Crawford MJ, Sanatinia R, Barrett B, et al. The clinical effectiveness and cost-effectiveness of brief intervention for excessive alcohol consumption among people attending sexual health clinics: A randomised controlled trial (SHEAR). *HTA* 2014;**18**.
57. Curry SJ, Ludman EJ, Grothaus LC, et al. A randomized trial of a brief primary-care-based intervention for reducing at-risk drinking practices. *Health Psychol* 2003;**22**:156-65.
58. Daepfen J, Gaume J, Bady P, et al. Brief alcohol intervention and alcohol assessment do not influence alcohol use in injured patients treated in the emergency department: a randomized controlled clinical trial [corrected] [published erratum appears in ADDICTION 2007 Dec;102(12):1995]. *Addiction* 2007;**102**:1224-33.
59. Daepfen JB, Bertholet N, Gaume J, et al. Efficacy of brief motivational intervention in reducing binge drinking in young men: A randomized controlled trial. *Drug Alcohol Depend* 2011;**113**:69-75.
60. Drummond C, Deluca P, Coulton S, et al. The effectiveness of alcohol screening and brief intervention in emergency departments: A multicentre pragmatic cluster randomized controlled trial. *PLoS One* 2014;**9**.
61. Field CA, Caetano R, Harris TR, et al. Ethnic differences in drinking outcomes following a brief alcohol intervention in the trauma care setting. *Addiction (Abingdon, England)* 2010;**105**:62-73.
62. Fleming M, Manwell LB. Brief intervention in primary care settings: a primary treatment method for at-risk, problem, and dependent drinkers. *Alcohol Res* 1999;**23**:128-37.
63. Fleming MF, Balousek SL, Grossberg PM, et al. Brief physician advice for heavy drinking college students: a randomized controlled trial in college health clinics. *J Stud Alcohol Drugs* 2010;**71**:23-31.
64. Freyer-Adam J, Coder B, Baumeister SE, et al. Brief alcohol intervention for general hospital inpatients: a randomized controlled trial. *Drug Alcohol Depend* 2008;**93**:233-43.
65. Gaume J, Gmel G, Faouzi M, et al. Is brief motivational intervention effective in reducing alcohol use among young men voluntarily receiving it? A randomized controlled trial. *Alcohol Clin Exp Res* 2011;**35**:1822-30.

66. Gaume J, Magill M, Longabaugh R, et al. Influence of counselor characteristics and behaviors on the efficacy of a brief motivational intervention for heavy drinking in young men—a randomized controlled trial. *Alcoholism: Clinical and Experimental Research* 2014;**38**:2138-47.
67. Gentilello LM, Rivara FP, Donovan DM, et al. Alcohol interventions in a trauma center as a means of reducing the risk of injury recurrence. *Ann Surg* 1999;**230**:473-80; discussion 80-3.
68. Gottlieb, Becker U, Nielsen AS, et al. Internet-based brief personalized feedback intervention in a non-treatment-seeking population of adult heavy drinkers: a randomized controlled trial. *J Med Internet Res* 2012;**14**:e98-e.
69. Heather N, Campion PD, Neville RG, et al. Evaluation of a controlled drinking minimal intervention for problem drinkers in general practice (the DRAMS scheme). *J R Coll Gen Pract* 1987;**37**:358-63.
70. Holloway AS, Watson HE, Arthur AJ, et al. The effect of brief interventions on alcohol consumption among heavy drinkers in a general hospital setting. *Addiction* 2007;**102**:1762-70.
71. Ingersoll KS, Ceperich SD, Hettema JE, et al. Preconceptional motivational interviewing interventions to reduce alcohol-exposed pregnancy risk. *J Subst Abuse Treat* 2013;**44**:407-16.
72. Juárez P, Walters ST, Daugherty M, et al. A randomized trial of motivational interviewing and feedback with heavy drinking college students. *J Drug Educ* 2006;**36**:233-46.
73. Kulesza M, Apperson M, Larimer ME, et al. Brief alcohol intervention for college drinkers: how brief is? *Addict Behav* 2010;**35**:730-3.
74. Kulesza M, McVay MA, Larimer ME, et al. A randomized clinical trial comparing the efficacy of two active conditions of a brief intervention for heavy college drinkers. *Addict Behav* 2013;**38**:2094-101.
75. Larimer ME, Turner AP, Anderson BK, et al. Evaluating a brief alcohol intervention with fraternities. *J Stud Alcohol* 2001;**62**:370-80.
76. Lock CA, Kaner E, Heather N, et al. Effectiveness of nurse-led brief alcohol intervention: a cluster randomized controlled trial. *J Adv Nurs* 2006;**54**:426-39.
77. Maisto SA, Conigliaro J, McNeil M, et al. Effects of two types of brief intervention and readiness to change on alcohol use in hazardous drinkers. *J Stud Alcohol* 2001;**62**:605-14.
78. Murphy JG, Duchnick JJ, Vuchinich RE, et al. Relative efficacy of a brief motivational intervention for college student drinkers. *Psychol Addict Behav* 2001;**15**:373-9.
79. Rubio G, Jiménez-Arriero MA, Martínez I, et al. Efficacy of physician-delivered brief counseling intervention for binge drinkers. *AJM* 2010;**123**:72-8.
80. Rubio DM, Day NL, Conigliaro J, et al. Brief motivational enhancement intervention to prevent or reduce postpartum alcohol use: a single-blinded, randomized controlled effectiveness trial. *J Subst Abuse Treat* 2014;**46**:382-9.
81. Saitz R, Palfai TP, Cheng DM, et al. Brief intervention for medical inpatients with unhealthy alcohol use: a randomized, controlled trial. [Summary for patients in *Ann Intern Med*. 2007 Feb 6;146(3):122; PMID: 17283343]. *Ann Intern Med* 2007;**146**:167-76.
82. Schaus JF, Sole ML, McCoy TP, et al. Alcohol screening and brief intervention in a college student health center: a randomized controlled trial. (College drinking: new research from the National Institute on Alcohol Abuse and Alcoholism's Rapid Response to College Drinking Problems initiative.). *J Stud Alcohol Drugs* 2009;**70**:131-41.
83. Senft RA, Polen MR, Freeborn DK, et al. Brief intervention in a primary care setting for hazardous drinkers. *Am J Prev Med* 1997;**13**:464-70.
84. Shiles CJ, Canning UP, Kennell-Webb SA, et al. Randomised controlled trial of a brief alcohol intervention in a general hospital setting. *Trials* 2013;**14**.
85. Smith AJ, Hodgson RJ, Bridgeman K, et al. A randomized controlled trial of a brief intervention after alcohol-related facial injury. *Addiction* 2003;**98**:43-52.
86. Wagener TL, Leffingwell TR, Mignogna J, et al. Randomized trial comparing computer-delivered and face-to-face personalized feedback interventions for high-risk drinking among college students. *J Subst Abuse Treat* 2012;**43**:260-7.

87. Walters ST, Vader AM, Harris TR, et al. Dismantling motivational interviewing and feedback for college drinkers: a randomized clinical trial. *J Consult Clin Psychol* 2009;**77**:64-73.
88. Watt K, Shepherd J, Newcombe R. Drunk and dangerous: a randomised controlled trial of alcohol brief intervention for violent offenders. *J Exp Criminol* 2008;**4**:1-19.
89. Saitz R, Svikis D, D'Onofrio G, et al. Challenges Applying Alcohol Brief Intervention in Diverse Practice Settings: Populations, Outcomes, and Costs. *Alcoholism: Clinical and Experimental Research* 2006;**30**:332-8.
90. Brooker C, Peters J, McCabe C, et al. The views of nurses to the conduct of a randomised controlled trial of problem drinkers in an accident and emergency department. *Int J Nurs Stud* 1999;**36**:33-9.
91. Desy PM, Perhats C. Alcohol screening, brief intervention, and referral in the emergency department: an implementation study. *J Emerg Nurs* 2008;**34**:11-9.
92. Rodriguez-Martos A, Castellano Y, Salmeron JM, et al. Simple advice for injured hazardous drinkers: an implementation study. *Alcohol Alcohol* 2007;**42**:430-5.
93. Lock CA. Screening and brief alcohol interventions: what, why, who, where and when? A review of the literature. *J Subst Use* 2004;**9**:91-101.
94. Werch CE, Owen DM, Carlson JM, et al. One-year follow-up results of the STARS for Families alcohol prevention program. *Health Educ Res* 2003;**18**:74-87.
95. Johansson K, Bendtsen P, Akerlind I. Early intervention for problem drinkers: readiness to participate among general practitioners and nurses in Swedish primary health care. *Alcohol Alcohol* 2002;**37**:38-42.
96. Lock CA, Kaner EF. Implementation of brief alcohol interventions by nurses in primary care: do non-clinical factors influence practice? *Fam Pract* 2004;**21**:270-5.
97. Owens L, Gilmore IT, Pirmohamed M. General practice nurses' knowledge of alcohol use and misuse: a questionnaire survey. *Alcohol Alcohol* 2000;**35**:259-62.
98. Kaner E, Bland M, Cassidy P, et al. Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): pragmatic cluster randomised controlled trial. *Bmj* 2013;**346**:e8501.
99. Newbury-Birch D, Coulton S, Bland M, et al. Alcohol screening and brief interventions for offenders in the probation setting (SIPS Trial): a pragmatic multicentre cluster randomized controlled trial. *Alcohol Alcohol* 2014;**49**:540-8.
100. Marsh HW, Bornmann L, Mutz R, et al. Gender Effects in the Peer Reviews of Grant Proposals: A Comprehensive Meta-Analysis Comparing Traditional and Multilevel Approaches. *Rev Educ Res* 2009;**79**:1290-326.