

Full title: The effect of twelve weeks of combined upper- and lower-body high intensity interval training on muscular and cardiorespiratory fitness in older adults

Running head: HIT in older adults

Authors: Christopher Hurst^{1,2}, Kathryn L Weston³ & Matthew Weston²

Affiliations: ¹School of Applied Social Sciences, Durham University, Durham, United Kingdom.

²Department of Psychology, Sport & Exercise, School of Social Sciences, Humanities and Law, Teesside University, Middlesbrough, United Kingdom.

³School of Health and Social Care, Teesside University, Middlesbrough, United Kingdom.

ORCID: Christopher Hurst: 0000-0002-7239-6599
Kathryn L Weston: 0000-0001-5918-6389
Matthew Weston: 0000-0002-9531-3004

Corresponding Author: Christopher Hurst
School of Applied Social Sciences
Durham University
42 Old Elvet
Durham
DH1 3HN
UNITED KINGDOM

Email: christopher.d.hurst@dur.ac.uk

Telephone: +44 (0)191 3341516

Fax: +44 (0)191 3346987

Full title: The effect of twelve weeks of combined upper- and lower-body high intensity interval training on muscular and cardiorespiratory fitness in older adults

Running head: HIT in older adults

Abstract (250-word limit)

Background: High-intensity interval training (HIT) can impact cardiorespiratory and muscular fitness simultaneously, yet protocols typically focus on lower-body exercise. For older adults however, performing activities of daily living requires upper- and lower-body fitness.

Aims: To assess the effects of combined upper- and lower-body HIT on fitness in adults aged >50 years.

Methods: Thirty-six adults (50-81y; 21 male) were assigned via minimisation to either HIT (n=18) or a no-exercise control group (CON, n=18) following baseline assessment of leg extensor muscle power, handgrip strength, cardiorespiratory fitness (predicted VO_{2max}) and health-related quality of life (HRQoL). The HIT group completed two training sessions per week for 12-weeks, performing a combination of upper-, lower- and full-body exercises using a novel hydraulic resistance ergometer. Data were analysed via ANCOVA with probabilistic inferences made about the clinical relevance of observed effects.

Results: All participants completed the intervention with mean ($82 \pm 6\%HR_{max}$) and peak ($89 \pm 6\%HR_{max}$) exercise heart rates confirming a high-intensity training stimulus. Compared with CON, HIT showed possibly small beneficial effects for dominant leg power (10.5%; 90% confidence interval 2.4 to 19.4%), non-dominant leg power (9.4%; 3.3 to 16.0%) and non-dominant handgrip strength (6.3%; 1.2 to 11.5%) while the intervention effect was likely trivial (5.9%; 0.5 to 11.5%) for dominant handgrip strength. There was a likely small beneficial effect for predicted VO_{2max} (8.4%; 1.8 to 15.4%) and small-moderate improvements across several domains of HRQoL.

Conclusion: Combined upper- and lower-body HIT has small clinically relevant beneficial effects on muscular and cardiorespiratory fitness in older adults.

Keywords: High-intensity interval training, Muscular strength, Muscular power, Physical performance, Cardiorespiratory fitness, Ageing

Introduction

Age-associated physiological changes in the neuromuscular and cardiorespiratory systems have important implications for maintaining an independent and healthy life into old age. Reduced muscular fitness is indicative of functional limitation [1] with lower body muscle power output of particular relevance because of its association with the capacity to perform the activities of daily living [2-4]. Moreover, reduced cardiorespiratory fitness (VO_{2max}) is associated with an increased risk of morbidity and mortality [5]. Exercise training offers a potential strategy for counteracting the deleterious effects of ageing with endurance and strength training capable of eliciting substantial fitness improvements in older adults [6, 7]. However, the divergent nature of physiological adaptation induced via endurance training and strength training means that older adults should perform both exercise modalities to maximise potential health and fitness benefits [8]. Despite this, the requirement to perform separate endurance and strength training activities places considerable time demands on individuals – an important consideration as lack of time remains a frequently cited barrier to exercise in a population where adherence to exercise guidelines remains poor [9, 10]. As such, training interventions which can elicit the benefits of endurance and strength training within a single exercise session may be an attractive proposition for potential exercisers.

High-intensity interval training (HIT), characterised by brief, intermittent bursts of vigorous activity interspersed with periods of rest or low-intensity exercise [11] offers an appealing training strategy because of its potential to induce positive adaptations in muscular and cardiorespiratory fitness simultaneously [12]. In older adults, HIT has previously been shown to be an effective strategy to improve cardiorespiratory [13, 14], muscular [15, 16] and functional fitness [17] with potentially greater effects for older and less fit individuals [18].

Despite emerging evidence supporting the effectiveness of HIT [15, 16], previous investigations have typically utilised cycle ergometry or treadmill walking/running as the preferred exercise mode, thereby providing a predominantly lower-body training stimulus. This is likely to be a sub-optimal training approach in this population as older adults need to maintain both upper- and lower-body fitness to perform the basic activities of daily living [1, 19], thereby suggesting a need for alternative exercise modes for performing HIT. Moreover, training modes which focus on lower-body exercise may be unsuitable or prohibitive for older adults with lower-body musculoskeletal or mobility complications. As such, innovative approaches to delivery of HIT are needed to increase accessibility for a greater number of individuals. Our study therefore, sought to evaluate the effects of 12-weeks of combined upper- and lower-body HIT performed using a novel exercise ergometer on measures of muscular and cardiorespiratory fitness as well as health-related quality of life in adults aged over 50 years.

Methods

Sample size estimation

Estimation of sample size for this investigation was performed using leg extensor muscle power as our primary outcome measure because of its importance for maintaining effective physical functioning in older adults {Bean:2010bz}{Foldvari:2000tc}. Sample size was estimated using a custom-made spreadsheet {Hopkins:2006we} by combining the smallest meaningful change in leg extensor power and the within-subject typical error as determined in our previous work {Hurst:2018bv}. Based on this, a sample size of 30 participants (15 per group) was recommended, however we aimed to recruit 36 participants to allow for dropout during the intervention period.

Participants

Initially, 44 potential participants were assessed for eligibility following recruitment via word of mouth and advertisement in local newspapers. Prior to enrolment, all potential participants completed a medical screening questionnaire to identify any medical conditions and current medication that could affect their ability to perform the required exercise training and testing. Those with pre-existing, neuromuscular or skeletal conditions, systemic disease (e.g. diabetes mellitus, cancer, heart disease) or those currently taking medications known to influence fitness or interpretation of the findings (n = 3) or who had engaged in formal and systematic (moderate to high intensity) endurance or strength training within the last year were excluded (n = 1). Following explanation of the study protocol, four participants declined to take part in the investigation. Participants meeting the inclusion criteria (n = 36) were physically active but were not currently, and had not in the previous year, engaged in structured exercise more than twice per week. All individual participants provided written, informed consent to participate in the study (www.clinicaltrials.gov identifier NCT02714088) which conformed to the requirements of The Declaration of Helsinki and was approved by Teesside University Research and Ethics Committee. Figure 1 documents the flow of participants through the study.

Figure 1 about here

Experimental design

Following baseline testing (April 2016), participants were allocated to either HIT (n = 18) or no-exercise control (CON; n = 18) using the minimisation approach. This method aims to ensure balance between groups at baseline across several prognostic factors even when sample sizes are small{Altman:2005hk}. In this investigation, we defined these factors as age, sex, peak power of dominant leg, peak handgrip strength of dominant hand and aerobic fitness. Minimisation was

performed using a custom-made spreadsheet {Hopkins:2010ua} by an investigator not involved in baseline testing (MW) with individual subject data coded to ensure blinded allocation. Participant characteristics at baseline are presented in Table 1. All participants (HIT and CON) were instructed to maintain habitual physical activity and not engage in any additional structured exercise outside of the intervention.

Table 1 about here

HIT intervention

Participants allocated to HIT completed two instructor-led, combined upper- and lower-body HIT sessions per week for 12-weeks (April - August 2016), performed in groups of two to five participants with 72 h recovery between sessions. As the aim of this study was to evaluate the effectiveness of the exercise stimulus, effort was made to promote maximal attendance for all participants. As such, when a participant was unable to attend a session for reasons such as work, family commitments or illness, the session was rescheduled wherever possible. To be included in the final analysis, participants were required to attend a minimum of 90% ($\geq 22/24$) of sessions.

The HIT intervention was performed using a novel double-concentric, hydraulic resistance ergometer (Speedflex, AlphaTech Inc, Nelson, NC). Each session began with a warm up of progressive intensity (~6 min) consisting of a combination of upper- (bent over row, shoulder press), lower- (squat, split squat) and full-body (power clean and press, step and press, pulldown to squat, high pull) exercises (See electronic supplementary material file 1 for images and descriptions of exercises). Participants then completed four sets of high-intensity exercise, with each set consisting of four exercise repetitions. Each set consisted of a combination of four of the exercises previously described with one exercise performed per repetition. In week one, repetition duration was 45-s with each repetition followed by a 15-s transition period allowing participants to move to the next exercise. Each set was followed by 3-min of passive rest. Repetition duration increased by 10 seconds at the end of every third week, with duration being 75-s by week 10. Transition period and rest duration remained constant over the course of the intervention with total exercise duration increasing from 12 to 20 minutes. Each session concluded with a cool down consisting of low intensity, full-body movements (~4 min).

During the intervention exercise order was randomised within- and between-sessions, with participants completing each exercise approximately the same number of times across the 12 weeks. This approach meant that participants could perform the same exercises in multiple consecutive sets, but not in consecutive repetitions, with all exercises not necessarily performed in each session. The aim of each exercise session was to deliver a full-body workout for all participants. As the nature of the Speedflex

machine does not permit the setting of a fixed external resistance, participants were asked to work at 'high-intensity' during exercise bouts, defined as peak heart rate (HR_{peak}) >90% of maximal heart rate ($\%HR_{max}$). Subjects were provided with strong verbal encouragement throughout each HIT session and were instructed to reach target heart rate by increasing the speed of movement during exercises.

Heart rate was monitored continuously during exercise at 5 s intervals using the Polar Team2 system (Polar Electro, Kempele, Finland) with maximal heart rate estimated using the formula: maximal heart rate (HR_{max}) = $207 - 0.7 \times \text{age}$ [21]. Where a participant exceeded this predicted value during training, HR_{max} was amended to the higher observed value [22]. Using the CR100[®] scale {Borg:2001wy} participants provided differential ratings of perceived exertion (RPE) ~10 minutes after the completion of exercise for upper-body muscle exertion (RPE-U), lower-body muscle exertion (RPE-L) and perceived sense of breathlessness in the chest (RPE-B) [24]. Compared with alternative scales (e.g. Borg 6-20 RPE scale[®]), the CR100[®] offers a larger and more finely graded numerical range (0-100 AU [arbitrary units]) with several verbal anchors placed along the scale corresponding to whole numbers (0, Nothing at all; 12, Easy; 22, Moderate; 35, Somewhat hard; 50, Hard; 70, Very hard; 100, Maximal). The use of the CR100[®] scale agrees with modern psychophysical theory {Borg:2001wy} and allows for a more sensitive assessment of perceived exertion while also having the advantage of providing association to a percentage scale {Fanchini:2016bc}. Participants were habituated with the CR100[®] scale and the concept of differential RPE at the study outset.

Outcome measures

At baseline, participants completed three pre-tests over several days (April 2016) with a minimum of 48 hours between sessions and at the same time of day to minimise the impact of circadian variation [26]. To reduce systematic bias and stabilise random variability [25] our primary outcome measures (leg extensor muscle power and handgrip strength) were assessed on three separate occasions and our secondary outcome (predicted VO_{2max}) was assessed twice. Health-related quality of life was evaluated once. Post-intervention (~3-7 days following final training session; August 2016) all outcome measures were assessed once. All testing was performed by the same researchers, strictly adhering to the prescribed standardised testing procedures with individual subject data coded to ensure blinding during data analysis. Participants were asked to avoid strenuous physical activity, alcohol and caffeine for 24h prior to each testing session.

Leg extensor muscle power

Leg power was assessed using the Nottingham leg extensor power rig (Medical Engineering Unit, University of Nottingham, Nottingham, UK), a reliable method for assessing lower body muscular

power [27]. Testing was performed in a randomised, counterbalanced order with participants performing all testing sessions in the same order (e.g., always dominant leg first or always non-dominant leg first based on initial randomisation). Participants completed a standardised warm-up consisting of three leg extensions at increasing submaximal intensity (~50, ~75 and ~90% of self-perceived maximal effort). Following this, ten maximal effort leg extensions, each separated by 30 s of passive rest were performed with participants instructed to extend their leg “as hard and as fast as possible”. This process was then repeated on the second leg. The highest value recorded over the ten leg extensions was taken as peak power output. Typical error (CV) was small on both the dominant (7.4%) and non-dominant legs (5.6%) for males and females combined after baseline test three [28].

Handgrip strength

Handgrip strength was assessed using a digital strain-gauge dynamometer (TKK 5401; Grip-D, Takei Scientific Instruments Co., Ltd., Tokyo, Japan). Testing was performed on both hands with testing order determined as previously described. Participants were instructed to maintain the standard bipedal position, with the arm in complete extension and feet positioned hip width apart. The dynamometer was adjusted appropriate to the individual’s hand size with scores recorded to the nearest 0.1 kg. Following a submaximal practice attempt, participants performed three maximal efforts, with 30 s rest following each attempt. The highest recorded value across the three attempts was used for analysis. Typical error was small on the dominant hand (7.1%) and small on the non-dominant hand (4.6%) after baseline test three.

Cardiorespiratory fitness

Cardiorespiratory fitness (predicted VO_{2max}) was assessed using the Chester Step Test [29], a submaximal, multi-stage step test which has previously been shown to be a reliable tool for the estimation of aerobic capacity [30]. The test consists of five stages, each of 2-minutes duration with a maximum test time of 10 minutes. Stepping cadence began at 15 steps/min (60bpm) and increased by 5 steps/min at the end of every 2-minute stage with step frequency controlled by an electronic metronome (Apple iPad, Apple, California, USA). The test was terminated if the participant reached >80% of predicted HR_{max} before the completion of stage five. Heart rate (Polar RS400, Polar Electro, Kempele, Finland) was recorded throughout each 2-minute stage with maximal heart rate calculated via the equation of Gellish et al. [21]. Following completion of the test, VO_{2max} was estimated using the Chester step test calculator (Assist Creative Resources, Wrexham, United Kingdom). After the second baseline test in this investigation [30], typical error was small (6.2%).

Health-related quality of life

Health-related quality of life (HRQoL) was evaluated using the Short Form-36 health questionnaire (SF-36v2, Optum, Eden Prairie, Minnesota, USA) which contains 36 items used to measure 8 domains of HRQoL [31]. The domains of physical functioning, role limitations due to physical health (role-physical), bodily pain and general health relate to the physical component of HRQoL. Domains of vitality, social functioning, role limitations due to emotional health (role-emotional) and mental health relate to the mental component. The SF-36v2 utilises norm-based scoring (NBS) whereby participant responses to each health domain scale are transformed to a T score with a mean value of 50 and a standard deviation of 10. Higher scores indicate better HRQoL.

Statistical analysis

Prior to all analyses plots of the residuals versus the predicted values revealed no evidence of non-uniformity of error. For training data, the proportion of HIT repetitions that met the pre-specified heart rate criteria for high-intensity was determined, with the median and interquartile range (IQR) for these proportions calculated subsequently. Linear mixed modelling, to allow for fixed (RPE) and random effects (within-participant) was used (SPSS v.23, Armonk, NY: IBM Corp) to examine the difference between differential RPE scores and to determine within-subject variability (expressed as a standard deviation [SD]) in RPE-U, RPE-L and RPE-B) with the SD doubled to interpret its magnitude [32].

Outcome measures were log transformed and then back transformed to obtain the percent difference between baseline and post-intervention, with uncertainty of the estimates expressed as 90% confidence intervals (CI). An analysis of covariance (ANCOVA) model was used (SPSS v.23, Armonk, NY: IBM Corp) to compare change scores between the two groups {Vickers:2001io} with age, sex and baseline value of the outcome measure included as covariates to control for any imbalances between the groups at baseline even after minimisation [34]. Adjusted mean intervention effects were evaluated for their practical/clinical significance by pre-specifying thresholds for small, moderate and large effects {Batterham:2006ts}. As robust clinical anchors for our outcome measures remain to be determined in this population, magnitude of effects were defined as standardised mean differences of 0.2, 0.6 and 1.2 between-subject standard deviations (SD) for small, moderate and large effects, respectively {Hopkins:2009dka}. The SD of the pooled baseline values was used for this purpose, as the post-intervention SD can be inflated by individual differences in response to the intervention. Using the mean intervention effect for each outcome, together with its uncertainty (i.e. the confidence interval), the probability of the true effect being trivial, beneficial or harmful was calculated then interpreted using the following scale: <0.5%, most unlikely or almost certainly not; 0.5–5%, very unlikely; 5–25%, unlikely or probably not; 25–75 %, possibly; 75–95%, likely; 95–99.5%, very likely; >99.5 %, most likely [35]. Effects were evaluated clinically given that interventions can be potentially harmful as well

as beneficial to individuals. The default probabilities for declaring an effect clinically beneficial are <0.5% (most unlikely) for harmful and >25% (possibly) for benefit; a clinically unclear effect is therefore possibly beneficial (>25%) with an unacceptable risk of harm (>0.5%) [35]. Data are presented as mean \pm standard deviation (SD) or, for adjusted mean change, 90% confidence interval (CI).

Results

Exercise training attendance and intensity

All 18 participants completed the HIT intervention with an overall attendance of 99% (429 out of a possible 432 sessions) across the 12-week period. There were 42 individual sessions rearranged (~10%) throughout the intervention period to offset participant unavailability and maximise attendance. Sixteen participants completed all twenty-four HIT sessions, one participant completed twenty-three sessions and one participant completed twenty-two. The reasons for missed sessions were: 1) injury unrelated to the study (two sessions missed) and 2) family commitments (one session missed). It was not possible to rearrange these three sessions because of a lack of time during the intervention period. Exercise intensity data are presented in Table 2. The proportion of repetitions meeting the high-intensity criterion (Peak heart rate [HR_{peak}] $\geq 90\%$ HR_{max}) was 62% (IQR 24%, 81%). There were most likely small differences between RPE-U and RPE-L (6 Arbitrary units [AU]; $\pm 90\%$ confidence limits 1 AU) and RPE-L and RPE-B (6 AU ± 1 AU). The difference between RPE-U and RPE-B was most likely trivial (0 AU; ± 1 AU). The magnitude of the within-subject variability for all RPE measures was moderate. No adverse events were reported during any of the exercise testing or training sessions.

Table 2 about here

Outcome measures

Baseline, adjusted mean change values and between-group comparisons (HIT v CON) are presented in Table 3. Compared to CON, HIT showed possibly small beneficial effects for dominant leg power, non-dominant leg power and non-dominant handgrip strength. For dominant handgrip strength, the effect was likely trivial. There was a likely small beneficial effect for predicted VO_{2max} .

Table 3 about here

For health-related quality of life (Table 4), there were possibly small beneficial effects for role-physical, general health, vitality and mental health and a likely small beneficial effect for bodily pain in the HIT group compared with CON. There was a possibly moderate beneficial effect for role-emotional. Between-group differences in physical functioning and social functioning were likely and possibly trivial, respectively.

Table 4 about here

Discussion

The present investigation has evaluated the effects of 12-weeks of combined upper- and lower-body HIT on muscular and cardiorespiratory fitness in older adults observing possibly beneficial improvements in leg power and handgrip strength as well as a likely beneficial improvement in cardiorespiratory fitness. These data provide support for HIT as a multicomponent training strategy in older adults.

Notwithstanding any potential training induced physiological and performance improvements, exercise training interventions must be acceptable and safe for participants. Despite there being no adverse events recorded during the training programme, the limited sample size and training programme duration mean that considerably more data is needed to fully quantify the risks associated with this training approach in older adults. Evaluation of attendance and adherence (i.e. meeting the prescribed exercise intensity [36]) can provide a useful quantification of the feasibility of an intervention. The attendance for the 18 participants across the 12-week intervention reported in this study is considerably higher than the 58% to 77% range for older adults undertaking exercise programmes reported in the systematic review of Picorelli and colleagues [37]. It is acknowledged however, that the structure of this training intervention – with extensive availability of the instructor and the exercise facility – meant a flexible approach to exercise session timing was possible to promote and achieve maximal possible attendance for all participants. Therefore, the present data represent a ‘best case’ scenario which may be considerably different from a ‘real-world’ scenario. Importantly, however, these data do provide support for the feasibility of instructor-led group-based HIT as a method for engaging older adults if a flexible approach to session delivery is possible.

As well as attendance, assessing if participants have performed the exercise training as intended is essential when evaluating interventions [36]. Using our pre-specified target threshold for high-intensity, 62% of repetitions met the criteria. Previous evaluation of intervention fidelity has suggested that high-intensity criterion attainment in 58% of HIT repetitions represented ‘moderate’ intervention fidelity [36], while Weston et al. [24] reported ‘low’ fidelity with ~23% compliance to HIT criteria. Based on

these previous interpretations, it seems reasonable to categorise intervention fidelity as ‘moderate’ in this study. Despite this however, between-study comparisons of intervention fidelity remain limited by a lack of available data. Wider translation of this approach to evaluating the fidelity of exercise training interventions may be enhanced through the use of clear thresholds and accompanying qualitative descriptors. As such, it is recommended that as per Taylor et al. [36] authors report the median along with the interquartile range of sessions/bouts meeting pre-specified criteria and qualify this using our proposed fidelity thresholds and corresponding qualitative descriptors (Median number of sessions/bouts meeting prespecified criteria: <50%, Low; 50-70%, Moderate; >70%, High). The thresholds suggested here represent the authors’ personal experiences of exercise training prescription and evaluation. Further investigation is needed to appraise these qualitative descriptors and their corresponding thresholds.

Mean RPE scores in this investigation fell in the range of ‘somewhat hard’ to ‘hard’, lower than previous studies prescribing HIT intensity as ‘hard’ to ‘very hard’ [24, 38] but similar to those reported following HIT in overweight young adults [39]. Between-study differences are likely multifactorial in nature with heterogeneous study populations as well as differences in exercise programming factors contributing to these observed differences [40]. Despite lower perceived exertion responses than typically prescribed during HIT, the present investigation has reported a clear beneficial effect on predicted $\text{VO}_{2\text{max}}$ – a finding with clear practical implications as exercise that is perceived to be less intense may be more palatable to potential exercisers [39].

Possibly small beneficial improvements in leg power on both the dominant and non-dominant leg were observed in this investigation. This finding has clear practical implications as improvements in leg power have been shown to make an important contribution to clinically meaningful improvements in gait speed and are associated with improved functional performance [2-4]. Previous investigations have shown potential for HIT to increase muscular power [15, 41]; however, these studies have typically employed training protocols classified as sprint-interval training (SIT) or performed at ‘all out’ intensity which may not be suitable or appealing to all. The present findings demonstrate that HIT does not need to be performed ‘all out’, with submaximal HIT capable of inducing improvements in muscular power. This finding may be of particular relevance in this population, where adherence to physical activity guidelines is poor {Jefferis:2014jz}, as previous work has suggested that exercise that is too strenuous is likely to be a deterrent to participation {Hardcastle:2014es}. The observed improvements in muscular power are of a similar magnitude (~10%) to those previously reported in young [41, 42] and older adults [16]. However, these studies assessed peak power using either the Wingate, or a maximal incremental test and as Sculthorpe and colleagues [15] have noted, the mechanical and metabolic differences between single expressions of power (e.g. during a leg extension) versus high-intensity cycling mean that between-study comparisons are limited. Our study is the first to demonstrate that submaximal HIT

increases explosive lower body muscle power in older adults. Further work is needed to understand the mechanistic basis of this adaptation.

Previous investigation has shown handgrip strength to be a strong predictor of mortality [43], while a reduction in grip strength is also related to difficulty in performing activities of daily living [1]. The present data indicate a possibly beneficial improvement in handgrip strength on the non-dominant hand of a similar magnitude to Pereira and colleagues [44] who reported improvements of 5% (dominant hand) and 6.9% (non-dominant hand) after 12-weeks of high-speed power training. Interestingly, both the current study and Pereira et al.[44] have shown greater improvements in handgrip strength on the non-dominant hand. This may be related to lower levels of baseline strength on the non-dominant hand as improvements in strength are related to initial strength level with greater increases reported in participants with lower baseline levels [45]. Our findings demonstrate that HIT could be an effective approach to increase handgrip strength when exercises are prescribed to target upper-body muscle groups. This supports the idea that creative approaches to exercise prescription are needed to maximise the potential for functionally relevant training induced adaptations.

The robust relationship between greater cardiorespiratory fitness and reduced mortality and morbidity means that the likely small beneficial improvement in predicted VO_{2max} reported in this investigation is an important and clinically relevant finding [5]. The observed improvement in VO_{2max} falls within the range of previous studies investigating HIT in older adults, employing the archetypal 4 x 4 min protocol of 6-15% [13, 14, 16]. It should be noted however, that the study reporting the largest improvement (15%; [13]) involved a 10-week training programme with three training sessions per week equating to 25% more training sessions performed than in the current investigation. As lack of time remains a commonly cited barrier to exercise participation {CohenMansfield:2003co}, training programmes which require a reduced time commitment, by prescribing a lower number of weekly sessions, may be more appealing to potential exercisers. Caution is also warranted when making between-study comparisons as improvements in VO_{2max} are affected by baseline fitness, with greater benefit observed for less fit participants [18]. Although speculative, it is possible that both central and peripheral adaptations contribute to increased cardiorespiratory fitness following HIT as reported in this investigation [46].

The present data suggest that HIT has potential to improve perceptions of health-related quality of life in older adults – an important aim for health interventions involving this population group [47]. Although previous work has suggested that HIT may have a beneficial effect on health-related quality of life in healthy [48] and clinical populations [49], comparison of our findings with previous investigations is challenging because of differences in methods of assessment and heterogeneous study populations. For example, the participants in the present investigation exhibited baseline values

considerably higher than the mean across seven of the eight health domains. It is likely that participants with higher perception of health-related quality of life at baseline will need to demonstrate improvements of a greater magnitude, compared with those reporting lower values at baseline, to have a meaningful effect. However, there remains no consensus on what represents the minimal clinically relevant change in health-related quality of life for healthy older adults and further work, involving longer duration follow-up should attempt to elucidate this. Caution is also warranted when interpreting our findings as short-term improvements in health-related quality of life may not translate into long-term changes.

Although this study has demonstrated that 12-weeks of HIT can elicit improvements in muscular power and strength as well as cardiorespiratory fitness, it is not without limitation. Firstly, this is a pragmatic, exploratory trial involving only a small number of participants. It is acknowledged that studies with low statistical power involving small sample sizes likely overestimate or exaggerate the magnitude of an effect [50]. A further definitive trial involving a greater number of participants is needed to accurately quantify training induced changes using this exercise mode. Moreover, a larger sample size would also allow for exploration of potential moderator variables (e.g. sex) which may influence the observed training response. Secondly, the outcome measures used in this investigation provide an understanding of the effects of HIT on isolated components of physical fitness (e.g. lower-body muscle power, upper-body muscle strength, cardiorespiratory fitness), yet performance on functional fitness tests (e.g. 30-s chair stand test, timed up and go) may be more relevant for assessing intervention effects as these tests provide composite measures of physical performance [51]. In addition, submaximal estimation of VO_{2max} is acknowledged as being an alternative to the gold standard approach for assessing cardiorespiratory fitness. However, the Chester Step test does quantify cardiorespiratory fitness in terms of VO_{2max} , facilitating comparison with previous investigations. Although laboratory-based determination of VO_{2max} via incremental exercise testing is the benchmark for assessing cardiorespiratory fitness, the pragmatic nature and logistical constraints of this investigation meant this was not possible. A future definitive trial should include measures of functional fitness and laboratory-based determination of VO_{2max} , while an increased follow-up period may help to understand the longer-term effects of this training approach. Finally, the results of the present investigation are representative of a healthy and relatively young older adult population and therefore should not be extrapolated to represent all older adults, including those with multimorbidity as well as individuals engaged in drug therapy. Consequently, further work is needed to evaluate the potential effectiveness of this training approach with these population groups.

Conclusions

Our study has demonstrated that 12-weeks of combined upper- and lower-body HIT is an effective method for inducing clinically meaningful improvements in aspects of muscular and cardiorespiratory fitness in adults aged over 50 years. In particular, the observed improvements in lower body muscle power are especially relevant because of the role this plays in maintaining functional performance in this population. Moreover, the present data provide evidence that HIT does not need to be performed at an ‘all out’ intensity to induce fitness improvements in this population. More broadly, this study provides further indication that HIT is a feasible and effective approach to exercise training in older adults with high intervention fidelity reported. These findings highlight potential for innovative approaches to training delivery and should encourage researchers to move beyond exercise modes traditionally associated with HIT.

Acknowledgements

The authors would like to thank the participants for their involvement in this study. We also acknowledge Tom W Macpherson and Shaun J McLaren for their assistance with data collection.

Compliance with ethical standards

Conflict of interest

This study was completed as part of Christopher Hurst's doctoral degree sponsored by Speedflex (UK & Ireland) Ltd. The funders had no involvement in study design, data collection and analysis, decision to publish or preparation of the manuscript. All other authors have no conflicts of interest to declare.

Ethical approval

All procedures performed within this study were in accordance with the ethical standards of the institutional research ethics committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent

Informed consent was obtained from all individual participants included in the study.

Data availability

The datasets generated and analysed during the current study are available from the corresponding author on reasonable request.

References

1. Hairi NN, Cumming RG, Naganathan V, et al (2010) Loss of Muscle Strength, Mass (Sarcopenia), and Quality (Specific Force) and Its Relationship with Functional Limitation and Physical Disability: The Concord Health and Ageing in Men Project. *J Am Geriatr Soc* 58:2055–2062. doi: 10.1111/j.1532-5415.2010.03145.x
2. Reid KF, Fielding RA (2012) Skeletal muscle power: a critical determinant of physical functioning in older adults. *Exerc Sport Sci Rev* 40:4–12. doi: 10.1097/JES.0b013e31823b5f13
3. Bean JF, Kiely DK, LaRose S, et al (2010) Are Changes in Leg Power Responsible for Clinically Meaningful Improvements in Mobility in Older Adults? *J Am Geriatr Soc* 58:2363–2368. doi: 10.1111/j.1532-5415.2010.03155.x
4. Foldvari M, Clark M, Laviolette LC, et al (2000) Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A Biol Sci Med Sci* 55:M192–199. doi:10.1093/gerona/55.4.m192
5. Kodama S, Saito K, Tanaka S, et al (2009) Cardiorespiratory Fitness as a Quantitative Predictor of All-Cause Mortality and Cardiovascular Events in Healthy Men and Women: A Meta-analysis. *JAMA* 301:2024–2035. doi: 10.1001/jama.2009.681
6. Liu CJ, Latham NK (2009) Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev*. doi: 10.1002/14651858
7. Huang G, Gibson CA, Tran ZV, Osness WH (2005) Controlled endurance exercise training and VO₂max changes in older adults: a meta-analysis. *Preventive cardiology* 8:217–225. doi: 10.1097%2F00005768-200405001-00676
8. Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, et al (2009) Exercise and Physical Activity for Older Adults. *Medicine & Science in Sports & Exercise* 41:1510–1530. doi: 10.1249/MSS.0b013e3181a0c95c
9. Jefferis BJ, Sartini C, Lee I-M, et al (2014) Adherence to physical activity guidelines in older adults, using objectively measured physical activity in a population-based study. *BMC Public Health* 2014 14:1 14:382. doi: 10.1186/1471-2458-14-382
10. Cohen-Mansfield J, Marx MS, Guralnik JM (2003) Motivators and barriers to exercise in an older community-dwelling population. *Journal of Aging and Physical Activity* 11:242–253. doi: 10.1123/japa.11.2.242

11. Gibala MJ, Little JP, MacDonald MJ, Hawley JA (2012) Physiological adaptations to low-volume, high-intensity interval training in health and disease. *The Journal of Physiology* 590:1077–1084. doi: 10.1113/jphysiol.2011.224725
12. Osawa Y, Tabata S, Katsukawa F, et al (2014) Effects of 16-week high-intensity interval training using upper and lower body ergometers on aerobic fitness and morphological changes in healthy men: a preliminary study. *OAJSM* 257. doi: 10.2147/OAJSM.S68932
13. Osteras H, Hoff J, Helgerud J (2005) Effects of high-intensity endurance training on maximal oxygen consumption in healthy elderly people. *Journal of Applied Gerontology* 24:377–387. doi: 10.1177/0733464804273185
14. Hwang C-L, Yoo J-K, Kim H-K, et al (2016) Novel all-extremity high-intensity interval training improves aerobic fitness, cardiac function and insulin resistance in healthy older adults. *Experimental Gerontology* 82:112–119. doi: 10.1016/j.exger.2016.06.009
15. Sculthorpe NF, Herbert P, Grace F (2017) One session of high-intensity interval training (HIIT) every 5 days, improves muscle power but not static balance in lifelong sedentary ageing men. *Medicine* 96:e6040–8. doi: 10.1097/MD.0000000000006040
16. Wang E, Næss MS, Hoff J, et al (2014) Exercise-training-induced changes in metabolic capacity with age: the role of central cardiovascular plasticity. *AGE* 36:665–676. doi: 10.1007/s11357-013-9596-x
17. Adamson SB, Lorimer R, Cobley JN, Babraj JA (2014) Extremely Short-Duration High-Intensity Training Substantially Improves the Physical Function and Self-Reported Health Status of Elderly Adults. *J Am Geriatr Soc* 62:1380–1381. doi: 10.1111/jgs.12916
18. Milanovic Z, Sporiš G, Weston M (2015) Effectiveness of High-Intensity Interval Training (HIT) and Continuous Endurance Training for VO₂max Improvements: A Systematic Review and Meta-Analysis of Controlled Trials. *Sports Med* 45:1469–1481. doi: 10.1007/s40279-015-0365-0
19. Lundgren-Lindquist B, Sperling L (1983) Functional studies in 79-year-olds. II. Upper extremity function. *Scandinavian Journal of Rehabilitation Medicine* 15:117–123.
20. Hopkins WG (2010) Assigning subjects to groups in a controlled trial. *Sportscience* 14:7–12.
21. Gellish RL, Goslin BR, Olson RE, et al (2007) Longitudinal modeling of the relationship between age and maximal heart rate. *Medicine & Science in Sports & Exercise* 39:822–829. doi: 10.1097/mss.0b013e31803349c6

22. Weston M, Helsen W, MacMahon C, Kirkendall D (2004) The impact of specific high-intensity training sessions on football referees' fitness levels. *The American journal of Sports Medicine* 32:54S–61S. doi: 10.1177/0363546503261421
23. Borg E, Borg G (2002) A comparison of AME and CR100 for scaling perceived exertion. *Acta psychologica* 109:157–175. doi:10.1016/s0001-6918(01)00055-5
24. Weston M, Batterham AM, Tew GA, et al (2017) Patients Awaiting Surgical Repair for Large Abdominal Aortic Aneurysms Can Exercise at Moderate to Hard Intensities with a Low Risk of Adverse Events. *Front Physiol* 7:1067–11. doi: 10.3389/fphys.2016.00684
25. Hopkins WG (2000) Measures of reliability in sports medicine and science. *Sports Med* 30:1–15.
26. Atkinson G, Reilly T (1996) Circadian variation in sports performance. *Sports Med* 21:292–312. doi: 10.2165/00007256-199621040-00005
27. Hurst C, Batterham AM, Weston KL, Weston M (2018) Short- and long-term reliability of leg extensor power measurement in middle-aged and older adults. *Journal of Sports Sciences* 36:970–977. doi: 10.1080/02640414.2017.1346820
28. Hopkins WG (2015) Spreadsheets for Analysis of Validity and Reliability. *Sportscience* 19:36–44.
29. Sykes K, Roberts A (2004) The Chester step test—a simple yet effective tool for the prediction of aerobic capacity. *Physiotherapy* 90:183–188. doi: 10.1016/j.physio.2004.03.008
30. Buckley JP, Sim J, Eston RG, et al (2004) Reliability and validity of measures taken during the Chester step test to predict aerobic power and to prescribe aerobic exercise. *British Journal of Sports Medicine* 38:197–205. doi: 10.1136/bjism.2003.005389
31. Ware JE, Sherbourne CD (1992) The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 30:473–483. doi: 10.1097/00005650-199206000-00002
32. Smith TB, Hopkins WG (2011) Variability and Predictability of Finals Times of Elite Rowers. *Med Sci Sport Exer* 43:2155–2160. doi: 10.1249/MSS.0b013e31821d3f8e
33. Atkinson G, Batterham AM (2015) True and false interindividual differences in the physiological response to an intervention. *Exp Physiol* 100:577–588. doi: 10.1113/EP085070
34. Senn S (2006) Change from baseline and analysis of covariance revisited. *Stat Med* 25:4334–4344. doi: 10.1002/sim.2682

35. Hopkins WG, Marshall SW, Batterham AM, Hanin J (2009) Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sport Exer* 41:3–13. doi: 10.1249/MSS.0b013e31818cb278
36. Taylor KL, Weston M, Batterham AM (2015) Evaluating Intervention Fidelity: An Example from a High-Intensity Interval Training Study. *PLoS ONE* 10:e0125166–8. doi: 10.1371/journal.pone.0125166
37. Picorelli AMA, Pereira LSM, Pereira DS, et al (2014) Adherence to exercise programs for older people is influenced by program characteristics and personal factors: a systematic review. *Journal of Physiotherapy* 60:151–156. doi: 10.1016/j.jphys.2014.06.012
38. Cassidy S, Thoma C, Hallsworth K, et al (2016) High intensity intermittent exercise improves cardiac structure and function and reduces liver fat in patients with type 2 diabetes: a randomised controlled trial. *Diabetologia* 59:56–66. doi: 10.1007/s00125-015-3741-2
39. Kilpatrick MW, Martinez N, Little JP, et al (2015) Impact of high-intensity interval duration on perceived exertion. *Med Sci Sport Exer* 47:1038–1045. doi: 10.1249/MSS.0000000000000495
40. McLaren SJ, Graham M, Spears IR, Weston M (2016) The Sensitivity of Differential Ratings of Perceived Exertion as Measures of Internal Load. *Int J Sports Physiol Perform* 11:404–406. doi: 10.1123/ijsp.2015-0223
41. Astorino TA, Allen RP, Roberson DW, et al (2011) Adaptations to high-intensity training are independent of gender. *Eur J Appl Physiol* 111:1279–1286. doi: 10.1007/s00421-010-1741-y
42. Astorino TA, Allen RP, Roberson DW, Jurancich M (2012) Effect of high-intensity interval training on cardiovascular function, VO₂max, and muscular force. *J Strength Cond Res* 26:138–145. doi: 10.1519/JSC.0b013e318218dd77
43. Leong DP, Teo KK, Rangarajan S, et al (2015) Prognostic value of grip strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. *The Lancet* 386:266–273. doi: 10.1016/S0140-6736(14)62000-6
44. Pereira A, Izquierdo M, Silva AJ, et al (2012) Effects of high-speed power training on functional capacity and muscle performance in older women. *Exp Gerontol* 47:250–255. doi: 10.1016/j.exger.2011.12.010
45. Lexell J (2000) Strength training and muscle hypertrophy in older men and women. *Topics in Geriatric Rehabilitation* 15:41–46. doi:10.1097/00013614-200003000-00007
46. Daussin FN, Zoll J, Dufour SP, et al (2008) Effect of interval versus continuous training on cardiorespiratory and mitochondrial functions: relationship to aerobic performance

- improvements in sedentary subjects. *AJP: Regulatory, Integrative and Comparative Physiology* 295:R264–R272. doi: 10.1152/ajpregu.00875.2007
47. Rejeski WJ, Mihalko SL (2001) Physical Activity and Quality of Life in Older Adults. *J Gerontol A Biol Sci Med Sci* 56:23–35. doi: 10.1093/gerona/56.suppl_2.23
 48. Knowles A-M, Herbert P, Easton C, et al (2015) Impact of low-volume, high-intensity interval training on maximal aerobic capacity, health-related quality of life and motivation to exercise in ageing men. *AGE* 37:25. doi: 10.1007/s11357-015-9763-3
 49. Wisloff U, Stoylen A, Loennechen JP, et al (2007) Superior Cardiovascular Effect of Aerobic Interval Training Versus Moderate Continuous Training in Heart Failure Patients: A Randomized Study. *Circulation* 115:3086–3094. doi: 10.1161/CIRCULATIONAHA.106.675041
 50. Button KS, Ioannidis JPA, Mokrysz C, et al (2013) Power failure: why small sample size undermines the reliability of neuroscience. *Nature Reviews* 14:365–376. doi: 10.1038/nrn3475
 51. Lord SR, Murray SM, Chapman K, et al (2002) Sit-to-stand performance depends on sensation, speed, balance, and psychological status in addition to strength in older people. *J Gerontol A Biol Sci Med Sci* 57:M539–43.

FIGURE LEGENDS

FIGURE 1 Flow of participants through the study