

[Click here to view linked References](#)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 **Mosquito larvicidal and pupicidal action of biosurfactant produced by *Bacillus subtilis* A1 and *Pseudomonas***  
2 ***stutzeri* NA3 against *Anopheles stephensi***

3

4 Punniyakotti Parthipan<sup>1</sup> · Raja Kumaresan Sarankumar<sup>1</sup> · Anitha Jaganathan<sup>2</sup> · Pandian Amuthavalli<sup>2</sup> · Ranganathan  
5 Babujanarthanam<sup>3</sup> · Pattanathu K.S.M. Rahman<sup>4</sup> · Kadarkarai Murugan<sup>2,5</sup> · Akon Higuchi<sup>6</sup> · Aruliah Rajasekar<sup>1\*</sup>

6

7 <sup>1</sup>Environmental Molecular Microbiology Research Laboratory, Department of Biotechnology, Thiruvalluvar  
8 University, Serkkadu, Vellore, Tamilnadu, India- 632 115.

9 <sup>2</sup>Division of Entomology, Department of Zoology, School of Life Sciences, Bharathiar University, Coimbatore,  
10 Tamilnadu, India-641046.

11 <sup>3</sup>Energy Bioscience Laboratory, Department of Biotechnology, Thiruvalluvar University, Serkkadu, Vellore,  
12 Tamilnadu, India- 632 115.

13 <sup>4</sup>Chemical and Bioprocess Engineering Group, School of Science and Technology, University of Teesside,  
14 Middlesbrough TS13BA Tees Valley, UK.

15 <sup>5</sup>Thiruvalluvar University, Serkkadu, Vellore, Tamilnadu, India- 632 115.

16 <sup>6</sup>Department of Chemical and Materials Engineering, National Central University, No. 300, Jhongli, Taoyuan  
17 32001, Taiwan

18

19 **\*Rajasekar Aruliah**

20 [rajasekargood@gmail.com](mailto:rajasekargood@gmail.com), [rajasekar.aruliah@gmail.com](mailto:rajasekar.aruliah@gmail.com)

21

22

23

24

25

26

27

28

1  
2  
3  
4 **29 Abstract**

5  
6 **30** *Anopheles stephensi* vector of a *Plasmodium parasite* which are responsible for spreading malaria in modern world.  
7  
8 **31** Presently, malaria suppression is desirable one because of insecticide resistance nature of vector, specifically  
9  
10 **32** improvement of *Plasmodium* species as highly resistant to a many antimalarial drugs. Present study focused as  
11  
12 **33** biosurfactant production using two potential biosurfactant producing bacterial strains such as *Bacillus subtilis* A1  
13  
14 **34** and *Pseudomonas stutzeri* NA3 for mosquitocidal application. Produced biosurfactant were characterized using  
15  
16 **35** fourier transform infrared (FTIR) spectroscopy and gas chromatography and mass spectrometry (GCMS) and  
17  
18 **36** confirmed that the produced biosurfactant were lipopeptide in nature. Different concentration of the biosurfactant  
19  
20 **37** ranged between 2-10 ppm was tested against different stages of *A. stephensi* and both biosurfactant were found as  
21  
22 **38** exterminating larval and pupal stages of mosquitoes. LC50 values were 3.58 (I), 4.92 (II), 5.73 (III), 7.10 (IV) and  
23  
24 **39** 7.99 (pupae) and 2.61 (I), 3.68 (II), 4.48 (III), 5.55 (IV) and 6.99 (pupa) for biosurfactant produced by *B. subtilis* A1  
25  
26 **40** and *P. stutzeri* NA3 respectively. Biosurfactant are eco-friendly and easily producible using low cost material. The  
27  
28 **41** toxic nature of these biosurfactant to the targeted organism like *A. stephensi* is lead to promising application in the  
29  
30 **42** medical field.  
31

32  
33 **43**  
34 **44** **Keywords:** Biosurfactant · *Anopheles stephensi* · mosquitocidal · Gas Chromatography · *Bacillus subtilis* ·  
35  
36 **45** Lipopeptide  
37

38  
39 **46**  
40  
41 **47 Introduction**

42 **48**  
43  
44 **49** Mosquitoes signify as an important menace for millions of animals and humans around the world, because they are  
45  
46 **50** active as vectors for dangerous pathogens and parasites, also responsible for millions of fatality per yearly.  
47  
48 **51** *Anopheles stephensi* vector of a *Plasmodium parasite* which are responsible for causing/spreading malaria. In 2013,  
49  
50 **52** about 198 million cases of malaria were recorded and 584,000 deaths were estimated among them. After 2000,  
51  
52 **53** malaria mortality rates have fallen by 47 % worldwide since specifically 54 % in the African region. Most fatality  
53  
54 **54** was recorded among African children, where a child dies was counted every minute due to malaria. A malaria  
55  
56 **55** mortality rate amongst brood in Africa has been abridged by an anticipated 58 % since 2000 (Mehlhorn, 2008).  
57  
58  
59  
60  
61

1  
2  
3  
4 56 However, the regenerations of malaria after suppression in many countries were still documented (Benelli et al.  
5  
6 57 2015a,b).

7  
8 58 Presently, malaria eradication is needed one due to insecticide resistance in vector, along with improvement  
9  
10 59 of *Plasmodium* species as resistant to a many antimalarial drugs. The most proficient way to managing of the vector  
11  
12 60 could be achieved at undeveloped stage of their life cycle (Mahesh Kumar et al. 2012). Herein position, eco-friendly  
13  
14 61 efficient controlling tools were urgently needed (Benelli et al. 2015a,b). Plant derivatives were used by many  
15  
16 62 researchers to control the vector growth, such as *Polygonum hydropiper* L., *Origanum scabrum*, *Clausena anisata*  
17  
18 63 and etc., (Maheswaran and Ignacimuthu, 2014; Govindarajan et al. 2016; Mukandiwa et al. 2016). Recently some  
19  
20 64 researchers used silver nanoparticles synthesized using plant material to control the mosquitoes (Poopathi et al.  
21  
22 65 2014; Murugan et al. 2015; Subramaniam et al. 2015; Subramaniam et al. 2016; Murugan et al. 2016). Ultimately,  
23  
24 66 bio-control of vectors is a suitable hopeful substitute to synthetic chemical pesticides. In this esteem, plentiful  
25  
26 67 biological materials have been tested to evaluate their probable to manage the mosquitos (Knight et al. 2003).  
27  
28 68 Toxins from bacterial strains *Bacillus sphaericus* and *B. thuringensis var.israelensis* were revealed to be useful  
29  
30 69 against mosquito larvae at very low dosage and harmless to non-targeted organisms (Das and Mukherjee, 2006).  
31  
32 70 Nonetheless, the biolarvicide product extracted from *B. sphaericus* strain is noted to be lesser effective against  
33  
34 71 *Anopheles culicifacies* and barely competent against *Aedes aegypti* (Mittal, 2003).

35  
36 72 Some of bacterial strains and their metabolites were used to control mosquitos, such as *Bacillus subtilis*  
37  
38 73 (Das and Mukherjee, 2006; Geetha et al. 2010) and *Bacillus circulans* (Darriet and Hougard, 2002). Rhamnolipid a  
39  
40 74 biosurfactant produced by *Pseudomonas aeruginosa* was potentially used to control the *Aedes aegypti* (Silva et al.  
41  
42 75 2015). Another biosurfactant ‘Di-rhamnolipid’ produced by bacterium *Pseudomonas fluorescens* was active against  
43  
44 76 the pupae of *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus* (Prabakaran et al. 2015). However,  
45  
46 77 many of these activities by bacterial strains and their products have not been completely studied for the  
47  
48 78 characteristic of their bio-control potential. *Bacillus subtilis* synthesis a wide range of biologically active compounds  
49  
50 79 such as fatty acids, lipopeptide which has immense prospective for biopharmaceutical importance for example their  
51  
52 80 use as antibacterial, antiviral, and antitumor agents (Cameotra and Makkar, 2004). Lipopeptides shows insecticide  
53  
54 81 potential against *Drosophila melanogaster* (Assie et al. 2002), their larvicidal properties besides mosquito vectors  
55  
56 82 has not much tested.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

83 Biosurfactants are biologically active compounds produced by groups of microorganisms that utilize their  
84 energy sources such as oils, hydrocarbons and simple sugars (Parthipan et al. 2017). They have the capabilities to  
85 reduce surface/interface tension with liquid and solid substances (Das and Mukherjee, 2007). Biosurfactants are  
86 extensively utilized for numerous intentions in different sectors like, oil recovery process, food processing industry,  
87 cleaning purpose, crude oil drilling lubricants, bioremediation of oil contaminated sites and pharmaceutical industry  
88 (Makkar et al. 2011; Freitas de Oliveira et al. 2013; Parthipan et al. 2017). Biosurfactants have many advantages,  
89 i.e., they are less toxic, eco-friendly, easily degradable, very stable/active in high temperature/salinity regions and  
90 can easily producible using cheap organic sources (Rienzo et al. 2016). The present research is deal with  
91 biosynthesis and characterization of biosurfactant using two bacterial strains such as *Bacillus subtilis* A1 and  
92 *Pseudomonas stutzeri* NA3 for mosquito control in laboratory conditions.

93  
94 **Materials and methods**

95  
96 **Bacterial strain and culture conditions**

97  
98 In this study, two bacterial strains were used, *Bacillus subtilis* A1 (KP895564) and *Pseudomonas stutzeri* NA3  
99 (KU708859), which are isolated in sample (A1 from crude oil and NA3 from injection water) collected at Indian  
100 crude oil reservoir. These bacterial strains were sub-cultured in Luria–Bertani (LB) medium (g/l 10.0 tryptone, 5.0  
101 yeast extract, 10.0 sodium chloride with 15.0 agar (Himedia, Mumbai, India)) and incubated for 24 hrs at 40°C for  
102 *B. subtilis* A1 and 30°C for *P. stutzeri* NA3 respectively. Further inoculums was prepared by picking single colonies  
103 and inoculated in LB broth (pH: 7.0) and incubated in orbital shaker incubator (150 rpm) at 37°C for 24 hrs  
104 (Parthipan et al. 2017).

105  
106 **Biosurfactant production**

107  
108 Biosurfactant production was carried as described in Parthipan et al. (2017). In brief, sterile minimal salt medium  
109 (MSM) (g/l: 0.5 FeCl<sub>3</sub>, 0.2 MgSO<sub>4</sub>, 1.0 NH<sub>4</sub>NO<sub>3</sub>, 1.0 KH<sub>2</sub>PO<sub>4</sub>, 1.0 K<sub>2</sub>HPO<sub>4</sub> and 0.02 CaCl<sub>2</sub> (Himedia, Mumbai,  
110 India)), supplemented with 2% sucrose as carbon source. Pre-cultured bacterial strains *B. subtilis* A1 and

1  
2  
3  
4 111 *Pseudomonas stutzeri* NA3 were inoculated (initial load:  $1.6 \times 10^4$  CFU ml<sup>-1</sup>) and incubated for 120 hrs at 40°C for  
5  
6 112 *B. subtilis* A1 and 30°C for *P. stutzeri* NA3 respectively in an orbital shaker (150 rpm). After incubation period,  
7  
8 113 bacterial biomass was separated by centrifugation with 3400 x g at 4°C for 20 min (refrigerated centrifuge, Remi-  
9  
10 114 India: R-248). Further biosurfactant containing solutions were acidified (pH 2.0) by help of 6 M HCl. The acidified  
11  
12 115 solutions were kept at 4 °C for overnight to complete precipitation. Further precipitated biosurfactants were  
13  
14 116 collected by centrifugation at 8000 g for 20 min at 4 °C and dissolved in deionized sterile water (pH 7.0), followed  
15  
16 117 by extraction using 65:15 ratio of chloroform:methanol. These solvents were removed using a rotary evaporator, and  
17  
18 118 the biosurfactant phase was sluiced with three volumes of hexane to eliminate free fatty acids, alcohols and alkanes.  
19  
20 119 This procedure was repeated three times. The biosurfactant was further characterized using FT-IR and GC-MS  
21  
22 120 spectroscopy. Both biosurfactant obtained in this method were checked for oil displacement test (Hassanshahian,  
23  
24 121 2014)  
25  
26 122

### 123 **Characterization of biosurfactant**

124  
125 The obtained biosurfactant was characterized by Fourier transform infrared spectrum (FT-IR) and gas  
126 chromatographic mass spectrum (GC-MS). The functional groups of the biosurfactant recovered from both bacterial  
127 strains *B. subtilis* A1 and *P. stutzeri* NA3 were characterized using FT-IR (Perkin–Elmer, Nicolet Nexus - 470)  
128 (Parthipan et al. 2017). In brief, biosurfactant was mixed with KBr in the ratio of 1:100 to make pellet. Further  
129 prepared pellet was set aside in the sample holder and analysed in the IR region ranged between 400- 4000 cm<sup>-1</sup>.  
130 Further biosurfactants were characterised by Gas chromatography as described by Parthipan et al (2017). Briefly 1  
131 µl of methanol diluted samples was injected into a gas chromatograph (Shimadzu QP2010 Ultra, Rtx-5Sil MS (30 m  
132 × 0.25 mm ID × 0.25 µm). Helium was used as carrier gas with the flow rate of 1.5 ml min<sup>-1</sup> and the temperature of  
133 the GC injector was set as 260°C. The gradient temperature was set between 60 to 260°C at a rate of 5°C min<sup>-1</sup>,  
134 through an isothermal phase 10 min at the end of the run. The electron impact ion basis was constant at 200°C. Mass  
135 spectra were observed at 70 keV. The mass spectra were acquired with a m/z range: 40–600 ultra-high resolution  
136 approach with an acquisition speed of 6 spectra/sec. The detection of components was made in scan mode by using  
137 NIST11 and Wiley8 library.  
138

1  
2  
3  
4 **139 Mosquito rearing**

5  
6 140  
7  
8 141 Eggs of *Anopheles stephensi* were collected from water reservoirs within Coimbatore (Tamil Nadu, India) using an  
9  
10 142 “O”-type brush. Batches of 100–110 eggs were moved to 18×13×4 cm<sup>3</sup> enamel trays with 500mL of water. Here  
11  
12 143 eggs were permissible to hatch in lab setup (75–85% relative humidity (RH); 27±2 °C; 14:10 (L/D) photoperiod). 5g  
13  
14 144 of ground dog biscuits (Pedigree, USA) and hydrolyzed yeast (Sigma-Aldrich, Germany) in ratio of 3:1 were  
15  
16 145 provided as feed for *A. stephensi* larvae. Freshly brewed larvae and pupae were carefully collected and will be used  
17  
18 146 in the toxicity experiments (Anitha et al. 2016).

19  
20 147  
21  
22 148 **In-vitro larvicidal and pupicidal toxicity assay**

23  
24 149  
25  
26 150 About 25 *A. stephensi* larvae (I, II, III or IV instar) and pupae were positioned for 24 h in a glass beaker containing  
27  
28 151 250 ml of dechlorinated water with desired concentration of the both biosurfactant separately. 0.5 mg of larval food  
29  
30 152 was supplied for each concentration of biosurfactant (Kovendan et al. 2012). Each concentration was repeated 5  
31  
32 153 times against all instars. Control mosquitoes were bared for 24 h to the solvent, mortality percentage was calculated  
33  
34 154 using following formula:

35  
36  
37 155 
$$\text{Percentage mortality} = (\text{number of dead individuals} / \text{number of treated individuals}) * 100$$

38  
39 156 **Data analysis**

40  
41 157 In mosquito controlling experiments, lethal concentration (LC) 50 and LC90 were calculated by probit  
42  
43 158 analysis, as described by Finney (1971). Mosquitocidal efficiency was analyzed using analysis of variance  
44  
45 159 (ANOVA,) means were separated using Tukey's HSD test. P<0.05 was considered as significance of differences  
46  
47 160 among means.

48  
49 161  
50  
51 162 **Results**

52  
53 163  
54  
55 164 **Biosurfactant characterization**

56  
57 165  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 166 Biosurfactants were produced effectively using both A1 and NA3 strains with the optimized production conditions  
5  
6 167 as reported in the Parthipan et al. (2017). Biosurfactant produced *B. subtilis* gave considerably level of  
7  
8 168 emulsification activity 84% (as reported in Parthipan et al. (2017)). Both strains showed the effective oil-  
9  
10 169 displacement activity as shown in Fig. 1a and 1b. Higher amount of biosurfactant was produced by the *B. subtilis*  
11  
12 170 A1 (4.85 g l<sup>-1</sup>) as reported earlier (Parthipan et al. 2017), similarly strain NA3 was produced 3.81 g l<sup>-1</sup> of  
13  
14 171 biosurfactant. FT-IR was analysed and observed that biosurfactant produced by A1 contains following functional  
15  
16 172 groups of -OH, P-H<sub>2</sub>, C=O, -CH<sub>3</sub>, -CH<sub>2</sub>-, -COOH, O-H, CH<sub>2</sub> and C-I (Parthipan et al. 2017).  
17

18 173 Similarly biosurfactant produced by NA3 also characterized by FT-IR and predicted the numerous peaks at  
19  
20 174 different positions were shown in Fig. 2. Peak at 1442cm<sup>-1</sup> was due to the presence of N-H; peaks at 2923, 2853,  
21  
22 175 1461 and 1391cm<sup>-1</sup> indicates the presence of aliphatic chains (-CH<sub>3</sub> and -CH<sub>2</sub>-). A strong peak at 1639 specified  
23  
24 176 that occurrence of CO-N bond. Presences of peaks at 1091 and 722cm<sup>-1</sup> may match to the C-N stretching vibrations.  
25  
26 177 With the previous literature about lipopeptide biosurfactant, these FT-IR descriptions confirmed that presence of the  
27  
28 178 aliphatic groups joined with a peptide moiety as distinguishing properties of lipopeptide biosurfactant (Zou et al.  
29  
30 179 2014).  
31

32 180 The gas chromatography and mass spectrum characterization further revealed that the biosurfactant  
33  
34 181 extracted from both strains were lipopeptide in nature. Compounds obtained from strain A1 were reported as fatty  
35  
36 182 acids, such as hexadecanoic acid, octadecadienoic acid and octadecenoic acid (Parthipan et al. 2017). Similarly NA3  
37  
38 183 also predicted with numerous fatty acid peaks at different retention time (RT) as below, 1-Dodecanol (Fig. 3a) (RT:  
39  
40 184 16.16, molecular weight (MW): 186, chemical formula (CF): C<sub>12</sub>H<sub>26</sub>O)), oleic acid (Fig. 3b) (RT: 20.55, MW: 282,  
41  
42 185 CF: C<sub>18</sub>H<sub>34</sub>O<sub>2</sub>), hexanoic acid, octadecyl ester (Fig. 3c) (RT: 22.82, MW: 368, CF: C<sub>24</sub>H<sub>48</sub>O<sub>2</sub>).  
43  
44 186

#### 45 187 **Biosurfactant toxicity assay against *A. stephensi***

46 188  
47  
48  
49 189 Table 1 indicates that the larvicidal and pupicidal activities of biosurfactant synthesized by *B. subtilis* A1 at different  
50  
51 190 concentrations in laboratory conditions. Biosurfactant from A1 was found to be highly toxic to the larva and pupa of  
52  
53 191 *A. stephensi* whose LC<sub>50</sub> values were 3.58 (I), 4.92 (II), 5.73 (III), 7.10 (IV) and 7.99 (V) for the different stages of  
54  
55 192 the life span. Similarly, mosquitocidal activity of NA3 (Table 2) showed their toxicity with the LC 50 values of  
56  
57 193 2.61 (I), 3.68 (II), 4.48 (III), 5.55 (IV) and 6.99 (pupa) which confirms that the both biosurfactant contains  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 194 mosquitocidal components and its efficiency of the extermination was increased with increasing of concentrations  
5  
6 195 (ppm).

7  
8 196

9  
10 197 **Discussion**

11 198

12  
13  
14 199 In recent times, plentiful reports were available to support that *Bacillus* sp. and *Pseudomonas* sp. were effective  
15  
16 200 biosurfactant producers and also widely used for extensive range of application in many field such as: oil recovery,  
17  
18 201 bioremediation, industrial application and biodegradation (Pereira et al. 2013; Pacwa-Plociniczak et al. 2014;  
19  
20 202 Cubitto et al. 2004 Greenwell et al. 2016; Ismail et al. 2013; Parthipan et al. 2017). Unfortunately very few reports  
21  
22 203 were available on the mosquitocidal application of the biosurfactant produced by *Bacillus* sp. and *Pseudomonas* sp.  
23  
24 204 (Das and Mukherjee, 2006; Geetha and Manonmani, 2010; Geetha et al. 2010). Biosurfactants are habitually a  
25  
26 205 mixture of molecules such as fatty acids, peptides and polysaccharide; it could be any one form as like lipopeptides,  
27  
28 206 lipoproteins, glycolipids, phospholipids and lipopolysaccharides based on the biosurfactant producers. Interestingly  
29  
30 207 some of these compounds were highly toxic to the many of organisms such as insects and mosquitos, which are very  
31  
32 208 dangerous to the human health (Das and Mukherjee, 2006; Geetha and Manonmani, 2010; Geetha et al. 2010). The  
33  
34 209 FTIR analysis reveals that occurrence of ester carbonyl groups, phosphines in phosphoserine, carboxylic acids and  
35  
36 210 Carbon–Iodine. Based on this observation biosurfactant obtained from bacterium *B. subtilis* A1 and *P. stutzeri* NA3  
37  
38 211 was categorized as lipopeptide in nature (Rodrigues et al. 2006; Parthipan et al. 2017).

39  
40 212 The LC50 values of the present study revealed that the significant difference in the production of  
41  
42 213 mosquitocidal metabolites of both biosurfactant produced by the bacterial strains *B. subtilis* A1 and *P. stutzeri* NA3.  
43  
44 214 Pupae of mosquitoes needed the accessible atmospheric oxygen for their respiration and other functions. Due to  
45  
46 215 reduction in surface tension of water by the auctions of lipopetides, the pupae were incapable to come up to the  
47  
48 216 surface of water for their oxygen needs and had to remain submerged in water, these unusual circumstances lead  
49  
50 217 them to fatality (Piper and Maxwell, 1971). Hence, the mosquito pupal mortality observed in our study could be  
51  
52 218 primarily by reduction in surface tension of the water caused by the bacterial biosurfactant. However, the possibility  
53  
54 219 of its action on the cuticle of the pupae cannot be ruled out as there are reports on the action of surfactin on  
55  
56 220 biological membranes (Vollenbroich et al. 1997a&b; Buchoux et al. 2008). Trace elements and carbon level in the  
57  
58 221 culture medium have been reported to enhance the lipopeptide production by the strains *B. subtilis* (Wei et al. 2007).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

222 These lipopeptides will get bind with the sulphur group of DNA, leading to the rapid denaturation of organelles and  
223 enzymes in mosquito. Subsequently, decreases in the membrane permeability and disturbance in proton motive force  
224 may cause loss of cellular function and also cell death. Further research on this issue is required (Benelli, 2016a,b).

225 Present research outcomes were supported by some of the previous research findings as the biosurfactant  
226 used as mosquito controlling agent. Deepali et al. (2014) isolated biosurfactant from *Stenotrophomonas maltophilia*,  
227 also identified that biosurfactant was rhamnolipid in nature and it has no activity for 4mg/l concentration but  
228 increasing the concentration to 10 mg/l gives larvicidal properties. Recently Geetha and Manonmani, (2010) isolated  
229 surfactin a biosurfactant compound from the strain *Bacillus subtilis* ssp. *subtilis* and reported that surfactin showed  
230 mosquito pupicidal activity (Geetha et al. 2010). Similarly Das and Mukherjee, (2006) used lipopeptides extracted  
231 from *B. subtilis* strain for mosquito larvicidal uses. Recently Silva et al. (2015) reported that the rhamnolipid as an  
232 eco-friendly surfactants and it has many activities like larvicidal, insecticidal, and repellent activities against *A.*  
233 *egypti*, but this achievements were obtained using very high level of biosurfactant 1 g/l as reported. As compared to  
234 these studies, present study achieved higher mortality rate in very low concentration (2-10ppm). These observations  
235 signify that the biosurfactant produced by strains A1 and NA3 were highly active at low concentrations and may be  
236 used as potential mosquitocidal agent.

237

238 **Conclusion**

239

240 In conclusion, these observations showed that the biosurfactant produced by both bacterial strains *B. subtilis* A1 and  
241 *P. stutzeri* NA3 were lipopeptide in nature and which are accountable of the *A. stephensi* mosquito larvicidal and  
242 pupicidal activities. This mosquitocidal action of the both biosurfactant on the larvae and pupae may be due to  
243 reduction in the surface tension of water, its direct to the oxygen deficiency at underwater where larvae and pupae  
244 exit and this abnormal condition lead them to dead. There are limited reports only available on the mosquitocidal  
245 effects of the biosurfactant produced by bacterial strains. As there are limited bio-control methods only available  
246 against mosquito to eradicate, it could be a hopeful method for control of the mosquito spread. Further studies  
247 needed to extend these observations to external uses in the form of effective controlling agent.

248

249

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**250 Acknowledgments**

251  
252 A. Rajasekar is thankful to the Department of Biotechnology (Government of India) for the award of the  
253 Ramalingaswami re-entry Fellowship (BT/RLF/Re-entry/17/2012), Department of Science and Technology for the  
254 young scientist award (SB/YS/LS-40/2013), University Grants Commission-MRP (MRP-MAJOR-MICRO-2013-  
255 31825) and Science and Engineering Research Board, Department of Science and Technology, Government of  
256 India (EEQ/2016/000449).

257

**258 Compliance with ethical standards**

259  
260 All applicable international and national guidelines for the care and use of animals were followed. All procedures  
261 performed in studies involving animals were in accordance with the ethical standards of the institution or practice at  
262 which the studies were conducted

263

**264 Conflict of interest**

265

266 The authors declare that they have no conflict of interest.

267

**268 References**

269

270 Anitha J, Murugan K, Panneerselvam C, Madhiyazhagan P, Dinesh D, Vadivalagan C, Aziz AT, Chandramohan  
271 B, Suresh U, Rajaganesh R, Subramaniam J, Marcello N, Higuchi A, Alarfaj AA, Munusamy MA, Kumar  
272 S, Benelli G (2016) Earthworm-mediated synthesis of silver nano-particles: a potent tool against  
273 hepatocellular carcinoma, pathogenic bacteria, Plasmodium parasites and malaria mosquitoes. Parasitol  
274 Int 65:276–284

275 Assie LK, Deleu M, Arnaud L, Paquot M, Thonart P, Gaspar Ch, Haubruge E (2002) Insecticide activity of  
276 surfactins and iturins from a biopesticide *Bacillus subtilis* Cohn (S499 strain). Meded Rijksuniv Gent Fak  
277 Landbouwkd Toegep Biol Wet 67:647–655

1  
2  
3  
4 278 Benelli G (2015a) Plant-borne ovicides in the fight against mosquito vectors of medical and veterinary importance:  
5  
6 279 a systematic review. *Parasitol Res* 114:3201–3212  
7  
8 280 Benelli G (2015b) Research in mosquito control: current challenges for a brighter future. *Parasitol Res* 114:2801–  
9  
10 281 2805  
11  
12 282 Benelli, G, LoIacono A, Canale A, Mehlhorn H (2016a) Mosquito vectors and the spread of cancer:an over looked  
13  
14 283 connection?. *Parasitol Res* 115:2131-2137  
15  
16 284 Benelli G, Pavela R, Canale A, Mehlhorn H (2016b) Tick repellents and acaricides of botanical origin: a green  
17  
18 285 road map to control tick- borne diseases?. *Parasitol Res* 115:2545-2560  
19  
20 286 Buchoux S, Lai-Kee-Him J, Garnier M, Tsan P, Besson F, Brisson A, Dufourc EJ (2008) Surfactin-triggered small  
21  
22 287 vesicle formation of negatively charged membranes: A novel membrane-lysis mechanism. *Biophys J*  
23  
24 288 95:3840–849  
25  
26 289 Cameotra SS, Makkar RS (2004) Recent applications of biosurfactants as biological and immunological  
27  
28 290 molecules. *Curr Opin Microbiol* 7:262–266  
29  
30 291 Cubitto MA, Moran AC, Commendatore M, Chiarello MN, Baldini MD (2004) Effects of *Bacillus subtilis* O9  
31  
32 292 biosurfactant on the bioremediation of crude oil-polluted soil. *Biodegradation* 15:281–287  
33  
34 293 Darriet F, Hougard JM (2002) An isolate of *Bacillus circulans* toxic to mosquito larvae. *J Am Mosq Control*  
35  
36 294 *Assoc* 18:65–67  
37  
38 295 Das K, Mukherjee AK (2006) Assessment of mosquito larvicidal potency of cyclic lipopeptides produced by  
39  
40 296 *Bacillus subtilis* strains. *Acta Trop* 97:168–173  
41  
42 297 Das K, Mukherjee AK (2007) Comparison of lipopeptide biosurfactants production by *Bacillus subtilis* strains in  
43  
44 298 submerged and solid state fermentation systems using a cheap carbon source: Some industrial  
45  
46 299 applications of biosurfactants. *Process Biochem* 42:1191–1199  
47  
48 300 Deepali K, Sneha P, Sucheta P (2014) Larvicidal activity of rhamnolipid biosurfactant produced by  
49  
50 301 *Stenotrophomonas maltophilia*. *Int J Sci Eng Res* 2229-5518  
51  
52 302 Finney DJ (1971) Probit analysis, 3rd ed. Cambridge University Press, Cambridge  
53  
54 303 Freitas de Oliveira DW, Franc IWL, Felix AKN, Martins JJJ, Giro MEA, Melob VMM, Goncalves LRB (2013)  
55  
56 304 Kinetic study of biosurfactant production by *Bacillus subtilis* LAMI005 grown in clarified cashew apple  
57  
58 305 juice. *Colloids Surf. B Biointerfaces* 101:34–43  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

306 Geetha I, Manonmani AM (2010) Surfactin: a novel mosquitocidal biosurfactant produced by *Bacillus subtilis* ssp.  
307 *subtilis* (VCRC B471) and influence of abiotic factors on its pupicidal efficacy. Lett Appl Microbiol  
308 51:406–412

309 Geetha I, Manonmani AM, Paily KP (2010) Identification and characterization of a mosquito pupicidal metabolite  
310 of a *Bacillus subtilis* subsp. *subtilis* strain. Appl Microbiol Biotechnol 86:1737–1744

311 Govindarajan M, Kadaikunnan S, Alharbi NS, Benelli (2016) Acute toxicity and repellent activity of the  
312 *Origanum scabrum* Boiss. & Heldr. (*Lamiaceae*) essential oil against four mosquito vectors of public  
313 health importance and its biosafety on non-target aquatic organisms. Environ Sci Pollut Res 23:23228-  
314 23238

315 Greenwell M, Sarker MH, Rahman PKSM (2016) Biosurfactant production and biodegradation of leather dust  
316 from tannery. Open Biotechnol J 10:1–14

317 Hassanshahian M (2014) Isolation and characterization of biosurfactant producing bacteria from Persian Gulf  
318 (Bushehr provenance). Mar Pollut Bull 86:361–366

319 Ismail W, Al-Rowaihi IS, Al-Humam AA, Hamza RY, El Nayal AM, Bououdina M (2013) Characterization of a  
320 lipopeptide biosurfactant produced by a crude-oil-emulsifying *Bacillus* sp. I-15. Int Biodeterior Biodegrad  
321 84:168–178

322 Knight RL, Walton WE, O’Meara GF, Reisen WK, Wass R (2003) Strategies for effective mosquito control in  
323 constructed treatments wetlands. Ecol Eng 21:211–232

324 Kovendan K, Murugan K, Vincent S, Barnard DR (2012) Studies on larvicidal and pupicidal activity of *Leucas*  
325 *aspera* Willd (*Lamiaceae*) and bacterial insecticide, *Bacillus* sp *haericus* against malarial vector  
326 *Anopheles stephensi* Liston (Diptera: *Culicidae*). Parasitol Res 110:195–203

327 Mahesh Kumar P, Murugan K, Kovendan K, Subramaniam J, Amaresan D (2012) Mosquito larvicidal and  
328 pupicidal efficacy of *Solanum xanthocarpum* (Family: *Solanaceae*) leaf extract and bacterial insecticide,  
329 *Bacillus thuringiensis*, against *Culex*. Parasitol Res 110:2541–2550

330 Maheswaran R, Ignacimuthu S (2014) Effect of confertifolin from *Polygonum hydropiper* L. against dengue  
331 vector mosquitoes *Aedes aegypti* L. Environ Sci Pollut Res 113:3143-3150

332 Makkar RS, Cameotra SS, Banat IM (2011) Advances in utilization of renewable substrates for biosurfactant  
333 production. AMB Express 1:5

- 1  
2  
3  
4 334 Mehlhorn H (2008) Encyclopedia of Parasitology, third ed. Springer, Heidelberg  
5  
6 335 Mittal PK (2003) Biolarvicides in vector control: challenges and prospects. J Vect Borne Dis 40:20–32  
7  
8 336 Mukandiwa L, Eloff JN, Naidoo V (2016) Repellent and mosquitocidal effects of leaf extracts of *Clausena anisata*  
9  
10 337 against the *Aedes aegypti* mosquito (Diptera: Culicidae), Environ Sci Pollut Res 23:11257-11266  
11  
12 338 Murugan K, Panneerselvam C, Subramaniam J, Madhiyazhagan P, Hwang JS, Wang L, Dinesh D, Suresh U, Roni  
13  
14 339 M, Higuchi A, Nicoletti M, Benelli G (2016) Eco-friendly drugs from the marine environment:  
15  
16 340 spongweed-synthesized silver nanoparticles are highly effective on *Plasmodium falciparum* and its  
17  
18 341 vector *Anopheles stephensi*, with little non-target effects on predatory copepods, Environ Sci Pollut Res  
19  
20 342 23:16671-16685  
21  
22 343 Murugan K, Venus JS, Panneerselvam C, Bedini S, Conti B, Nicoletti M, Sarkar SK, Hwang JS, Subramaniam J,  
23  
24 344 Madhiyazhagan P, Kumar PM, Dinesh D, Suresh U, Benelli G (2015) Biosynthesis, mosquitocidal and  
25  
26 345 antibacterial properties of *Toddalia asiatica*-synthesized silver nanoparticles: do they impact predation of  
27  
28 346 guppy *Poecilia reticulata* against the filariasis mosquito *Culex quinquefasciatus*?. Environ Sci Pollut Res  
29  
30 347 22:17053-17064  
31  
32 348 Pacwa-Plociniczak M, Płaza GA, Poliwoda A, Piotrowska-Seget Z (2014) Characterization of hydrocarbon-  
33  
34 349 degrading and biosurfactant-producing *Pseudomonas* sp. P-1 strain as a potential tool for bioremediation  
35  
36 350 of petroleum-contaminated soil. Environ Sci Pollut Res 21:9385–9395  
37  
38 351 Parthipan P, Preetham E, Machuca LL, Rahman PKSM, Murugan K, Rajasekar A (2017) Biosurfactant and  
39  
40 352 degradative enzymes mediated crude oil degradation by bacterium *Bacillus subtilis* A1. Front Microbiol  
41  
42 353 8:193  
43  
44 354 Pereira JFB, Gudina EJ, Vitorino CR, Teixeira JA, Coutinho JAP (2013) Optimization and characterization of  
45  
46 355 biosurfactant production by *Bacillus subtilis* isolates towards microbial enhanced oil recovery  
47  
48 356 applications. Fuel 111:259–268  
49  
50  
51 357 Piper WD, Maxwell KE (1971) Mode of action of surfactants on mosquito pupae. J Econ Entomol 64:601–606  
52  
53 358 Poopathi S, De Britto LJ, Praba VL, Mani C, Praveen M (2015) Synthesis of silver nanoparticles from *Azadirachta*  
54  
55 359 *indica*—a most effective method for mosquito control, Environ Sci Pollut Res 22:2956-2963  
56  
57 360 Prabakaran G Hoti SL, Rao HS, Vijjapu S (2015) Di-rhamnolipid is a mosquito pupicidal metabolite from  
58  
59 361 *Pseudomonas fluorescens* (VCRC B426). Acta Trop 148:24-31  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

362 Rienzo MADD, Kamalanathan ID, Martin PJ (2016) Comparative study of the production of rhamnolipid  
363 biosurfactants by *B. thailandensis* E264 and *P. aeruginosa* ATCC 9027 using foam fractionation. Process  
364 Biochem 51:820-827

365 Rodrigues LR, Teixeira JA, Van Der Mei HC, Oliveira R (2006) Isolation and partial characterization of a  
366 biosurfactant produced by *Streptococcus thermophilus* A. Colloids Surf. B Biointerfaces 53:105–112

367 Silva VL, Lovaglio RB, Von Zuben CJ and Contiero J (2015) Rhamnolipids:solution against *Aedes aegypti*?. Front  
368 Microbiol 6:88

369 Subramaniam J, Murugan K, Panneerselvam C, Kovendan K, Madhiyazhagan P, Dinesh D, Mahesh Kumar P,  
370 Chandramohan B, Suresh U, Rajaganesh R, Saleh Alsalhi M, Devanesan S, Nicoletti M, Canale A,  
371 Benelli G Multipurpose effectiveness of *Couroupita guianensis*-synthesized gold nanoparticles: high  
372 antiplasmodial potential, field efficacy against malaria vectors and synergy with *Aplocheilus lineatus*  
373 predators. Environ Sci Pollut Res 23:7543-7558

374 Subramaniam J, Murugan K, Panneerselvam C, Kovendan K, Madhiyazhagan P, Kumar PM, Dinesh D,  
375 Chandramohan B, Suresh U, Nicoletti M, Higuchi A, Hwang JS, Suresh K, Alarfaj AA, Munusamy MA,  
376 Messing RH, Benelli G (2015) Eco-friendly control of malaria and arbovirus vectors using the  
377 mosquitofish *Gambusia affinis* and ultra-low dosages of *Mimusops elengi*-synthesized silver  
378 nanoparticles: towards an integrative approach?. Environ Sci Pollut Res 22:20067-20083

379 Vollenbroich D, Ozel M, Vater J, Kamp RM, Pauli G (1997) Mechanism of inactivation of enveloped viruses by  
380 the biosurfactant surfactin from *Bacillus subtilis*. Biologicals 25:289–97

381 Vollenbroich D, Pauli G, Ozel M, Vater J (1997) Antimycoplasma properties and application in cell culture of  
382 surfactin, a lipopeptide antibiotic from *Bacillus subtilis*. Appl Environ Microbiol 63:44–49

383 Wei YH, Lai CC, Chang JS (2007) Using taguchi experimental design methods to optimize trace element  
384 composition for enhanced surfactin production by *Bacillus subtilis* ATCC 21332. Process Biochem  
385 42:40–45

386 Zou C, Wang M, Xing Y, Lan G, Ge T, Yan X, Gu T (2014) Characterization and optimization of biosurfactants  
387 produced by *Acinetobacter baylyi* ZJ2 isolated from crude oil-contaminated soil sample towards  
388 microbial enhanced oil recovery applications. Biochem Eng J 90:49-58

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure Captions:**

390  
391  
392  
393  
394  
395  
396

**Fig. 1** Oil displacement activity of the biosurfactant: (a) *B. subtilis* A1; (b) *Pseudomonas stutzeri* NA3.

**Fig. 2** FT-IR spectrum of biosurfactant produced by strain *Pseudomonas stutzeri* NA3.

**Fig. 3** Mass spectrum of the biosurfactant isolated from *Pseudomonas stutzeri* NA3: (a) 1-Dodecanol; (b) oleic acid; (c) hexanoic acid, octadecyl ester.

**Fig. 1**








Fig. 2



**Fig. 3**



**Table 1** Mosquitocidal effect of the biosurfactant isolated from *B. subtilis* A1 against the malarial vector *Anopheles stephensi*.

No mortality was observed in the control.

Mosquito life stages	Percentage of larval and pupal mortality					LC <sub>50</sub> and (LC <sub>90</sub> )	95% confidence Limit		Regression equation	$\chi^2$ (d.f.=4)
	Concentration (ppm)						LC <sub>50</sub> (LC <sub>90</sub> )			
	2	4	6	8	10		Lower	Upper		
I	37.2 ±1.30	51 ± 2.12	70.2±1.92	87.6 ±2.07	98.6 ±1.14	3.589 (8.211)	2.987 (7.548)	4.085 (9.108)	$y = -0.995 + 0.277 x$	3.352 <i>n.s</i>
II	29.6 ±1.67	35.2±1.64	61.8±1.30	73.6 ±1.67	89.8±0.83	4.925 (10.596)	4.335 (9.652)	5.453 (11.939)	$y = -1.113 + 0.226 x$	3.494 <i>n.s</i>
III	22.2± 1.30	34.4±1.51	56.2 ±1.48	67.8±2.86	78.8±1.64	5.733 (12.128)	5.126 (10.918)	6.318 (13.930)	$y = -1.149 + 0.200 x$	1.015 <i>n.s</i>
IV	19.2 ±1.09	31 ±1.22	43.4±2.30	59.4±1.14	65±1.58	7.109 (15.036)	6.397 (13.112)	7.953 (18.224)	$y = - 1.149 + 0.162 x$	1.066 <i>n.s</i>
Pupa	15 ± 1.87	26.6± 0.89	37 ± 2.23	51.4± 1.67	61±0.70	7.993 (15.778)	7.247 (13.737)	8.979 (19.170)	$y = - 1.316 + 0.165 x$	0.548 <i>n.s</i>

LC50= lethal concentration (ppm) that kills 50% of the exposed organisms.

LC90= lethal concentration (ppm) that kills 90% of the exposed organisms.

 $\chi^2$ =chi-square.

d.f.= degrees of freedom.

n.s.= not significant ( $\alpha=0.05$ ).

**Table 2** Mosquitocidal effect of the biosurfactant isolated from *P. stutzeri* NA3 against the malarial vector *Anopheles stephensi*.

Mosquito life stages	Percentage of larval and pupal mortality					LC <sub>50</sub> and (LC <sub>90</sub> )	95% confidence Limit		Regression equation	$\chi^2$ (d.f.=4)	
	Concentration (ppm)						LC <sub>50</sub> (LC <sub>90</sub> )	Lower			Upper
	2	4	6	8	10						
I	45.4± 1.34	63.6± 0.89	80.8± 1.30	94.4± 1.94	99.8± 0.44	2.618 (7.151)	1.874 (6.534)	3.186 (7.986)	y = -0.740 +0.283 x	0.846 n.s	
II	39 ± 0.70	51.6± 1.51	64.2± 1.92	83± 1.87	92.6±1.51	3.680 (9.709)	2.893 (8.784)	4.299 (11.054)	y= -0.782 +0.213 x	1.815 n.s	
III	32.8± 1.09	47.6± 1.14	58 ± 2.91	75.6± 2.07	84± 1.58	4.481 (11.541)	3.679 (10.287)	5.137 (13.480)	y= -0.813 +0.182 x	0.553 n.s	
IV	27.4± 1.14	39.2± 1.92	54.6± 2.07	68.4± 2.50	73.6±1.81	5.553 (13.490)	4.786 (11.825)	6.264 (16.216)	y= -0.897 +0.161x	1.024 n.s	
Pupa	22.6± 1.81	32.8± 1.48	44± 1.58	57.6± 2.30	66± 2.54	6.999 (15.641)	6.231 (13.464)	7.912 (19.408)	y= - 1.038 +0.148 x	0.187 n.s	

No mortality was observed in the control.

LC50= lethal concentration (ppm) that kills 50% of the exposed organisms.

LC90= lethal concentration (ppm) that kills 90% of the exposed organisms.

$\chi^2$ =chi-square.

d.f.= degrees of freedom.

n.s.= not significant ( $\alpha=0.05$ ).