

Title:

Applied Physiology of Female Soccer: An Update

Authors:

Naomi Datson^{1,2}, Andrew Hulton², Helena Andersson³, Tracy Lewis¹, Matthew Weston⁴, Barry Drust² and Warren Gregson^{2,5}

¹ The English Football Association, St. George's Park, Burton Upon Trent, UK

² Research Institute of Sport Sciences, Liverpool John Moores University, Liverpool, UK

³ The Swedish Football Association, Stockholm, Sweden

⁴ School of Social Sciences and Law, Teesside University, Teesside, UK

⁵ ASPIRE, Academy for Sports Excellence, Doha, Qatar

Corresponding Author:

Name: Naomi Datson

Email Address: Naomi.Datson@thefa.com

Telephone Number: +44844 980 8200 (6940)

Abstract

The popularity and professionalism of female soccer has increased markedly in recent years, with elite players now employed on either a professional or semi-professional basis. The previous review of the physiological demands of female soccer was undertaken two decades ago when the sport was in its relative infancy. Increased research coupled with greater training and competition demands warrants an updated review to consider the effect on physical performance and injury patterns.

The physical demands of match-play along with the influence of factors such as the standard of competition, playing position and fatigue have been explored. Total distance covered for elite female players is approximately 10 km, with 1.7 km completed at high-speed ($>18 \text{ km}\cdot\text{h}^{-1}$). Elite players complete 28% more high-speed running and 24% more sprinting than moderate level players. Decrements in high-speed running distance have been reported between and within halves, which may indicate an inability to maintain high-intensity activity. Although physical capacity of female players is the most thoroughly researched area comparisons are difficult due to differing protocols. Elite players exhibit maximal oxygen uptake ($\text{VO}_{2\text{max}}$) values of $49.4 - 57.6 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$, Yo Yo Intermittent Endurance test level 2 (YYIE2) scores of $1774 \pm 532 \text{ m}$ (mean \pm SD) and 20 m sprint times of $3.17 \pm 0.03 \text{ s}$ (mean \pm SD). Reasons for the increased prevalence of anterior cruciate ligament injuries in females (2-6 times greater than males) are discussed with anatomical, biomechanical loading and neuromuscular activation differences being cited in the literature. This review presents an in-depth contemporary examination of the applied physiology of the female soccer player.

1) Introduction

The last review of the physiological demands of female soccer was undertaken two decades ago [1]. At the time of that review, women's soccer was in its relative infancy with the inaugural Women's World Cup having taken place in 1991. The popularity of women's soccer has increased markedly in recent years with 29 million participants recorded in 2011, an increase of 34% from 2000 [2]. The 2015 Women's World Cup will have 24 participating nations, an increase of eight from the previous tournament.

Since the previous review, increasing attention in the scientific literature has focused on women's soccer with over two thousand scientific articles published. There has been a drive towards professionalism in the sport with players at the highest level now employed on a professional or semi-professional basis. Current elite players are exposed to greater training volumes and competition demands than ever before, consequently these increased physical demands may have implications for both physical performance levels and injury patterns. In-line with these changes, advances in technology available for monitoring athletes has led to greater opportunities to scientifically support players and thus information regarding the modern elite female player is increasingly available. It is evident that an updated review into the applied physiology of the female game and its players is warranted to examine the recent research while also highlighting directions for future work. To ensure all relevant empirical studies on female soccer players were examined, a search on PubMed and Google Scholar was performed using a combination of the following terms: 'soccer', 'football', 'female' and 'women'. Manual secondary searches were undertaken on the reference lists of recovered articles. As this was a narrative and not a systematic review, the papers that were believed to be the most relevant were selected. No date limit was set for the search period and only articles written in English were considered. Articles that did not focus on the physical aspects of performance and injury were excluded.

2) Match Physical Performance

A comprehensive understanding of the demands of the game is necessary to apply a systematic approach to training [3]. The use of technologies such as Global Positioning System (GPS) devices and semi-automated camera systems are now relatively commonplace within men's soccer. These technologies are more time efficient than the traditional video-based time motion analysis systems and offer greater objectivity and volume of information [4]. Furthermore, more detailed evaluations of specific elements of a player's physical performance as opposed to generic time-motion analysis can be undertaken [5-8]. Despite advancements in the understanding of match-play in elite male players, published research on elite female players is lacking [9], predominantly due to stadium access and financial considerations.

At the time of the previous review [1] there were no published findings on the activity profile of female players. Early unpublished data collected via video-tape [1] ($n = 7$) of national team players showed individuals covered a total distance of 8.47 ± 2.2 km (mean \pm SD) and average sprint distances were 14.9 ± 5.6 m (mean \pm SD), although the speed zone for sprinting was not defined. More recent studies [10-13], using larger sample sizes ($n = 13 - 58$) have reported an increase in total distance covered (~ 10 km) for national team players when measured using video-tape and GPS technology. Whilst a trend exists for an increase in total distance covered compared to early reports, these differences may also to some extent reflect differences in data collection methods [4]. Total distance covered provides a global estimation of physical match-play but it is high-speed running which is deemed to be more informative and reflective of the game [5,7]. In the previous review [1], the only reference to high-speed activity was the average sprinting distance. However, since this time national team players have been reported to cover a distance of 1.53 – 1.68 km at high speeds (>18 km.h⁻¹) [10,12,14]. Repeated sprinting and high-intensity bouts have also been examined with elite players performing 5.1 ± 5.1 and 31.2 ± 18.7 (mean \pm SD) respectively [15]. Explosive events and game specific skill involvements are also deemed important markers of physical match performance yet these areas have been subject to little attention in the literature. Game specific skill involvements (including passing, dribbling, tackling and trapping) have been quantified as 76 ± 30 events per match [11] and specifically 11 ± 1 headers and 16 ± 1 tackles [14] (mean \pm SD) with players also undertaking between 1350 and 1650 changes of activity [10,11,14].

The amount of high-speed running and sprinting undertaken by female players increases at higher standards of competition. Top-level players perform 28% more high-speed running (1.68 ± 0.09 km v 1.30 ± 0.10 km; effect size 4.0 (mean \pm SD)) and 24% more sprinting (0.46 ± 0.02 km v 0.38 ± 0.05 km; effect size 2.2 (mean \pm SD)) than moderate-level players [12]. Players produce more repeated-sprint bouts during international matches compared to national league (4.8 ± 2.8 bouts/match v 1.0 ± 1.0 bouts/match; effect size 2.0 (mean \pm SD)) [11]. Additionally, female players complete more high-speed running (13%) and sprinting (14%) when playing an international game compared to a domestic game [10]. An overview of the demands of elite female match-play taken from recent studies is shown in Table 1.

An evaluation of the demands of different playing positions is essential to ensure specificity of training and potential talent identification of players. Despite the previous review [1] stating that further investigation of positional variations was warranted there still appears to be a lack of information in this area. Observations indicate that midfielders cover greater total distance (10.67 ± 1.34 km) compared to defenders (9.62 ± 1.20 km) and attackers (9.61 ± 0.36 km) (mean \pm SD) [11]. These findings have been confirmed by GPS technology with midfielders covering over 1000 m more than attackers (9.64 km v 8.51 km) and 600 m more than defenders (9.64 km v 9.01 km) (mean \pm SD) [13]. Differences in high-speed activity also exist with central defenders performing less high-speed running (1.26 ± 0.11 km) than both midfielders and attackers (1.65 ± 0.11 km and 1.63 ± 0.10 km, respectively) (mean \pm SD) [14]. Attackers complete a greater amount of sprinting ($>25\text{km}\cdot\text{h}^{-1}$) (0.52 ± 0.03 km) than both midfielders and defenders (0.43 ± 0.04 km and 0.33 ± 0.05 km, respectively) (mean \pm SD) [14]. Future studies investigating the physical demands of different playing positions should consider the use of more specific positions (e.g. central defenders, wide defenders, central midfielders, wide midfielders and attackers) that are now commonly applied to research focussed on the male game [5,7]. The need to differentiate between central and wide positions in female soccer research is in accordance with recent studies, which have observed differences between these positions [16].

A change in physical performance, particularly high-speed running within and between each 45-minute half has frequently been used as a marker of progressive match fatigue, with results indicating that players experience temporary fatigue during and towards the end of a game [16]. Total distance covered has been shown to be consistent between halves (5.23 ± 0.09 km v 5.21 ± 0.08 km; effect size 0.2), however, significant reductions in high-speed running (0.68 ± 0.06 km v 0.62 ± 0.04 km; effect size 1.2) and sprinting (0.20 ± 0.03 km v 0.17 ± 0.02 km; effect size 1.0) occur (mean \pm SD) [17]. Some alterations in game specific skill involvements have also been demonstrated between halves with the number of tackles being greater in the first half compared to the second [14]. Reductions in repeated sprint and repeated high-intensity activity have not been observed between halves in elite players, however, an increased amount of low-intensity recovery between efforts occurs [15]. In an attempt to further examine the occurrence of fatigue, previous research has examined changes in physical performance within each half. Reductions in high-speed running distance have been reported with a 30% (0.27 to 0.19 km) and 34% (0.24 to 0.16 km) decrement from the first to the last 15-minute period observed in both the first and second half, respectively [18]. This reduction in high-speed running distance is even more pronounced when the same players compete in different levels of competition, with an 18% and 40% decline in the last to first 15 minutes of the second half for domestic and international competition, respectively [12]. The 5-minutes subsequent to the peak 5-minute period of high-speed running during match-play has frequently been used as an indicator of temporary fatigue and a 17% decrement in high-speed running was apparent in top-level players but not moderate-level players [14]. These findings suggest an inability to maintain high-intensity running and thus a failure to perform at the required intensity for the duration of the match. However, due to the tactical constraints of match-play these findings do not provide direct evidence of fatigue and as such a number of studies have assessed isolated physical performance parameters before and after match-play since these measures are not affected by tactical and technical decisions [19]. Decrements in sprint ($3.0 \pm 0.5\%$) [20], counter movement jump ($4.4 \pm 0.8\%$) [20], repeated sprint (4%) [21], intermittent endurance as measured by the Yo Yo Intermittent Endurance test level 2 (YYIE2) (62%) [21], peak torque knee extension ($7.1 \pm 1.9\%$) [20] and peak torque knee flexion ($9.4 \pm 1.8\%$) [20] have been highlighted (mean \pm SD). These types of assessment may give a clearer indication of fatigue development following a game.

The published studies to date indicate differences in the motion analysis of soccer players based upon the level of competition, stage of the game and playing position. The majority of these studies have been conducted using traditional single camera time-motion analysis methods pioneered in the 1970s. Modern technologies such as GPS and semi-automated camera systems are able to provide more detailed information across large samples of players relative to video-based approaches [4,22]. Consequently, more comprehensive evaluations of the physical characteristics of elite female match-play using contemporary analysis methods and more specific playing position classifications are warranted. The defining of individual speed threshold zones relative to a player's physical capabilities could also be considered for future studies. Male players have been shown to have individual thresholds for high-intensity running, as defined by the speed at the point of the second ventilatory threshold [23]. Accordingly, there is an apparent need to individualise the speed thresholds utilised. Furthermore it has been demonstrated that the use of default thresholds in semi-automated camera systems substantially underestimated the amount of high-intensity running distance completed [23]. Recent research [24] has examined maximum sprinting speed for female soccer players and these values could be used to contribute to a framework of speed threshold classification for female players. However, consideration must be given to classification and justification of the different match activity zones.

3) Physiological Demands of Match Play

3.1) Aerobic

Heart rate measurement is often used as an indirect measure of intensity of exercise and the validity of heart rate monitoring in soccer has formerly been described [25]. In the previous review [1], the only referenced heart rate data was during small-sided games. Since then, several studies have determined heart rate during competitive female match-play and noted average and peak heart rate values of $86 \pm 3\%$ of maximum heart rate (HR_{max}) and $98 \pm 1\%$ HR_{max} respectively (mean \pm SD). The assessment of heart rate and subsequent use of pre-established heart rate and maximal oxygen uptake (VO_{2max}) relationships are frequently used to estimate the aerobic contribution to match-play [26]. Average VO_2 during match-play in elite female players has been estimated at 77-80% of VO_{2max} [17,18], with peak values of 96% of VO_{2max} [18]. These values for heart rate and VO_2 suggest the aerobic energy system is highly taxed throughout female soccer match-play with periods of near maximal exertion exhibited [18]. It should be noted, however, that heart rate values during match-play might overestimate the actual physical demands [10,17,21].

Heart rate responses to match-play do not appear to vary for position during domestic competition with values of 86 - 88% of peak heart rate (HR_{peak}) reported for defenders, midfielders and attackers [18]. Slightly lower ($82 \pm 3\%$ HR_{peak}) (mean \pm SD) average heart rate values were reported during an international friendly [27]. Peak heart rate values achieved during match-play have been shown to be similar for international ($97 \pm 3\%$ HR_{peak}) and domestic games ($97 \pm 2\%$ HR_{peak}) (mean \pm SD), despite differences in workload as measured by high-speed running [9]. Heart rate values in females are reported as similar during the first and second halves or indeed in any 15-minute period of the game [21]. Furthermore, match-play heart rate was not found to correlate with VO_{2max} , which is likely due to the limitations of heart rate for assessing exercise-intensity during match-play [10,17,21].

3.2) Anaerobic

Still little information exists on the anaerobic energy contribution to female match-play. Traditionally anaerobic capacity has been evaluated using blood lactate with elite female players exhibiting mean blood lactate concentrations of 5.1 ± 0.5 $mmol.L^{-1}$ and 2.7 ± 0.4 $mmol.L^{-1}$ (mean \pm SD) following the first and second halves respectively [21]. Decreased blood lactate concentrations following the second half have been attributed to a reduction in distance covered and intensity [28]. However, it is important to highlight that blood lactate concentration depends largely on the activity pattern of the player in the minutes preceding blood sampling and consequently may over or underestimate the overall demands [28]. Further insight into the anaerobic demands of female match-play is warranted, particularly in relation to playing position and standards of competition.

4) Physical Capacity

Identifying the physical capacity of an individual player represents a central component of the ergonomic model discussed by Reilly [3]. A comprehensive understanding of a player's strengths and weaknesses informs selection, tactical decisions and ultimately the specifics of a detailed training programme. Moreover, the testing of physical capacity can help identify talent and differentiate between standards of competition [29].

4.1) Anthropometry

Female national team players and those competing in the highest domestic leagues are on average 20 - 27 years old, 1.61 – 1.70 m in stature, weighing 57 – 65 kg, with a body fat percentage of 14.6 - 20.1% and lean muscle mass of 45.7 ± 3.9 kg (mean \pm SD) [10,13,17,18,21,27,30-36] (see Table 2). Height and body mass of elite female players remains comparable to earlier reports, however, percentage body fat is now lower than previously observed (20.8 - 22%) [1]. These alterations in body composition likely reflect greater training volume and improved access to support staff such as nutritionists.

The large variation in anthropometric measures highlights the heterogeneity amongst the elite female population. Height and body mass have previously been reported to be similar between playing positions in elite Norwegian [33] and North American [37] players. However, contradictory evidence exists showing that defenders tend to be taller than attackers and heavier than all other outfield positions [33,35]. Similarly, goalkeepers have been shown to be taller and heavier than midfielders and attackers [33,35]. Goalkeepers were reported to have the highest fat mass, which could be attributed to reduced energy expenditure in training and games [35]. Full backs were found to have the lowest body fat percentage and highest muscle mass [35] which may be related to differing physical demands of their playing position although this finding cannot be corroborated as current match analysis literature fails to differentiate between wide and central defenders.

Conversely, older studies [38,39] have found no significant differences in body fat percentage between outfield playing positions. It can be speculated that the positional differences demonstrated in recent studies are a result of increased training specificity for playing positions or indeed that player's with particular anthropometric characteristics are assigned to certain playing positions.

The basic physical characteristics of female players across a spectrum of playing standards have been documented within the literature. Researchers have described anthropometric characteristics of high school [40], university [37,41,42], lower division domestic league [32], higher division domestic league [21,33] and national team players [10,13]. However, the anthropometric profile of elite youth players has yet to be fully examined. Some information exists regarding differences in anthropometric variables based on the level of competition, with differences in body mass and body fat percentage being demonstrated between elite and non-elite players [35,38]. Specifically, elite Spanish players were found to be lighter and have lower percentage body fats than their non-elite counterparts [35].

4.2) Maximal Aerobic Power

The aerobic system is highly stressed during soccer match-play [28]. The direct measurement of VO_{2max} in female soccer players has been reported in a number of studies with values for elite players ranging from 49.4 - 57.6 ml.kg⁻¹.min⁻¹ and are similar to those reported in the previous review [1]. Some studies [37,43,44] have estimated VO_{2max} using the equation derived from the distance achieved in the multistage fitness test [45]. However, the predictive power of the multistage fitness test to estimate VO_{2max} in young women soccer players has recently been questioned [46]. A summary of direct and indirect values for VO_{2max} values in elite female soccer players is shown in Table 3.

Similar VO_{2max} levels have been observed between playing positions [33] although midfielders often record higher values than those in other positions [33]. When using the multistage fitness test to estimate VO_{2max} the findings are equivocal with some studies showing no differences in aerobic capacity between playing positions [37,38], whereas one study highlighted significant differences between midfielders and defenders [39]. The VO_{2max} of female players has been reported by a number of researchers at various levels of competition and descriptions for university [37,42,47] and lower division domestic league players [44] generally show a lower capacity than higher division domestic league [17,18] and national team players [48,49]. In addition, a positive correlation has been demonstrated between VO_{2max} and the amount of high-intensity running completed during a match [18]. However, when VO_{2max} for international and regional English players was directly compared, no significant differences were reported [38]. The failure to discriminate between standards of competition in this study may be a result of both groups being non-professional at this time and therefore the differentiation between weekly training volume and intensity may have been slight. Furthermore, the specificity of VO_{2max} for soccer performance has been questioned [50,51], consequently more intermittent based assessment tools are now commonly used.

4.3) High-Intensity Endurance Capacity

It has previously been described [1] that in order to be successful a player must have the capacity to recover rapidly from high-intensity exercise. The use of the Yo Yo Intermittent Recovery (YYIR) [53] and Yo Yo Intermittent Endurance tests (YYIE) [54] are now commonplace in field-based testing protocols for team sports. An overview of published literature can be found in Table 4.

Published data for national team female players shows distance of 1774 ± 532 m being attained on the YYIE2 test [54], however, non-published data reports distances of up to 2182 ± 89 (mean ± SD) [55]. There appears to be no published data for national team players using the Yo Yo Intermittent Recovery test level 1 (YYIR1) test, however, domestic players from the Danish and Spanish leagues cover 1379m (600 – 1960 m) [18] and 1224 ± 255 m (mean ± SD) [32], respectively. Some positional differences in YYIE2 have been identified [54], with senior wide midfielders covering more distance than central defenders and attackers, similarly, wide and central midfielders covered more distance than central defenders at under nineteen (U19) level. Moreover, large differences (48%) in YYIR1 performance have been demonstrated (1224 m versus 826 m) between first and second division female Spanish players [32]. These large differences can likely be attributed to female players in lower leagues operating at a recreational level with a lower training volume.

4.4) Speed

A plethora of studies [22,29,37,40-44,56,57] have undertaken speed testing with female soccer players, however, the use of different assessment protocols limits comparison. For example, the distance that a player stands before the first timing gate is variable with 0 cm [41,58] and 88 cm [21] being used in the literature. Nevertheless, Table 5 highlights the studies to date that have described speed characteristics of female players.

The previous review [1], could only reference one report of sprint time in elite players 3.31 ± 0.11 s (mean \pm SD) for 20 m by Australian national team players [43]. More recent findings have shown 20 m sprint times of 3.26 ± 0.06 s [57] with the same protocol and 3.17 ± 0.03 s [21] when using a greater distance between the start line and first timing gate (mean \pm SD).

Limited information exists on the differences in speed characteristics between playing positions. Vescovi and colleagues [37] evaluated differences in speed variables (10 - 40 yard sprints) and they concluded there were no significant differences between playing positions, despite a trend for defenders to be slower. It has been proposed that sprint performance can distinguish between standards of competition [24,29]. Selected players from trials for an American professional soccer league were between 0.5 and 0.8 km.h⁻¹ faster than their non-selected equivalents [24]. Similar findings were reported during an Australian talent identification project with selected players recording faster times over 5 m, 10 m and 20 m, respectively than the non-selected players [29].

4.5) Repeated Sprint Ability

Repeated sprint ability is acknowledged as an important aspect of team sports [15,22]. However, very few studies have reported assessments of this fitness characteristic in female soccer players. Furthermore, the published papers to date [44,47,58] have used different protocols making comparison difficult. One study [44] adopted a method that involved players completing a 34.2 m sprint course seven times with 25 seconds rest between sprints [59]. The reported mean sprint time to complete the course ranged from 8.79 - 9.13 seconds following a training intervention. Gabbett [58] designed a repeated sprint test specifically to reflect the demands of international female soccer. The test involved 6 repetitions of a 20 m maximal sprint, on a 15 second recovery cycle which represents the most extreme repeated-sprint demands of the international game [11]. The total sprint time was found to be reliable (1.5% typical error of measurement) and able to discriminate between elite and sub-elite players and thus this protocol could be used in future to advance research in this area.

4.6) Power and Strength

The assessment of jumping capability is an accepted functional measure of power in soccer players [28]. Furthermore, a significant relationship has been demonstrated between team success and average jump height and leg extension power, thus demonstrating the importance of this physical component for soccer-specific performance [60]. There are numerous reports within the literature detailing the power performance of female players across differing competition levels; university [37,41,42], high school [56,61], lower division domestic league [32,44], higher division domestic [20,21,35], national team youth [36] and senior national team players [11,36,43,49]. As with the other components of physical fitness previously described, the ability to compare data between studies is hindered by the different protocols and equipment adopted by researchers. A summary of results for vertical jump ability in elite players can be found in Table 6.

Playing position does not seem to be a distinguishing factor relating to performance in jumping assessments. However, trends are apparent with goalkeepers recording the lowest scores [35,37] and midfielders and forwards recording greater scores than defenders [37]. Vertical jump performance has been shown to differentiate between age groups [36,61]. Specifically, U19 Italian national team players produced higher countermovement jumps (CMJ's) and squat jumps (SJ's) than their under seventeen (U17) counterparts [36]. Additionally, an improvement in CMJ performance has been described until players reach an age of 15 - 16 and then a plateau in performance is shown until 21 years of age [61]. The data regarding competitive standard and jumping performance show no apparent differences between Spanish [35] or English [38] elite and sub-elite players.

Insightful recommendations have been made within the literature regarding the specifics of protocols that should be adopted for female players. It has been suggested that a drop jump (DJ) may not be appropriate as DJ values have been shown to be lower than CMJ scores despite the increased activation of muscle fibres resulting in increased force production [35]. Justifications for this finding include the increased technical requirement with a DJ compared to a CMJ, furthermore the CMJ has been described as being more similar to physical skills executed within soccer. Consequently, consideration should be given before including the DJ in a testing protocol [35]. Unilateral jump performance has a stronger correlation with sprint performance than bilateral jump performance and is a useful tool for assessing asymmetries, therefore the inclusion of unilateral jumps in a testing protocol is warranted [41]. Castagna and Castellini [36] have attempted to describe thresholds of acceptable vertical jump values for elite females using the data gathered from Italian national team players. The authors have suggested scores over 34.4 cm and 32.9 cm for CMJ and static jump should be regarded as superior vertical jump abilities. In addition, a CMJ of 29.8cm should be considered a threshold measure for discriminating between competitive levels in elite female players. This work could be expanded to include thresholds for different playing positions and age groups.

Investigations into muscle strength in female players are limited and the methods of data collection appear varied, thus making comparisons difficult. Limited recent [21] and historical [48] isokinetic data is available for national team players but the majority of research has been conducted using sub-elite and recreational players to examine injury risk. The scarcity of information available in this area is surprising, as female players have been regularly completing this component of training for many years.

5) Training

Effective conditioning programmes that carefully balance training and recovery are the cornerstone of athletic development. Moreover, in soccer, those that can integrate both physical and tactical/technical training are considered superior in enhancing the effectiveness of the available training time. The issue of managing a player's training load is important for the elite player to ensure optimal preparation for performance and potentially reduce the susceptibility to injury [4,62].

5.1) Training Interventions

As the aerobic system is highly stressed during a soccer game [28], many soccer training programmes serve to increase the ability to repeatedly perform high-intensity exercise [63]. The volume of female specific training literature is sparse but two recent studies [62,64] have shown increased VO_{2max} using interval training methodologies. Specifically, an 8-week mixed-intensity interval programme significantly improved VO_{2max} capacity ($49.69 \pm 1.15 \text{ mmol.kg}^{-1}$ to $62.13 \pm 0.96 \text{ mmol.kg}^{-1}$) (mean \pm SD) [64]. Interval training sessions lasted between 12 – 36 minutes with work durations of between 30 – 90 seconds, sessions were repeated three times per week. Similarly, a 12-week individualised interval training programme consisting of intervals of between 5 – 120 seconds increased mean values for VO_{2max} (13.2%), anaerobic threshold (16.8%) and 1-minute heart rate recovery (36.9%) [62]. However, the participants in these two studies were recreational players and therefore large increases in physical markers would be expected when adhering to an organised training regime. Currently no information is available regarding high-intensity training interventions for elite female players.

It is recognised that to be successful in soccer, an individual must be able to cover short distances quickly [65], as high-intensity activities are most often associated with decisive moments in a game [1]. To date, few studies [44,65,66] have examined components of speed training using a female soccer population. Polman [44] evaluated the use of the 'Speed, Agility, Quickness' (SAQ) method of training and demonstrated improved mean performances in sprint to fatigue (11.6%), 25-m sprint (4.4%), left and right side agility (4.5% and 4.0% respectively) and vertical and horizontal power tests (18.5% and 7.7% respectively) following 12 weeks of SAQ training compared to a control group. The concept of assisted and resisted sprint training has been assessed and these methods were found to be more affective than traditional sprint training [65]. Furthermore, assisted sprint training was more suitable for improving acceleration (≤ 13.7 m) and resisted sprint training for improving maximum velocity (≥ 13.7 m). Bartolini [66] highlighted that when completing overspeed training (2 maximal sprints) with an elastic cord, sprint times were lowest following the use of 30% body weight assistance over distances up to 15 yards. The studies to date have examined sub-elite and collegiate level players and have expressed benefits of various forms of speed training on physical testing values. As with the high-intensity endurance training there appears to be limited available research on the interventions regarding the elite player and this is an area requiring further examination.

Explosive strength is an important determinant of high-level performance in soccer and plyometric activity is cited as a suitable training modality for improving jumping performance and explosiveness [67]. Specifically, elite female players who completed 12-weeks of plyometric training three times per week had mean increases of 3.3 cm and 4.5 cm for CMJ and DJ respectively [67]. Furthermore, female players completing combined plyometric, resistive and anaerobic programmes twice per week also experienced improvements in time to fatigue during a simulated soccer running test [68], a reduction in mean 20-m sprint time (0.10 s) [68] and a significant decrease in the percentage of VO_{2peak} at a set running speed [69]. Plyometric training has also been shown to have a role in injury prevention for female soccer players by enhancing the functional joint stability in the lower extremity [70]. These studies highlight the potential advantages of including plyometric-based activity as part of a training programme, however, as with the aforementioned training techniques further investigation using elite players is required.

6) Female Differences

There are some physiological areas of research that are exclusive to females. A full appraisal of these topics is beyond the scope of this review and in some instances has already been conducted [71-76]. However, the importance of these variables to female soccer players and the potential impact on training, performance and injury risk should not be underestimated.

6.1) Menstrual Cycle

Women aged ~13 – 50 years, experience a circa-mensual rhythm termed the menstrual cycle [77]. Alterations in the cycle can occur and cause menstrual disturbances [78]. High training loads can have an adverse effect on the normal menstrual cycle. Young females with low body mass and body fat levels who undertake intense training may experience delayed menarche [77,79]. Similarly, menstruating athletes with high training loads, low body mass, low body fat levels and/or energy imbalance [77] may suffer secondary amenorrhea, although this is uncommon in soccer players.

Physiological responses may be influenced by the variations in endocrine hormones that occur according to menstrual cycle phase [77]. As women train and compete at all stages of their menstrual cycle the possible effect on performance should be considered. These effects have received a limited amount of research attention and the findings to date have been unclear. VO_{2max} appears to be largely unaffected by the phase of the menstrual cycle [80,81]. However, some studies have shown increased minute ventilation [39,82-84], heart rate [84-86] and rating of perceived exertion [84,85] during the luteal phase, which could be attributed to an increased core temperature (0.3 - 0.5 °C) during this phase [84]. Lactate threshold findings are equivocal with some reports of stable values throughout the cycle [87,88], whilst others describe lower blood lactate concentrations in the luteal phase during moderate-intensity activity [89]. Sprint performance as measured via power output seems unaffected by menstrual cycle phase [90]. Investigations into muscle contractile strength during the cycle have produced conflicting results, however, when methodological shortcomings are addressed it appears that fluctuations in female steroid hormones do not affect muscle strength and fatigability [73]. Limited soccer-specific research exists but one study showed no performance differences in high-intensity intermittent shuttle running throughout the menstrual cycle [91], although differences in core temperature were not found in this study which may explain these findings. At present there are a number of conflicting reports within the literature, methodological concerns and a lack of specificity to team sports, specifically soccer activity.

6.2) Oral Contraceptive Pill

The oral contraceptive pill (OCP) is available in single (progesterone only) and combined (oestrogen and progesterone) formulations. Non-contraceptive benefits include cycle control and reductions in dysmenorrhoea, iron deficient anaemia and bone mineral density loss [92]. In recent years there has been an increase in the use of OCP by the athletic population with reports stating 83% of elite athletes use an OCP [93].

Despite the increased prevalence of OCP usage in athletic populations there is still limited research regarding its effects on athletic performance [94]. Aerobic performance as measured by VO_{2max} has been shown to reduce by 5 - 15% in trained and active women using a triphasic OCP [95,96], however, research remains equivocal when considering the monophasic pill [72]. Anaerobic performance, high-intensity intermittent performance and strength appear to be largely unaffected by OCP, however, more well-controlled studies are required [72,93]. During an investigation into the effects of OCP on team sport athletes, reactive strength index as measured during a drop jump was significantly lower during the withdrawal phase, i.e. when exogenous oestrogen and progesterone were at their lowest. The mechanism for this finding could not be fully explained but it was suggested that increased endogenous oestrogen may have a negative impact on neuromuscular timing, muscle activation and performance [94]. It is clear that further studies are necessary to ascertain the effects of OCP on soccer-specific performance.

7) Injury

Injury is an accepted part of sports participation at all levels of competition. Variations in injury definition and methods of calculating incidence mean that it is difficult to make valid inter-study comparisons [97]. The majority of literature focuses on adult male professional players with only a few studies assessing injuries in female soccer players [98-105].

The injury incidence for females is between 1.2 and 7.0 injuries per 1000 training hours and 12.6 - 24.0 per 1000 match hours [103-107]. Longitudinal injury patterns in female soccer are of interest to assess if soft tissue injuries are becoming more frequent as the intensity of the female game increases or if the increased

professionalism and training status of players decreases injury rates. An 8-season domestic French study reported no significant differences between 1998 and 2006 with injury rates fluctuating between ~4.5 - 8.0 injuries per 1000 hours [105]. Conversely, recorded injuries in the U19 European Championships decreased between 2006 and 2008 from 13.5 to 4.9 injuries per 1000 hours [108].

Soccer injuries predominately affect the ankle, knee and muscles of the thigh and calf [101]. Injuries to the hip and groin are less common in female players [98,103,106,109,110] than males [111,112] and this is attributed to the abdominal wall deficiency apparent in males [113]. Knee injuries are more frequent in females than males [114], with the knee [98,107] and ankle [110] often being cited as the most common injury sites in females. Incidences of ankle injuries have been shown as similar between elite males and females during a competitive season in the Swedish Premier League [114]. Furthermore, ankle sprains accounted for 40% of all injuries in an 8-season study in to female youth soccer [105].

Knee injuries in females are well documented [103] with ligament injuries, specifically injury to the anterior cruciate ligament (ACL) predominant in the literature [75,98,99,115,116]. The large body of evidence is in part due to the financial and emotional cost of an ACL as well as the potential impact upon the team [117]. Female athletes are reported to have between 2 and 6 fold higher incidence of non-contact ACL injury than their male counterparts [115,118]. In soccer, this has been translated to ACL injury rates of 0.4 and 0.7 injuries per season in males and females respectively [115]. Match-play poses a substantially higher risk of ACL injury compared to training (0.09 per 1000 hours versus 0.90 per 1000 hours respectively) [107]. In addition, female players sustain their knee injuries at an earlier age (14 - 19 years) [75] compared to males (~23 years) [120]. The majority of ACL injuries in soccer are non-contact and occur during landing and cutting manoeuvres [120]. Discrete classification is not always possible as ACL injuries often involve perturbation by another player [120]. The risk factors associated with ACL injury are often multi-factorial. External risk factors include environmental conditions, footwear and surface. Internal risk factors include anatomical, neuromuscular, biomechanical and hormonal variables [74,75]. Studies often concentrate on intrinsic risk factors as they can provide differentiation between the sexes.

Evidence suggests the anatomy of the female knee, specifically a smaller intercondylar notch width and ACL, are contributing factors to an increased risk of ACL injury [74,75]. A link has been established between hypermobile individuals and higher risk of ACL injuries in female soccer and basketball players [122]. A greater quadriceps angle (Q-angle) has been linked to increased injury risk in female basketball players [123] but this has not been replicated in elite female soccer players [110]. Whilst it is important to recognise these anatomical factors they should not be focussed upon as they cannot be addressed via training interventions [74]. Specific studies assessing differences between the sexes have highlighted dynamic alterations in biomechanical loading and patterns of neuromuscular activation, which may expose females to a higher predisposition to injury [124]. Knee stability during dynamic movement and landing is underpinned by muscular strength and recruitment patterns [124]. Hamstring activation patterns and knee abduction movements differ between males and females, with females exhibiting slower and less frequent contractions and increased knee valgus moments [118,125]. Specifically, female soccer players land with decreased hip and knee flexion compared to males, thus increasing force transmitted through the quadriceps relative to the hamstrings and consequently stressing the ACL [126]. Alterations in neuromuscular and biomechanical strategies may be present under fatigue [127] and contribute to increased risk of injury but cannot be considered as an isolated risk factor for ACL [128].

A recent review combined data from seven studies and concluded that female athletes may be more predisposed to non-contact ACL injuries during the pre-ovulatory phase of the menstrual cycle [129]. It has been suggested that hormonal fluctuations may have an indirect effect on neuromuscular control and consequently injury [129]. Furthermore, OCP use combined with neuromuscular training may be advantageous for increasing the dynamic stability of the knee and thus reducing injury risk [130-132]. Indirect support also exists for a link between the use of OCP and reduced soft tissue [133] and traumatic injuries [134].

8) Conclusions and Future Research

Women's soccer has changed dramatically since the previous review [1] two decades ago. The increased professionalism within the sport is represented by the increased physicality of match-play as total distances covered have increased from ~8.5 km [1] to ~10 km [10-13]. Differences in time-motion analysis have also been identified between standards of competition [11], playing position [11,13,14] and at different time-points within a match [12,14,17,18]. The majority of time-motion studies in female soccer to date have used traditional single camera analysis methods as such the ability to examine specific physical elements in detail are limited [5,22]. Consequently, more comprehensive evaluations into the physical characteristics of elite female

match-play using contemporary analysis methods and larger sample sizes are warranted.

The physical capacity of female players is probably the most thoroughly researched area within both this and also the previous review [1]. Data is available for most components of physical fitness; however, research focused upon the influence of competitive standard, age and playing position on performance characteristics is still lacking. This largely reflects a lack of consistency between protocols and equipment used. Data for the elite female player is lacking for YYIR and for repeated sprint activity and as such warrants further investigation. Other areas of investigation include tracking physical capacity and injury rates over time to examine potential relationships as well as the utilisation of dual-energy x-ray absorptiometry (DXA) as a sophisticated measurement of body composition to better understand lean mass.

There is still equivocal research on how the menstrual cycle and/or the usage of OCP may affect physical performance. Some studies have indicated that physiological fluctuations that occur throughout the menstrual cycle and/or as a result of OCP usage may affect physical performance [84,93,95]. One study has revealed that the use of the OCP (during the withdrawal phase) has been shown to elicit significantly lower reactive strength index scores during a drop jump. Injury data in female players has largely focussed on ACL injuries, as it is a recognised issue within female team sport athletes, with the incidence quoted as 2-6 times greater compared to males [115,118]. Numerous explanations exist for this increased incidence, some of which are anatomical [74,75,123], whilst others focus on dynamic differences in biomechanical loading and patterns of neuromuscular activation [118,124,126]. A key recent finding is that female athletes may be more predisposed to non-contact ACL injuries during the pre-ovulatory phase of the menstrual cycle [129] and therefore the advantages and disadvantages for training and competing during this phase should be considered.

Acknowledgements

No sources of funding were used to assist in the preparation of this review. The authors have no potential conflicts of interest that are directly relevant to the content of this review.

Table 1. Match physical demands for elite female soccer players

| Reference | Year | N | Competition level | Nationality / region | Distance covered (km) | Number of activity changes | Number of high intensity bouts | High intensity running (km) | High intensity % | Sprinting % | Sprint distance (m) (>25km.h ⁻¹) | Number of sprinting bouts (>25km.h ⁻¹) | Method of data capture |
|-------------------------|------|----|--------------------|----------------------|-----------------------|----------------------------|--------------------------------|-----------------------------|------------------|-------------|--|--|------------------------|
| Andersson et al. [10] | 2010 | 17 | International | Scandinavian | 9.9 ± 1.8 | 1641 ± 41 | 187 ± 15 | 1.53 ± 0.1 | | 0.65 ± 0.06 | 256 ± 57 | | Video tape |
| | | 17 | Domestic league | Scandinavian | 9.7 ± 1.4 | 1593 ± 30 | 168 ± 12 | 1.33 ± 0.9 | | 0.54 ± 0.05 | 221 ± 45 | | Video tape |
| Andersson et al. [135] | 2008 | 21 | Highest division | Swedish | 9.9 | | | 1.15 | | | | | Video tape |
| Gabbett and Mulvey [11] | 2008 | 13 | v male youth teams | Australian | 9.32 ± 0.84 | | | 1.86 ± 0.48 ^a | | | | | Video tape |
| | | 13 | National league | Australian | 9.70 ± 0.48 | | | 2.01 ± 0.30 ^a | | | | | Video tape |
| | | 13 | International | Australian | 9.97 ± 1.14 | | | 2.46 ± 0.49 ^a | | | | | Video tape |
| Hewitt et al. [13] | 2008 | 15 | International | Australian | 9.14 ± 1.03 | | | 0.62 ± 0.11 ^b | | | | 280 ± 80 ^c | GPS |
| Krustrup et al. [12] | 2008 | 12 | International | Scandinavian | 10.0 ± 0.5 | | | 1.6 ± 0.4 | | | | | Video tape |
| | | 12 | Domestic league | Scandinavian | 9.7 ± 0.6 | | | 1.4 ± 0.4 | | | | | Video tape |
| Mohr et al. [14] | 2008 | 19 | Highest division | All – USA pro league | 10.33 ± 0.15 | 1379 ± 34 | 154 ± 7 | 1.68 ± 0.09 | 6.0 ± 0.3 | 1.2 ± 0.1 | 460 ± 20 | 30 ± 2 | Video tape |
| | | 15 | Domestic league | Scandinavian | 10.44 ± 0.15 | 1326 ± 24 | 125 ± 7 | 1.3 ± 0.10 | 4.4 ± 0.5 | 0.9 ± 0.1 | 380 ± 50 | 26 ± 1 | Video tape |
| Krustrup et al. [18] | 2005 | 14 | Highest division | Danish | 10.3 | 1459 | 125 | 1.31 | | | 160 | 26 | Video tape |

Data are mean ± standard deviation

GPS = global positioning system

^a = descriptive definition^b = >16km.h⁻¹^c = >20km.h⁻¹

Table 2. Basic physical characteristics for elite female soccer players

| Reference | Year | N | Competition level | Nationality / region | Age (yrs) | Height (m) | Weight (kg) | Body fat content (%) |
|--------------------------|------|-----|-----------------------|----------------------|--------------|-------------|--------------|----------------------|
| Castagna et al. [36] | 2013 | 21 | National team players | Italian | 25.8 ± 3.9 | 1.67 ± 0.04 | 59.9 ± 3.8 | |
| Gravina et al. [31] | 2011 | 14 | Highest division | Spanish | 25 ± 5 | | 61 ± 7.4 | 15.5 ± 2.9 |
| Ingebrigsten et al. [33] | 2011 | 29 | Highest division | Norwegian | 20.8 ± 3.7 | 1.66 ± 0.05 | 60.7 ± 6.6 | |
| Andersson et al. [10] | 2010 | 17 | National team players | Scandinavian | 27 ± 1 | 1.68 ± 0.02 | 61.0 ± 1.4 | |
| Andersson et al. [27] | 2010 | 16 | Highest division | Scandinavian | 22 ± 3 | 1.67 ± 0.05 | 64 ± 2 | |
| Krustrup et al. [21] | 2010 | 23 | Highest division | Danish | 23 | 1.69 | 60.1 | 18.50 |
| Mujika et al. [32] | 2009 | 17 | Highest division | Spanish | 23.1 ± 2.9 | 1.65 ± 0.04 | 56.8 ± 5.7 | |
| Sedano et al. [35] | 2009 | 100 | Highest division | Spanish | 22.1 ± 1.1 | 1.61 ± 0.06 | 57.7 ± 7.5 | 20.1 ± 5.5 |
| Hewitt et al. [13] | 2008 | 15 | National team players | Australian | 23.5 ± 2.5 | 1.70 ± 0.05 | 64.88 ± 4.6 | |
| Krustrup et al. [18] | 2005 | 14 | Highest division | Danish | 24 | 1.67 | 58.5 | 14.6 |
| Can et al. [34] | 2004 | 17 | Highest division | Turkish | 20.73 ± 2.09 | 1.62 ± 0.06 | 56.63 ± 5.03 | 19.75 ± 0.69 |
| Bunc et al. [30] | 2004 | 14 | Elite | Czech | 25.3 ± 4.8 | 1.65 ± 0.06 | 64.5 ± 9.9 | 14.9 ± 5.7 |
| Mohr et al. [17] | 2004 | 18 | Highest division | Scandinavian | 24 ± 1 | 1.68 ± 0.02 | 61 ± 1 | 15 ± 1 |

Data are mean ± standard deviation

Table 3. Maximal oxygen uptake for elite female soccer players

| Reference | Year | N | Competition level | Nationality / region | Maximal oxygen uptake (ml.kg.min ⁻¹) | Estimated maximal oxygen uptake (ml.kg.min ⁻¹) |
|--------------------------|------|----|-----------------------|----------------------|--|--|
| Ingebrigsten et al. [33] | 2011 | 29 | Highest division | Norwegian | 51.85 ± 5.05 (def) | |
| | | | | | 55.36 ± 5.65 (mid) | |
| | | | | | 52.94 ± 3.17 (att) | |
| | | | | | 50.70 ± 4.96 (gk) | |
| Andersson et al. [27] | 2010 | 16 | Highest division | Scandinavian | 54 ± 3 | |
| Krustrup et al. [21] | 2010 | 23 | Highest division | Danish | 52.3 ± 1.3 | |
| Andersson et al. [20] | 2008 | 8 | Highest division | Scandinavian | 55.4 ± 3.6 | |
| Andersson et al. [20] | 2008 | 9 | Highest division | Scandinavian | 53.8 ± 2.3 | |
| Gabbett et al. [11] | 2008 | 13 | Scholarship holders | Australian | 51.4 ± 5.4 | |
| Krustrup et al. [18] | 2005 | 14 | Highest division | Danish | 49.4 | |
| Bunc et al. [30] | 2004 | 14 | Elite | Czech | 53.9 ± 5.7 | |
| Mohr et al. [16] | 2004 | 18 | Highest division | Scandinavian | 49.5 ± 1 | |
| Tumilty et al. [57] | 2000 | 17 | National team players | Australian | | 50.3 ± 5.1 |
| Tamer et al. [136] | 1997 | 22 | Elite | Turkish | | 43.15 ± 4.06 |
| Davis et al. [48] | 1992 | 14 | National team players | English | 52.2 ± 5.1 ^a | |
| Evangelista et al. [137] | 1992 | 12 | Elite | Italian | 49.75 ± 8.3 | |
| Jensen et al. [49] | 1992 | 10 | National team players | Danish | 57.6 ^a | |
| Tumilty et al. [43] | 1992 | 20 | National team players | Australian | | 48.5 ± 4.8 |

Data are mean ± standard deviation

^a = following an intervention

def = defenders

mid = midfielders

att = attackers

gk = goalkeepers

Table 4. Soccer specific endurance test results for elite female soccer players

| Reference | Year | N | Competition level | Nationality / region | YYIR1 (m) | YYIE2 (m) |
|----------------------|------|----|---------------------------|----------------------|------------|------------|
| Bradley et al. [54] | 2012 | 92 | National team players | European | | 1774 ± 532 |
| | | 46 | Highest division | European | | 1261 ± 449 |
| | | 42 | U20 national team players | European | | 1490 ± 447 |
| | | 19 | Lowest division | European | | 994 ± 373 |
| Krustrup et al. [21] | 2010 | 23 | Highest division | Danish | | 1213 ± 90 |
| Mujika et al. [32] | 2009 | 17 | Highest division | Spanish | 1224 ± 255 | |
| | | 17 | Second division | Spanish | 826 ± 160 | |
| Krustrup et al. [18] | 2005 | 14 | Highest division | Danish | 1379 | |

Data are mean ± standard deviation

YYIR1 = Yo-yo intermittent recovery test level 1

YYIE2 = Yo-yo intermittent endurance test level 2

U20 = Under 20

Table 5. Speed test results for female soccer players

| Reference | Year | N | Competition level | Nationality / region | Time 5m (s) | Flying time 9.14m (s) | Time 9.14m (s) | Time 10m (s) | Flying time 18.2m (s) | Time 18.28m (s) | Flying time 20m (s) | Time 20m (s) | Time 25m (s) | Time 27.3m (s) | Time 36.56m (s) |
|-----------------------|------|----|---|----------------------|-------------|-----------------------|----------------|--------------|-----------------------|-----------------|--------------------------|--------------|--------------------------|----------------|-----------------|
| Sjokvist et al. [42] | 2011 | 14 | University | USA | | | | | | | | 3.59 ± 0.17 | | | |
| McCurdy et al. [41] | 2010 | 15 | University | USA | | | | 2.31 ± 0.25 | | | | | 4.52 ± 0.20 | | |
| Andersson et al. [20] | 2008 | 8 | Highest division | Scandinavian | | | | | | | | 3.18 ± 0.03 | | | |
| | | 9 | Highest division | Scandinavian | | | | | | | | 3.17 ± 0.03 | | | |
| Vescovi et al. [56] | 2008 | 83 | High school | USA | | 1.37 ± 0.06 | 1.96 ± 0.10 | | 2.67 ± 0.13 | 3.33 ± 0.15 | | | | 4.63 ± 0.21 | 5.94 ± 0.28 |
| | | 51 | College | USA | | 1.38 ± 0.07 | 2.00 ± 0.11 | | 2.75 ± 0.44 | 3.38 ± 0.17 | | | | 4.69 ± 0.23 | 5.99 ± 0.29 |
| Vescovi et al. [37] | 2006 | 64 | University | USA | | | 1.98 ± 0.11 | | | 3.34 ± 0.17 | | | | | 5.90 ± 0.31 |
| Polman et al. [44] | 2004 | 36 | Second division | English | | | | | | | | | 4.12 ± 0.13 ^a | | |
| Siegler et al. [40] | 2003 | 17 | High school | USA | | | | | | | 2.90 ± 0.13 ^a | | | | |
| Hoare et al. [29] | 2000 | 17 | Talent identification – selected triallists | Australian | 1.18 ± 0.06 | | | 2.01 ± 0.08 | | | | 3.47 ± 0.14 | | | |
| Tumilty [57] | 2000 | 20 | National team players | Australian | 1.14 ± 0.04 | | | 1.91 ± 0.04 | | | | 3.26 ± 0.06 | | | |

Data are mean ± standard deviation

^a = following an intervention

Table 6. Vertical jump test results for elite female soccer players

| Reference | Year | N | Competition level | Nationality | Squat jump (no arms) (cm) | Drop jump (no arms) (cm) | Countermovement jump (no arms) (cm) | Countermovement jump (with arms) (cm) |
|-----------------------|------|-----|-----------------------|--------------|---------------------------|--------------------------|-------------------------------------|---------------------------------------|
| Castagna et al. [36] | 2013 | 21 | National team players | Italian | 30.1 ± 3.7 | | 31.6 ± 4.0 | |
| | | 20 | U19 national team | Italian | 32.8 ± 2.9 | | 34.3 ± 3.9 | |
| | | 21 | U17 national team | Italian | 28.2 ± 2.5 | | 29 ± 2.1 | |
| Krustrup et al. [21] | 2010 | 23 | Highest division | Danish | | | 35 ± 1.0 | |
| Mujika et al. [32] | 2009 | 17 | Highest division | Spanish | | | 32.6 ± 3.7 | 38 ± 4.8 |
| Sedano et al. [35] | 2009 | 100 | Highest division | Spanish | | 25.3 ± 5.6 | 26.1 ± 4.8 | |
| Andersson et al. [20] | 2008 | 8 | Highest division | Scandinavian | | | 30.5 ± 1.2 | |
| Krustrup et al. [13] | 2008 | 12 | National team players | Scandinavian | | | 35 ± 1.0 | |
| Can et al. [34] | 2004 | 17 | Highest division | Turkish | | | 34.48 ± 7.11 ^a | |
| Tumilty [57] | 2000 | 20 | National team players | Australian | | | 51 ± 5 | |
| Jensen et al. [49] | 1992 | 10 | National team players | Danish | | | 37.8 ^{b,c} | |
| Tumilty et al. [43] | 1992 | 20 | National team players | Australian | | | 40.5 ± 4.5 ^a | |

Data are mean ± standard deviation

^a = recorded as "vertical jump" but assumed to be countermovement jump

^b = jump height estimated from flying time

^c = following an intervention

U19 = under 19

U17 = under 17

References

1. Davis JA, Brewer J. Applied physiology of female soccer players. *Sports Med.* 1993;16(3):180-189.
2. Fahmy M. Increase participation and competitions. In: 5th FIFA Women's Football Symposium. FIFA. 2011.
http://www.fifa.com/mm/document/footballdevelopment/women/01/51/51/64/presentation_increasparticipation_e.pdf. Accessed 19 Aug 2013.
3. Reilly T. An ergonomics model of the soccer training process. *J Sports Sci.* 2005;23(6):561–72.
4. Randers M, Mujika I, Hewitt A, et al. Application of four different match analysis systems: A comparative study. *J Sports Sci.* 2010;28(2):171-182.
5. Di Salvo V, Gregson W, Atkinson G, et al. Analysis of high intensity activity in Premier League soccer. *Int J Sports Med.* 2009;30:205–12.
6. Harley JA, Lovell RJ, Barnes CA, et al. The interchangeability of global positioning system and semiautomated video-based performance data during elite soccer match play. *J Strength Cond Res.* 2011;25(8):2334–6.
7. Bradley PS, Sheldon W, Wooster B, et al. High-intensity running in English FA Premier League soccer matches. *J Sports Sci.* 2009;27(2):159–68.
8. Di Salvo V, Baron R, González-Haro C, et al. Sprinting analysis of elite soccer players during European Champions League and UEFA Cup matches. *J Sports Sci.* 2010;28(14):1489–94.
9. Vescovi JD. Sprint profile of professional female soccer players during competitive matches: Female Athletes in Motion (FAiM) study. *J Sports Sci.* 2012;30(12):1259–65.
10. Andersson HÅ, Randers MB, Heiner-Møller A, et al. Elite female soccer players perform more high-intensity running when playing in international games compared with domestic league games. *J Strength Cond Res.* 2010;24(4):912-919.
11. Gabbett TJ, Mulvey MJ. Time-motion analysis of small-sided training games and competition in elite women soccer players. *J Strength Cond Res.* 2008;22(2):543-552.
12. Krstrup P, Andersson H, Mohr M, et al. Match activities and fatigue development of elite female soccer players at different levels of competition. In: Reilly T and Korkusuz F editors. *Science and Football VI*. New York: Routledge; 2008. pp.205-211.
13. Hewitt A, Withers R, Lyons K. Match analyses of Australian international female soccer players using an athlete tracking device. In: Reilly T and Korkusuz F editors. *Science and Football VI*. New York: Routledge; 2008. pp.224-228.
14. Mohr M, Krstrup P, Andersson H, et al. Match activities of elite women soccer players at different performance levels. *J Strength Cond Res.* 2008;22(2):341–9.
15. Gabbett TJ, Wig, H, Spencer, M. Repeated high-intensity running and sprinting in elite women's soccer competition. *Int J Sports Physiol Perform.* 2013;8(2):130-8.
16. Mohr M, Krstrup P, Bangsbo J. Match performance of high-standard soccer players with special reference to development of fatigue. *J Sports Sci.* 2003;21(7):519–28.
17. Mohr M, Ellingsgaard H, Andersson H, et al. Physical demands in high-level female soccer - Applications of fitness tests to evaluate match performance. *J Sports Sci.* 2004;22(6):552–3.
18. Krstrup P, Mohr M, Ellingsgaard H, et al. Physical demands during an elite female soccer game: importance of training status. *Med Sci Sports Exerc.* 2005;37(7):1242–8.

19. Gregson W, Drust B, Atkinson G, et al. Match-to-match variability of high-speed activities in Premier League soccer. *Int J Sports Med.* 2010;31(04):237–42.
20. Andersson H, Raastad T, Nilsson J, et al. Neuromuscular fatigue and recovery in elite female soccer. *Med Sci Sports Exerc.* 2008;40(2):372–80.
21. Krustup P, Zebis M, Jensen JM, et al. Game-induced fatigue patterns in elite female soccer. *J Strength Cond Res.* 2010;24(2):437-441.
22. Spencer M, Bishop D, Dawson B, et al. Physiological and metabolic responses of repeated-sprint activities: specific to field-based team sports. *Sports Med.* 2005;35(12):1025–44.
23. Abt G and Lovell R. The use of individualized speed and intensity thresholds for determining the distance run at high-intensity in professional soccer. *J Sports Sci.* 2009;27(9):893–98.
24. Vescovi JD. Sprint speed characteristics of high-level American female soccer players: Female Athletes in Motion (FAiM) Study. *J Sci Med Sport.* 2012;15(5):474–8.
25. Helgerud J, Engen LC, Wisloff U, et al. Aerobic endurance training improves soccer performance. *Med Sci Sports Exerc.* 2001;33(11):1925–31.
26. Reilly T. Energetics of high-intensity exercise (soccer) with particular reference to fatigue. *J Sports Sci.* 1997;15(3):257–63.
27. Andersson H, Karlsen A, Blomhoff R, et al. Plasma antioxidant responses and oxidative stress following a soccer game in elite female players. *Scand J Med Sci Sports.* 2010;20(4):600–8.
28. Stolen T, Chamari K, Castagna C, et al. Physiology of soccer: an update. *Sports Med.* 2005;35(6):501–36.
29. G Hoare D, Warr CR. Talent identification and women's soccer: An Australian experience. *J Sports Sci.* 2000;18(9):751–8.
30. Bunc V, Psotta R. Functional characteristics of elite Czech female soccer players. *J Sports Sci.* 2004;22(6):528.
31. Gravina L, Ruiz F, Lekue JA, et al. Metabolic impact of a soccer match on female players. *J Sports Sci.* 2011;29(12):1345–52.
32. Mujika I, Santisteban J, Impellizzeri FM, et al. Fitness determinants of success in men's and women's football. *J Sports Sci.* 2009;27(2):107–14.
33. Ingebrigtsen J, Dillern T, Shalfawi SAI. Aerobic capacities and anthropometric characteristics of elite female soccer players. *J Strength Cond Res.* 2011;25(12):3352-3357.
34. Can F, Yilmaz I, Erden Z. Morphological characteristics and performance variables of women soccer players. *J Strength Cond Res.* 2004;18(3):480–5.
35. Sedano S, Vaeyens R, Philippaerts RM, et al. Anthropometric and anaerobic fitness profile of elite and non-elite female soccer players. *J Sports Med Phys Fitness.* 2009;49(4):387–94.
36. Castagna C, Castellini E. Vertical jump performance in Italian male and female national teams soccer players. *J Strength Cond Res.* 2013;27(4):1156–61.
37. Vescovi JD, Brown TD, Murray TM. Positional characteristics of physical performance in Division I college female soccer players. *J Sports Med Phys Fitness.* 2006;46(2):221–6.
38. Todd MK, Scott D, Chisnall PJ. Fitness characteristics of English female soccer players: an analysis by position and playing standard. In: Spinks W, Reilly T, Murphy A, editors. New York: Routledge:

- Science and Football IV*; 2002. pp. 374–81.
39. Wells C, Reilly T. Influence of playing position on fitness and performance measures in female soccer players. In: Spinks W, Reilly T, Murphy A, editors. New York: Routledge; 2002. pp. 369–73.
 40. Siegler J, Gaskill S, Ruby B. Changes evaluated in soccer-specific power endurance either with or without a 10-week, in-season, intermittent, high-intensity training protocol. *J Strength Cond Res.* 2003;17(2):379–87.
 41. McCurdy KW, Walker JL, Langford GA, et al. The relationship between kinematic determinants of jump and sprint performance in division I women soccer players. *J Strength Cond Res.* 2010;24(12):3200–3208.
 42. Sjökvist J, Laurent MC, Richardson M, et al. Recovery from high-intensity training sessions in female soccer players. *J Strength Cond Res.* 2011;25(6):1726–35.
 43. Tumilty D, Darby S. Physiological characteristics of Australian female soccer players. *J Sports Sci.* 1992;10:145.
 44. Polman R, Walsh D, Bloomfield J, et al. Effective conditioning of female soccer players. *J Sports Sci.* 2004;22(2):191–203.
 45. Ramsbottom R, Brewer J, Williams C. A progressive shuttle run test to estimate maximal oxygen uptake. *Br J Sports Med.* 1988;22(4):141–4.
 46. Castagna C, Impellizzeri FM, Manzi V, et al. The assessment of maximal aerobic power with the multistage fitness test in young women soccer players. *J Strength Cond Res.* 2010;24(6):1488–1494.
 47. Rhodes E, Mosher R. Aerobic and anaerobic characteristics of elite female university soccer players. *J Sports Sci.* 1992;10:143–4.
 48. Davis JA, Brewer J. Physiological characteristics of an international female soccer squad. *J Sports Sci.* 1992;10:142–3.
 49. Jensen K, Larsson B. Variations in physical capacity among the Danish national soccer team for women during a period of supplemental training. *J Sports Sci.* 1992;10:144.
 50. Castagna C, Abt G, D'Ottavio S. Competitive-level differences in Yo-Yo intermittent recovery and twelve minute run test performance in soccer referees. *J Strength Cond Res.* 2005;19(4):805–9.
 51. Krustup P, Bangsbo J. Physiological demands of top-class soccer refereeing in relation to physical capacity: effect of intense intermittent exercise training. *J Sports Sci.* 2001;19(11):881–91.
 52. Bangsbo J, Iaia FM, Krustup P. The Yo-Yo intermittent recovery test : a useful tool for evaluation of physical performance in intermittent sports. *Sports Med.* 2008;38(1):37–51.
 53. Castagna C, Impellizzeri FM, Belardinelli R, et al. Cardiorespiratory responses to yo-yo intermittent endurance test in nonelite youth soccer players. *J Strength Cond Res.* 2006;20(2):326–30.
 54. Bradley PS, Bendiksen M, Dellal A, et al. The application of the yo-yo intermittent endurance level 2 test to elite female soccer populations. *Scand J Med Sci Sports.* 2014;24(1):43–54.
 55. Scott D and Andersson H. Women Soccer. In: Williams M, editor. *Science and Soccer: Developing Elite Players.* London: Routledge; 2012. pp. 237–257.
 56. Vescovi JD, Mcguigan MR. Relationships between sprinting, agility, and jump ability in female athletes. *J Sports Sci.* 2008;26(1):97–107.
 57. Tumilty, D. Protocols for the physiological assessment of male and female soccer players. In: Gore, CJ, editors. *Physiological tests for elite athletes.* Champaign, Illinois: Human Kinetics; 2000. pp. 356–

- 362.
58. Gabbett TJ. The development of a test of repeated-sprint ability for elite women's soccer players. *J Strength Cond Res.* 2010;24(5):1191-1194.
 59. Bangsbo J. The physiology of soccer-with special reference to intense intermittent exercise. *Acta Physiol Scand.* 1994;619:1–155.
 60. Arnason A, Sigurdsson SB, Gudmundsson A, et al. Physical fitness, injuries and team performance in soccer. *Med Sci Sports Exerc.* 2001;36(2):278–85.
 61. Vescovi JD, Rupf R, Brown TD, et al. Physical performance characteristics of high-level female soccer players 12-21 years of age. *Scand J Med Sci Sports.* 2011;21(5):670–8.
 62. Rhea MR, Lavinge DM, Robbins P, et al. Metabolic conditioning among soccer players. *J Strength Cond Res.* 2009;23(3):800–6.
 63. Little T, Williams AG. Suitability of soccer training drills for endurance training. *J Strength Cond Res.* 2006;20(2):316–9.
 64. Clark JE. The use of an 8-week mixed-intensity interval endurance-training program improves the aerobic fitness of female soccer players. *J Strength Cond Res.* 2010;24(7):1773–81.
 65. Upton DE. The effect of assisted and resisted sprint training on acceleration and velocity in Division I female soccer athletes. *J Strength Cond Res.* 2011;25(10):2645–52.
 66. Bartolini JA, Brown LE, Coburn JW, et al. Optimal elastic cord assistance for sprinting in collegiate women soccer players. *J Strength Cond Res.* 2011;25(5):1263-1270.
 67. Campo SS, Vaeyens R, Philippaerts RM, et al. Effects of lower-limb plyometric training on body composition, explosive strength and kicking speed in female soccer players. *J Strength Cond Res.* 2009;23(6):1714-22.
 68. Siegler J, Gaskill S, Ruby R. Changes evaluated in soccer-specific power endurance either with or without a 10-week, in-season, intermittent, high-intensity training protocol. *J Strength Cond Res.* 2003;17(2):379-87.
 69. Grieco CR, Cortes N, Greska EK, et al. Effects of a combined resistance-plyometric training program on muscular strength, running economy, and $VO_{2\text{ peak}}$ in division 1 female soccer players. *J Strength Cond Res.* 2012;26(9):2570-6.
 70. Chimera NJ, Swanik KA, Swanik CB, et al. Effects of plyometric training on muscle-activation strategies and performance in female athletes. *J Athl Train.* 2004;39(1):24-31.
 71. Oosthuysen T, Bosch AN. The effect of the menstrual cycle on exercise metabolism: implications for exercise performance in eumenorrhoeic women. *Sports Med.* 2010;40(3):207–27.
 72. Burrows M, Peters CE. The influence of oral contraceptives on athletic performance in female athletes. *Sports Med.* 2007;37(7):557–74.
 73. Janse de Jonge XAK. Effects of the menstrual cycle on exercise performance. *Sports Med.* 2003;33(11):833–51.
 74. Alentorn-Geli E, Myer GD, Silvers HJ, et al. Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc.* 2009;17(7):705–29.
 75. Renstrom P, Ljungqvist A, Arendt E, et al. Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. *Br J Sports Med.* 2008;42(6):394–412.

76. Vescovi JD. The menstrual cycle and anterior cruciate ligament injury risk: implications of menstrual cycle variability. *Sports Med.* 2011;41(2):91–101.
77. Reilly T. The menstrual cycle and human performance: an overview. *Biol Rhythm Res.* 2000;31(1):29–40.
78. Loucks AB. Effects of exercise training on the menstrual cycle: existence and mechanisms. *Med Sci Sports Exerc.* 1990;22(3):275-280.
79. Warren MP, Perloth NE. The effects of intense exercise on the female reproductive system. *J Endocrinol.* 2001;170(1):3–11.
80. De Souza MJ, Maguire MS, Rubin KR, et al. Effects of menstrual phase and amenorrhea on exercise performance in runners. *Med Sci Sports Exerc.* 1990;22(5):575–80.
81. Beidleman BA, Rock PB, Muza SR, et al. Exercise VE and physical performance at altitude are not affected by menstrual cycle phase. *J Appl Physiol.* 1999;86(5):1519–26.
82. Stachenfeld NS, Silva C, Keefe DL. Estrogen modifies the temperature effects of progesterone. *J Appl Physiol.* 2000;88(5):1643–9.
83. Williams TJ, Krahenbuhl GS. Menstrual cycle phase and running economy. *Med Sci Sports Exerc.* 1997;29(12):1609-18.
84. Janse de Jonge XAK, Thompson MW, Chuter VH, et al. Exercise performance over the menstrual cycle in temperate and hot, humid conditions. *Med Sci Sports Exerc.* 2012;44(11):2190–8.
85. Pivarnik JM, Marichal CJ, Spillman T, et al. Menstrual cycle phase affects temperature regulation during endurance exercise. *J Appl Physiol.* 1992;72(2):543–8.
86. Hessemer V, Bruck K. Influence of menstrual cycle on thermoregulatory, metabolic, and heart rate responses to exercise at night. *J Appl Physiol.* 1985;59(6):1911-7.
87. Smekal G, Duvillard von SP, Frigo P, et al. Menstrual Cycle: no effect on exercise cardiorespiratory variables or blood lactate concentration. *Med Sci Sports Exerc.* 2007;39(7):1098–106.
88. Dean TM, Perreault L, Mazzeo RS, et al. No effect of menstrual cycle phase on lactate threshold. *J Appl Physiol.* 2003;95(6):2537–43.
89. Zderic TW, Coggan AR, Ruby BC. Glucose kinetics and substrate oxidation during exercise in the follicular and luteal phases. *J Appl Physiol.* 2001;90(2):447–53.
90. Tsampoukos A, Peckham EA, James R, et al. Effect of menstrual cycle phase on sprinting performance. *Eur J Appl Physiol.* 2010;109(4):659–67.
91. Sunderland C, Nevill M. Effect of the menstrual cycle on performance of intermittent, high-intensity shuttle running in a hot environment. *Eur J Appl Physiol.* 2003;88(4-5):345-52.
92. Maia H Jr, Casoy J. Non-contraceptive health benefits of oral contraceptives. *Eur J Contracept Reprod Health Care.* 2008;13(1):17–24.
93. Rechichi C, Dawson B, Goodman C. Athletic performance and the oral contraceptive. *Int J Sports Physiol Perform.* 2009;4(2):151–62.
94. Rechichi C, Dawson B. Effect of oral contraceptive cycle phase on performance in team sport players. *J Sci Med Sport.* 2009;12(1):190–5.
95. Casazza GA, Suh S-H, Miller BF, et al. Effects of oral contraceptives on peak exercise capacity. *J Appl Physiol.* 2002;93(5):1698–702.

96. Lebrun CM, Petit MA, McKenzie DC, et al. Decreased maximal aerobic capacity with use of a triphasic oral contraceptive in highly active women: a randomised controlled trial. *Br J Sports Med.* 2003;37(4):315–20.
97. Brooks JHM, Fuller CW. The influence of methodological issues on the results and conclusions from epidemiological studies of sports injuries: illustrative examples. *Sports Med.* 2006;36(6):459–72.
98. Ostenberg A, Roos H. Injury risk factors in female European football. A prospective study of 123 players during one season. *Scand J Med Sci Sports.* 2000;10(5):279–85.
99. Söderman K, Pietilä T, Alfredson H, et al. Anterior cruciate ligament injuries in young females playing soccer at senior levels. *Scand J Med Sci Sports.* 2002;12(2):65–8.
100. Soligard T, Grindem H, Bahr R, et al. Are skilled players at greater risk of injury in female youth football? *Br J Sports Med.* 2010;44(15):1118–23.
101. Junge A, Dvorak J. Soccer injuries. *Sports Med.* 2004;34(13):929–38.
102. Junge A, Dvorak J. Injuries in female football players in top-level international tournaments. *Br J Sports Med.* 2007;41(Supplement 1):i3–i7.
103. Jacobson I, Tegner Y. Injuries among Swedish female elite football players: a prospective population study. *Scand J Med Sci Sports.* 2007;17(1):84–91.
104. Tegnander A, Olsen OE, Moholdt TT, et al. Injuries in Norwegian female elite soccer: a prospective one-season cohort study. *Knee Surg Sports Traumatol Arthrosc.* 2008;16(2):194–8.
105. Le Gall F, Carling C, Reilly T. Injuries in young elite female soccer players: an 8-season prospective study. *Am J Sports Med.* 2008;36(2):276–84.
106. Faude O. Injuries in female soccer players: a prospective study in the German national league. *Am J Sports Med.* 2005;33(11):1694–700.
107. Giza E, Mithöfer K, Farrell L, et al. Injuries in women’s professional soccer. *Br J Sports Med.* 2005;39(4):212–6.
108. Häggglund M, Waldén M, Ekstrand J. UEFA injury study—an injury audit of European Championships 2006 to 2008. *Br J Sports Med.* 2009;43(7):483–9.
109. Engström B, Johansson C, Törnkvist H. Soccer injuries among elite female players. *Am J Sports Med.* 1991;19(4):372–5.
110. Söderman K, Adolphson J, Lorentzon R, et al. Injuries in adolescent female players in European football: a prospective study over one outdoor soccer season. *Scand J Med Sci Sports.* 2001;11(5):299–304.
111. Hawkins RD, Fuller CW. A prospective epidemiological study of injuries in four English professional football clubs. *Br J Sports Med.* 1999;33(3):196–203.
112. Häggglund M, Waldén M, Bahr R, et al. Methods for epidemiological study of injuries to professional football players: developing the UEFA model. *Br J Sports Med.* 2005;39(6):340–6.
113. Ekstrand J, Hilding J. The incidence and differential diagnosis of acute groin injuries in male soccer players. *Scand J Med Sci Sports.* 2007;9(2):98–103.
114. Häggglund M, Waldén M, Atroshi I. Preventing knee injuries in adolescent female football players – design of a cluster randomized controlled trial. *BMC Musculoskelet Disord.* 2009;10(1):75.
115. Waldén M, Häggglund M, Magnusson H, et al. Anterior cruciate ligament injury in elite football: a

- prospective three-cohort study. *Knee Surg Sports Traumatol Arthrosc.* 2011;19(1):11–9.
116. Wild CY, Steele JR, Munro BJ. Why do girls sustain more anterior cruciate ligament injuries than boys? *Sports Med.* 2012;42(9):733–49.
 117. Gottlob CA, Baker CL. Anterior cruciate ligament reconstruction: socioeconomic issues and cost effectiveness. *Am J Orthop.* 2000;29(6):472–6.
 118. Hewett TE. Anterior cruciate ligament injuries in female athletes: Part 1, Mechanisms and risk factors. *Am J Sports Med.* 2006;34(2):299–311.
 119. Roos H, Ornell M, Gärdsell P, et al. Soccer after anterior cruciate ligament injury— an incompatible combination? A national survey of incidence and risk factors and a 7-year follow-up of 310 players. *Acta Orthop Scand.* 1995;66(2):107-12.
 120. Steffen K, Andersen TE, Krosshaug T, et al. ECSS Position Statement 2009: Prevention of acute sports injuries. *Eur J Sport Sci.* 2010;10(4):223–36.
 121. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med.* 2006;35(3):359–67.
 122. Myer GD, Ford KR, Paterno MV, et al. The effects of generalized joint laxity on risk of anterior cruciate ligament injury in young female athletes. *Am J Sports Med.* 2008;36(6):1073-80.
 123. Shambaugh JP, Klein A, Herbert JH. Structural measures as predictors of injury in basketball players. *Med Sci Sports Exerc.* 1991;23(5):522-7.
 124. Silvers HJ, Mandelbaum BR. Prevention of anterior cruciate ligament injury in the female athlete. *Br J Sports Med.* 2007;41(Supplement 1):i52–9.
 125. Malinzak RA, Colby SM, Kirkendall DT, et al. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech.* 2001;16(5):438–45.
 126. Yu B, McClure SB, Onate JA, et al. Age and gender effects on lower extremity kinematics of youth soccer players in a stop-jump task. *Am J Sports Med.* 2005;33(9):1356–64.
 127. Ryder SH, Johnson RJ. Prevention of ACL Injuries. *J Sport Rehabil.* 1997;6(2):80-96.
 128. Faunø P, Wulff Jakobsen B. Mechanism of anterior cruciate ligament injuries in soccer. *Int J Sports Med.* 2006;27(1):75–9.
 129. Hewett TE, Zazulak BT, Myer GD. Effects of the menstrual cycle on anterior cruciate ligament injury risk: a systematic review. *Am J Sports Med.* 2007;35(4):659–68.
 130. Hewett TE. Neuromuscular and hormonal factors associated with knee injuries in female athletes. Strategies for intervention. *Sports Med.* 2000;29(5):313–27.
 131. Nielsen JM, Hammar M. Sports injuries and oral contraceptive use. *Sports Med.* 1991;12(3):152–60.
 132. Martineau PA, Al-Jassir F, Lenczner E, et al. Effect of the oral contraceptive pill on ligamentous laxity. *Clin J Sport Med.* 2004;14(5):281–6.
 133. Bennell K, White S, Crossley K. The oral contraceptive pill: a revolution for sportswomen? *Br J Sports Med.* 1999;33(4):231–8.
 134. Möller-Nielsen J, Hammar M. Women's soccer injuries in relation to the menstrual cycle and oral contraceptive use. *Med Sci Sports Exerc.* 1989;21(2):126-129.
 135. Andersson H, Ekblom B, Krstrup P. Elite football on artificial turf versus natural grass: Movement

- patterns, technical standards, and player impressions. *J Sports Sci.* 2008;26(2):113–22.
136. Tamer K, Gunay M, Tiryaki G, et al. Physiological characteristics of Turkish female soccer players. In: Reilly T, Bangsbo, J and Hughes, M editors. London: E and FN Spon: *Science and Football III*; 1997. pp37-39.
137. Evangelista M, Pandolfi O, Fanton F, et al. A functional model of female soccer players: analysis of functional characteristics. *J Sports Sci.* 1992;(10):165.