

Effects of low-volume high-intensity interval training (HIT) on fitness in adults: A meta-analysis of controlled and non-controlled trials

Matthew Weston¹, Kathryn L. Taylor², Alan M. Batterham² and Will G. Hopkins³

¹Department of Sport & Exercise Sciences, School of Social Sciences & Law, Teesside University, Middlesbrough, UK

²School of Health and Social Care, Teesside University, Middlesbrough, UK

³Sports Performance Research Institute New Zealand, AUT University, Auckland, New Zealand

Corresponding author:

Dr Matthew Weston, Department of Sport & Exercise Sciences, School of Social Sciences & Law, Teesside University, Middlesbrough, UK

Email: m.weston@tees.ac.uk

Telephone: ++44 (0)1642 342380

Fax: ++44 (0)1642 342317

Abstract

Background: Low-volume high-intensity interval training (HIT) appears to be an efficient and practical way to develop physical fitness.

Objective: To estimate meta-analysed mean effects of HIT on aerobic power (maximum oxygen consumption [VO_{2max}] in an incremental test) and sprint fitness (peak and mean power in a 30-s Wingate test).

Data Sources: Five databases (PubMed, MEDLINE, Scopus, BIOSIS and Web of Science) were searched for original research articles published up to January 2014. Search terms included *high intensity, HIT, sprint, fitness, and VO_{2max}* .

Study Selection: Inclusion criteria were: fitness assessed pre and post training; training period ≥ 2 wk; repetition duration 30 to 60 s; work/rest ratio < 1.0 ; exercise intensity described as maximal or near maximal; adult subjects of age > 18 y.

Data Extraction: The final data set consisted of 55 estimates from 32 trials for VO_{2max} , 23 estimates from 16 trials for peak sprint power, and 19 estimates from 12 trials for mean sprint power. Effects on fitness were analysed as percentages via log transformation. Standard errors calculated from exact p values (where reported) or imputed from errors of measurement provided appropriate weightings. Fixed effects in the meta-regression model included type of study (controlled, uncontrolled), subject characteristics (sex, training status, baseline fitness), and training parameters (number of training sessions, repetition duration, work/rest ratio). Probabilistic magnitude-based inferences for meta-analysed effects were based on standardized thresholds for small, moderate and large changes (0.2, 0.6 and 1.2, respectively) derived from between-subject standard deviations (SDs) for baseline fitness.

Results: A mean low-volume HIT protocol (13 training sessions, 0.16 work/rest ratio) in a controlled trial produced a likely moderate improvement in the VO_{2max} of active non-athletic males (6.2%; 90% confidence limits $\pm 3.1\%$), when compared with control. There were possibly moderate improvements in the VO_{2max} of sedentary males (10.0%; $\pm 5.1\%$) and active non-athletic females (3.6%; $\pm 4.3\%$), and a likely small increase for sedentary females (7.3%; $\pm 4.8\%$). The effect on the VO_{2max} of athletic males was unclear (2.7%; $\pm 4.6\%$). A possibly moderate additional increase was likely for subjects with a $10 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ lower baseline VO_{2max} (3.8%; $\pm 2.5\%$), whereas the modifying effects of sex and difference in exercise dose were unclear. The comparison of HIT with traditional endurance training was unclear (-1.6%; $\pm 4.3\%$). Unexplained variation between studies was 2.0% (SD). Meta-analysed effects of HIT on Wingate peak and mean power were unclear.

Conclusions: Low-volume high-intensity interval training produces moderate improvements in the aerobic power of active non-athletic and sedentary subjects. More studies are needed to resolve the unclear modifying effects of sex and HIT dose on aerobic power and the unclear effects on sprint fitness.

1 Introduction

High-intensity interval training (HIT), which involves alternating bouts of intensive exercise with low-intensity recovery periods,[1] is considered one of the most effective means of improving cardiorespiratory and metabolic function [2]. Athletes and coaches have historically used HIT to improve exercise performance, but the effectiveness of HIT to improve health-related outcomes has recently generated new interest [3]. In recent reviews there appears to be a consensus for the benefit of high-intensity aerobic interval training in patient populations [3-6]. Weston et al. [6] meta-analysed ten studies and reported that high-intensity aerobic interval training, typically performed at 85-95% maximal heart rate ($\%HR_{max}$), increased cardiorespiratory fitness by almost double that of moderate-intensity continuous training in patients with lifestyle-induced chronic disease. In contrast, HIT of similar intensity elicits improvements in maximal oxygen uptake (VO_{2max}) slightly greater than those typically reported with continuous training in healthy, active adults [7].

High-intensity interval training can encompass a considerable range of exercise intensities. For example, Buchheit and Laursen [8] recently defined HIT as “either repeated short (<45 s) to long (2-4 min) bouts of rather high- but not maximal-intensity exercise, or short (<10 s, repeated-sprint sequences) or long (>20-30 s, sprint interval session) all-out sprints, interspersed with recovery periods”. As such, maximal, all-out sprint training is classified as a form of high-intensity training at the highest end of the intensity spectrum [9,10]. Here, the repeated bouts of relatively brief all-out (maximal) intermittent exercise necessitate shorter interval durations and longer recovery periods than those of traditional high-intensity aerobic interval programming, and the total weekly volume (duration) of exercise is therefore lower. There is accumulating evidence supporting improved aerobic exercise performance following this form of training. Kessler et al. [3] reviewed five studies with exercise intensity described as all-out and concluded that it was an effective means of improving VO_{2max} . Sloth et al. [10] meta-analysed standardized effects of low-volume all-out interval training on VO_{2max} in 13 studies and reported an overall moderate effect (standardised change in the mean of 0.63). However, modifying effects of study and subject characteristics were not accounted for in their meta-analysis, and neither were studies with reference groups representing traditional endurance training, rather than no training. Using similar inclusion criteria (e.g., 30-s all-out sprints) Gist and colleagues [11] meta-analysed 16 randomised controlled trials and reported a moderate effect (0.69) of HIT on VO_{2max} in comparison with no-exercise control groups and a trivial effect (0.04) when compared with endurance-training controls. Effects on physical performance should, however, be meta-analysed in percent units before assessment via standardization [12]. Gist et al. [11] reported no significant effects of initial fitness, intervention length, inclusion of additional training or mode of training in response to HIT, but they did not report the effect of sex or work to rest ratio. The magnitude of the benefit of low-volume HIT on aerobic power, therefore, has still to be summarised adequately.

Low-volume HIT may also have the potential to improve sprint power, as it increases enzymatic activities of anaerobic metabolism [13]. Most sporting activities depend upon the expression of power for short or sustained periods of time [14]. Furthermore, many basic daily activities are dependent on the ability to generate force at high velocity, and power training can improve mobility-related outcomes in the elderly [15] as well as increasing self-efficacy, satisfaction with physical function and overall life satisfaction [16]. A meta-analysis of the effect of HIT on sprint power is therefore timely. Our aim for this review was to use a mixed-model meta-analysis to provide estimates of the effect of low-volume HIT on fitness (VO_{2max} , 30-s Wingate power) along with the modifying effects of study and subject characteristics.

2 Methods

2.1 Literature search

A search of five databases (PubMed, MEDLINE, Scopus, BIOSIS and Web of Science), along with the reference lists of original research and review articles published in English up to January 2014 was conducted by two of the authors (KT, MW). Our independent variable search terms were ‘aerobic high intensity’, ‘high intensity’, ‘HIT’, ‘intervals’, ‘intensive’, ‘sprint’, ‘repeated sprint’, and the dependent variable search terms were ‘fitness’, ‘aerobic fitness’, ‘anaerobic fitness’, ‘ VO_{2max} ’, ‘performance’, ‘endurance’ and ‘adaptations’. Independent variable search terms were combined with dependent variable search terms, giving a total of 49 combinations.

2.2 Study selection

The most common model employed in low-volume HIT studies consists of four to six 30 s ‘all-out’ efforts separated by ~4 min of recovery, for a total of 2-3 intense exercise during a single training session [9]. As such, our study selection criteria were: VO_{2max} or 30-s Wingate power assessed pre and post training, training period ≥ 2 weeks, repetition duration 30 to 60 s, work/rest ratio < 1.0 , exercise intensity described as maximal or near maximal, and adult subjects of age > 18 y. No inclusion criteria were used for participant fitness. Using the subject characteristic information provided by each study, participants were assigned to one of three groups; sedentary, active non-athletic, or athletic.

The selection of studies for our meta-analysis was confined to studies predominantly utilising the classic Wingate protocol. In doing so, we acknowledge the exclusion of a large body of laboratory- and field-based HIT research utilising longer interval durations (1-4 min) performed at high, but not maximal intensity, and with a work:rest ratio > 1.0 [8]. Furthermore, by selecting VO_{2max} as our measure of aerobic fitness, we excluded field-relevant performance measures, which may limit the application of our findings to athletic populations and sports performance. While the recent meta-analysis of Bacon et al. [7] has, to an extent, addressed this gap in the literature, the number of studies excluded from the present study on interval duration, intensity and other measures of aerobic fitness (e.g., velocity at VO_{2max} , speed at lactate threshold, running

economy and sports-specific tests) underscores the need for a dedicated review of studies using longer intervals at lower intensities.

To select relevant papers all titles were initially screened by two authors (KT, MW) during the electronic searches to exclude studies that were beyond the scope of this meta-analysis. Following this initial selection process there were 550 potentially eligible studies (Figure 1). All study titles and abstracts were then screened independently by the same authors. Full text versions of the remaining papers that met each of the eligibility criteria were then reviewed by these authors to determine final inclusion in the meta-analysis. Any disputed studies were taken to a third reviewer (AMB) for resolution. The final data set for VO_{2max} consisted of 55 estimates from 32 trials, of which 11 were controlled trials. For peak sprint power, the final data set consisted of 23 estimates from 16 trials, of which 3 were controlled trials, and for mean sprint power the data set consisted of 19 estimates from 12 trials, of which 3 were controlled trials.

2.3 Data extraction

Graph digitizer software (DigitizeIt, Germany) was used to obtain data values in studies where only plots were published. Accuracy was confirmed via intra- and inter-individual reassessments of data extraction. Mean effects on VO_{2max} , peak and mean sprint power in training and control groups were converted to a percentage change. For each converted effect, standard errors were calculated to indicate the level of imprecision. In studies where exact p-values were given (VO_{2max} $n = 7$; peak power $n = 4$; mean power $n = 5$) standard errors were calculated directly via the corresponding t-statistic and its degrees of freedom. Under the assumption that studies with similar test protocols and subject characteristics would have similar typical errors of measurement, the typical errors from these studies were then averaged (via the weighted mean variance) and assigned to the studies that did not report an exact p-value. The standard error was then calculated via the relationship between typical error and standard error [17,18]. Descriptive statistics for studies included in the meta-analysis for VO_{2max} , peak and mean sprint power are shown in Tables 1, 2 and 3, respectively.

2.4 Publication bias and outliers

To investigate the extent of publication bias, we examined the standard error against the t-value for each predicted effect for each outcome, and inspected the plot for signs of asymmetrical scatter [12]. Such a plot is an improved version of the funnel plot, as the scatter of the effects is adjusted for any uncertainty in the estimates and also for the contribution of study covariates. Examination of these plots revealed no evidence of the asymmetrical scatter associated with publication bias.

2.5 Meta-analytic model

The general linear mixed-model procedure (Proc Mixed) in the Statistical Analysis System (Version 9.2, SAS Institute, Cary, NC) was used to perform the meta-analysis. Fixed effects in the model included type of study (controlled, uncontrolled), study-level subject characteristics (sex, training status, baseline VO_{2max} , peak and mean sprint

power), and training parameters (number of training sessions, repetition duration, work/rest ratio). We determined the predicted effect of reference training conditions on VO_{2max} , peak and mean sprint power using mean values of baseline fitness, number of sessions, and work/rest ratio from all eligible studies. Performance effects were then calculated as the predicted effect under these reference training conditions. The modifying effects of predictors were also calculated, either as differences between levels of a nominal covariate (i.e., male/female, non-athletic/sedentary), or as the effect of approximately two standard deviations of a numeric covariate (i.e., a typically high value minus a typically low value) [12]. Random effects in the model were the usual between-subject random effect and a novel within-study random effect to account for within-study repeated measurements (a control treatment and/or more than one training treatment). The residual was set to unity to properly weight the estimates by the inverse of the square or their standard errors. Unexplained true variation within and between studies was estimated by combining the variances for the random effects and was expressed as a standard deviation. The standard deviation was doubled before interpreting its magnitude with the scale used to interpret fixed effects [19], for the same reason that the magnitude of the effect of a linear covariate is evaluated with two standard deviations of the covariate [12].

2.5.1 Outcome statistics

We expressed the uncertainty in the estimates of effects on fitness as 90% confidence limits (CL) and as probabilities that the true value of the meta-analysed effect was trivial, beneficial or harmful in relation to threshold values for benefit and harm. Probabilities were then used to make a qualitative probabilistic inference about the effect [12]. Given that improved aerobic functioning and power output have clinical application [3-7,15,16] main treatment effects were considered unclear if the chance of benefit (improved fitness) was high enough to warrant use of the intervention but with an unacceptable risk of harm (reduced fitness). An odds ratio of benefit to harm of <66 was used to identify such unclear effects. This ratio corresponds to a borderline possibly beneficial effect (25% chance of benefit) and a borderline most unlikely harmful effect (0.5% risk of harm). All other effects were deemed clinically clear and inference made via estimation of the probability that the true magnitude of the effect was at least as large as our pre-specified thresholds. In the absence of robust anchors for the smallest worthwhile clinical and practical effect on VO_{2max} and sprint power, our inferences were based on standardised thresholds for small, moderate and large changes of 0.2, 0.6 and 1.2 standard deviations, respectively [12] and derived by averaging appropriate between-subject variances for baseline VO_{2max} , peak and mean sprint power. For VO_{2max} , magnitude thresholds were 3.2, 9.6 and 19.2% for sedentary subjects, 1.4, 4.1 and 8.1% for active non-athletic subjects, and 1.4, 4.2 and 8.4% for athletic subjects. For peak and mean sprint power, thresholds were 1.7, 5.1 and 10.3% and 1.7, 5.2, and 10.5%, respectively for male subjects. The chance of the true effect being trivial, beneficial or harmful was interpreted using the following scale: $<0.5\%$, most unlikely; 0.5–5%, very unlikely; 5–25%, unlikely; 25–75%, possibly; 75–95%, likely; 95–99.5%, very likely; $>99.5\%$, most likely [12]. Modifying effects were evaluated non-clinically

and deemed unclear if the 90% confidence limit overlapped the thresholds for the smallest worthwhile positive and negative effects [12].

3 Results

3.1 Maximum oxygen consumption

The meta-analysed effects on VO_{2max} of an average low-volume HIT protocol in a controlled trial are shown in Table 4. When compared with control, moderate improvements in VO_{2max} were likely for active non-athletic males and possible for sedentary males and active non-athletic females. A small improvement in VO_{2max} was likely for sedentary females. The effect on athletic males was unclear. With the exception of a possible moderate additional increase in VO_{2max} for subjects with a lower baseline value, the effects of all modifiers were unclear. The comparison of HIT with endurance training was unclear (-1.6%; 90% confidence limits $\pm 4.3\%$). Unexplained variation expressed as a between-study SD was 2.0% ($\pm 2.7\%$).

3.2 Sprint power

The meta-analysed effects of low-volume HIT on 30-s Wingate peak and mean sprint power in a controlled trial are shown in Table 5 and 6, respectively. With the exception of a possibly moderate improvement in the peak sprint power of controls, all mean effects on sprint power were unclear. There were possibly moderate and likely small improvements in mean and peak sprint power, respectively following a 3-fold increase in the number of training sessions. A moderately beneficial improvement in peak sprint power with a greater work/rest ratio was likely and a small additional increase in mean sprint power was possible for subjects with a lower baseline value. All other modifying effects were unclear. Unexplained variation between studies was 2.4% ($\pm 2.5\%$) and 1.0% ($\pm 2.9\%$) for peak and mean sprint power, respectively.

4 Discussion

In the previous meta-analyses of Sloth et al. [10] and Gist et al. [11], low-volume HIT improved aerobic fitness and Wingate sprint power, but the effects on different subject groups and other modifying effects were either not analysed or presented. Our meta-analysis broadens the scope of these previous reviews as it is the first to include study and subject characteristics in the analysis. Our data revealed HIT to have an apparent adaptive effect on VO_{2max} that favours the less fit. Despite HIT effectively representing repeated Wingate tests, there was no clear effect on measures of performance in the test.

We found that a mean protocol of 13 HIT sessions with a work/rest ratio of 0.16 led to moderate improvements in the VO_{2max} of sedentary and non-athletic males and females. This main finding is consistent with the recent work of Sloth et al. [10], Gist et al. [11] and Bacon et al. [7], who reported standardised moderate effects on VO_{2max} for HIT and high-intensity aerobic interval training, respectively. A combination of central and peripheral adaptations promoting an enhanced availability, extraction and utilisation of oxygen may explain such improvements following intensive interval-training protocols.

Mechanisms responsible for increased $\text{VO}_{2\text{max}}$ following HIT were not the focus of our review, however. Comprehensive reviews of the possible underlying mechanisms are available elsewhere [9,10].

Gibala et al. [58] reported low-volume HIT to be a time-efficient strategy for rapid physiological and performance improvements that are comparable to improvements following traditional endurance training. The random effects component of our mixed-model enabled us to include studies where the reference group was traditional endurance training rather than no training. Here, the comparison between the two types of training was unclear. This finding is consistent with that of Gist et al. [11] who reported a trivial effect of HIT on $\text{VO}_{2\text{max}}$ when compared with endurance training controls. More studies are therefore required to examine the effectiveness of HIT against traditional endurance training for training-induced endurance gains. The effect of HIT on $\text{VO}_{2\text{max}}$ was greater for the less fit, which is consistent with training in general having greater effects on the less fit [59]. For already highly trained athletes who replaced their usual training with HIT, as opposed to the HIT being added, the effect on $\text{VO}_{2\text{max}}$ was unclear. This finding also indicates the need for more research, providing elite athletes can be convinced to experiment with their normal training programmes [9]. Despite reporting no analytical data for the potentially modifying effect of training duration, Sloth et al. [10] and Gist et al. [11] reported no clear effects of the length of HIT intervention on the magnitude of $\text{VO}_{2\text{max}}$ improvement. The data presented in our more extensive meta-analysis has still not resolved this issue.

On the basis of the confidence limits low-volume HIT had an unclear effect on peak and mean sprint power that could at most be a moderate beneficial or a small harmful effect. These results are consistent with those of Sloth et al. [10], who reported enhanced peak and mean power following HIT. Their assertion was based on nine studies [24,28-30,33,36,41,42,52], without a meta-analysis of the mean effect and its uncertainty. Three of these studies [30,36,42] were excluded from our analysis, owing to difficulties in obtaining precise baseline and post-intervention data during the data extraction process. An enhanced sprint power following HIT was expected, given that all-out training increases enzymatic activity related to anaerobic metabolism [13]. Furthermore, studies showing strong similarities between testing and training routines are more likely to show training improvements [60]. However, when measured relative to controls, the meta-analysed effect of HIT on sprint power was unclear. Improvements of 4.5% in peak sprint power and 2.8% in mean sprint power of control subjects may have represented a learning effect on the Wingate test or provide some evidence of compensatory rivalry (e.g., greater effort by controls). There was some evidence of a dose-reponse relationship and a greater effect for the less fit. The finding of a possibly moderate enhancement in peak sprint power with a 5-fold increase in repetition work/rest ratio could be explained by greater phosphocreatine resynthesis in the recovery phase [61,62].

There was considerable uncertainty in the standard deviations representing the residual between-study variation in the mean effect of the treatment on the three measures of fitness, but in this sample of studies the observed magnitudes (after doubling the standard deviations) were small to moderate, depending on the measure of performance and the subject group. This standard deviation needs to be added to and subtracted from the main effect to evaluate the magnitude of the HIT treatment in a specific setting. For example, the mean effect of HIT on $\text{VO}_{2\text{max}}$ for active non-athletic males (6.2%; moderate) in any given setting could be anywhere from 4.2% (very likely small) to 8.2% (possibly large). Such differences between the effects of training in the different studies presumably reflect differences in subject characteristics and training protocols that are not properly accounted for by the published data. Some data may also have been analysed or reported erroneously.

We propose several areas for future research along with suggestions for those publishing research in this area. Given the age of participants included within our meta-analysis was mainly young adults, it is evident that research is required to clarify the effects of low-volume HIT in older populations. Moderating effects of changing the exercise dose on $\text{VO}_{2\text{max}}$ were unclear, as was the replacement of athletes' usual aerobic training with HIT, indicating that more research is necessary to investigate these predictors. We do, however, recommend that modifying effects are interpreted with slight caution as when a covariate is a subject characteristic averaged over study subjects, the observed meta-regression relationships might not hold at the individual study level [63]. The practicality of low-volume HIT warrants further investigation, given that repeated bouts of maximal exercise require high levels of motivation [9]. Adherence to unsupervised training also needs investigation [29]. We concur with the need to test the effectiveness of low-volume HIT via large-scale, multi-centre, randomised clinical trials in various clinical populations and on long-term clinical outcome measures [64]. Of further benefit would be the reporting of full inferential statistics, such as SD of change scores or exact p values in training and control groups, to enable meta-analysis of the magnitude of individual responses. Finally, the findings of a training study are of very little or no value without precise information of the training itself [65]. We therefore encourage authors to report physiological responses during HIT sessions, as this practice will help to demonstrate that the fidelity of an intervention has been upheld for all subjects.

5 Conclusions

Low-volume HIT is increasingly being used for aerobic adaptations previously achieved with traditional endurance training. Our meta-analysis provides evidence of substantial improvements in the endurance fitness of sedentary and non-athletic subjects following repeated bouts of brief maximal intermittent exercise. The effect of HIT on sprint power should be determined with more studies.

Acknowledgements

No sources of funding were used to assist in the preparation of this review. The authors have no conflicts of interest that are directly relevant to the content of this review.

References

1. Helgerud J, Høydal K, Wang E, et al. Aerobic high-intensity intervals improve VO₂max more than moderate training. *Med Sci Sports Exerc.* 2007;39(4):665-71.
2. Buchheit M, Laursen P. High-intensity interval training, solutions to the programming puzzle. *Sports Med.* 2013;43:313-38
3. Kessler HS, Sisson SB, Short KR. The potential for high-intensity interval training to reduce cardiometabolic disease risk. *Sports Med.* 2012;42(6):489-509.
4. Guiraud T, Nigam A, Gremeaux V, et al. High-intensity interval training in cardiac rehabilitation. *Sports Med.* 2012;42(7):587-605.
5. Hwang CL, Wu YT, Chou CH. Effect of aerobic interval training on exercise capacity and metabolic risk factors in people with cardiometabolic disorders: a meta-analysis. *J Cardiopulm Rehabil Prev.* 2011;31(6):378-85.
6. Weston KS, Wisløff U, Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med.* 2013 Oct 21. doi:10.1136/bjsports-2013-092576. [Epub ahead of print].
7. Bacon AP, Carter RE, Ogle EA, et al. VO₂max trainability and high intensity interval training in humans: a meta-analysis. *PLoS One.* 2013 Sep 16;8(9):e73182. doi: 10.1371/journal.pone.0073182.
8. Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle. *Sports Med.* 2013;43:927-954.
9. Gibala MJ, Little JP, Macdonald MJ, et al. Physiological adaptations to low-volume, high-intensity interval training in health and disease. *J Physiol.* 2012;590(Pt 5):1077-84.
10. Sloth M, Sloth D, Overgaard K, et al. Effects of sprint interval training on VO₂max and aerobic exercise performance: A systematic review and meta-analysis. *Scand J Med Sci Sports.* 2013 Jul 25. doi: 10.1111/sms.12092. [Epub ahead of print]
11. Gist NH, Fedewa MV, Dishman RK, et al. Sprint interval training effects on aerobic capacity: A systematic review and meta-analysis. *Sports Med.* 2014;44(2):269-79.
12. Hopkins WG, Marshall SW, Batterham AM, et al. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3-12.
13. Rodas G, Ventura JL, Cadefau JA, et al. A short training programme for the rapid improvement of both aerobic and anaerobic metabolism. *Eur J Appl Physiol.* 2000;82(5-6):480-6.

14. Davison RR, Van Someren KA, Jones AM. Physiological monitoring of the Olympic athlete. *J Sports Sci.* 2009;27(13):1433-42.
15. Katula JA, Rejeski WJ, Marsh AP. Enhancing quality of life in older adults: a comparison of muscular strength and power training. *Health Qual Life Outcomes.* 2008;6:45.
16. Hruda KV, Hicks AL, McCartney N. Training for muscle power in older adults: effects on functional abilities. *Can J Appl Physiol.* 2003;28(2):178-89.
17. Vandenberghe TJ, Hopkins WG. Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis. *Sports Med.* 2011;41(9): 773-92.
18. Carr AJ, Hopkins WG, Gore CJ. Effects of acute alkalosis and acidosis on performance: a meta-analysis. *Sports Med.* 2011;41(10):801-14.
19. Smith TB, Hopkins WG. Variability and predictability of finals times of elite rowers. *Med Sci Sports Exerc.* 2011;43:2155-60.
20. Rakobowchuk M, Harris E, Taylor A, et al. Heavy and moderate interval exercise training alters low-flow-mediated constriction but does not increase circulating progenitor cells in healthy humans. *Exp Physiol.* 2012;97(3):375-85.
21. Siahkouhian M, Khodadadi D, Shahmoradi K. Effects of high-intensity interval training on aerobic and anaerobic indices: Comparison of physical active and inactive men. *Sci Sports.* 2013;28:e119-e125.
22. Trilk JL, Singhal A, Bigelman KA, et al. Effect of sprint interval training on circulatory function during exercise in sedentary, overweight/obese women. *Eur J Appl Physiol.* 2011;111:1591-7.
23. Allemeier CA, Fry AC, Johnson P, et al. Effects of sprint cycle training on human skeletal muscle. *J Appl Physiol.* 1994;77(5):2385-90.
24. McKenna MJ, Heigenhauser GJ, McKelvie RS, et al. Enhanced pulmonary and active skeletal muscle gas exchange during intense exercise after sprint training in men. *J Physiol.* 1997;501(Pt 3):703-16.
25. Macpherson RE, Hazell TJ, Olver TD, et al. Run sprint interval training improves aerobic performance but not maximal cardiac output. *Med Sci Sports Exerc.* 2011;43: 115-22.
26. Esfandiari S, Sasson Z, Goodman JM. Short-term high-intensity interval and continuous moderate-intensity training improve maximal aerobic power and diastolic filling during exercise. *Eur J Appl Physiol.* 2014; 114 (2):331-43.
27. Dunham C, Harms CA. Effects of high-intensity interval training on pulmonary function. *Eur J Appl Physiol.* 2012;112:3061-8.
28. Bayati M, Farzad B, Gharakhanlou R, et al. A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble “all-out” sprint interval training. *J Sports Sci Med.* 2011;10:571-6.
29. Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism.* 2010;59:1421-8.
30. Hazell TJ, Macpherson RE, Gravelle BM, et al. 10 Or 30-S sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol.* 2010;110:153-60.

31. Jacobs RA, Flück D, Bonne TC, et al. Improvements in exercise performance with high-intensity interval training coincide with an increase in skeletal muscle mitochondrial content and function. *J Appl Physiol.* 2013;115:785-93.
32. Sharp RL, Costill DL, Fink WJ, et al. Effects of eight weeks of bicycle ergometer sprint training on human muscle buffer capacity. *Int J Sports Med.* 1986;7(1):13-7.
33. Barnett C, Carey M, Proietto J, et al. Muscle metabolism during sprint exercise in man: influence of sprint training. *J Sci Med Sport.* 2004;7:314-22.
34. Harmer AR, Ruell PA, McKenna MJ, et al. Effects of sprint training on extrarenal potassium regulation with intense exercise in Type 1 diabetes. *J Appl Physiol.* 2006;100(1):26-34.
35. Shepherd SO, Cocks M, Tipton KD, et al. Sprint interval and traditional endurance training increase net intramuscular triglyceride breakdown and expression of perilipin 2 and 5. *J Physiol.* 2013;591(3):657-75.
36. Burgomaster KA, Howarth KR, Phillips SM, et al. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *J Physiol.* 2008;586:151-60
37. Bailey SJ, Wilkerson DP, Dimenna FJ, et al. Influence of repeated sprint training on pulmonary O₂ uptake and muscle deoxygenation kinetics in humans. *J Appl Physiol.* 2009;106:1875-87.
38. MacDougall JD, Hicks AL, MacDonald JR, et al. Muscle performance and enzymatic adaptations to sprint interval training. *J Appl Physiol.* 1998;84:2138-42.
39. Tong TK, Chung PK, Leung RW, et al. Effects of non-wingate-based high-intensity interval training on cardiorespiratory fitness and aerobic-based exercise capacity in sedentary subjects: a preliminary study. *Journal of Exercise Science & Fitness.* 2011;9(2):75-81.
40. Esfarjani F, Laursen PB. Manipulating high-intensity interval training: effects on VO₂max, the lactate threshold and 3000 m running performance in moderately trained males. *J Sci Med Sport.* 2007;10(1):27-35.
41. Burgomaster KA, Heigenhauser GJ, Gibala MJ. Effect of short-term sprint interval training on human skeletal muscle carbohydrate metabolism during exercise and time-trial performance. *J Appl Physiol.* 2006;100:2041-7.
42. Astorino TA, Allen RP, Roberson DW, et al. Effect of high-intensity interval training on cardiovascular function, VO₂max, and muscular force. *J Strength Cond Res.* 2012;26:138-45.
43. Sandvei M, Jeppesen PB, Støen L, et al. Sprint interval running increases insulin sensitivity in young healthy subjects. *Arch Physiol Biochem.* 2012;118(3):139-47.
44. Rowan AE, Kueffner TE, Stavrianeas S. Short duration high-intensity interval training improves aerobic conditioning of female college soccer players. *Int J Exerc Sci.* 2012;5(3): 232-8.

45. Stathis CG, Febbraio MA, Carey MF, et al. Influence of sprint training on human skeletal muscle purine nucleotide metabolism. *J Appl Physiol.* 1994;76(4):1802-9.
46. Barnes KR, Hopkins WG, McGuigan MR, et al. Effects of different uphill interval-training programs on running economy and performance. *Int J Sports Physiol Perform.* 2013;8:639-47.
47. Laursen PB, Shing CM, Peake JM, et al. Interval training program optimization in highly trained endurance cyclists. *Med Sci Sports Exerc.* 2002;34(11):1801-7.
48. Harmer AR, McKenna MJ, Sutton JR, et al. Skeletal muscle metabolic and ionic adaptations during intense exercise following sprint training in humans. *J Appl Physiol.* 2000;89(5):1793-803.
49. Laursen PB, Blanchard MA, Jenkins DG. Acute high-intensity interval training improves \dot{V}_{O2} and peak power output in highly trained males. *Can J Appl Physiol.* 2002;27(4):336-48.
50. Dalleck L, Bushman TT, Crain RD, et al. Dose-response relationship between interval training frequency and magnitude of improvement in lactate threshold. *Int J Sports Med.* 2010;31(8):567-71.
51. Iaia FM, Hellsten Y, Nielsen JJ, et al. Four weeks of speed endurance training reduces energy expenditure during exercise and maintains muscle oxidative capacity despite a reduction in training volume. *J Appl Physiol.* 2009;106(1):73-80.
52. Burgomaster KA, Hughes SC, Heigenhauser GJ, et al. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *J Appl Physiol.* 2005;98:1985-90.
53. Forbes SC, Slade JM, Meyer RA. Short-term high-intensity interval training improves phosphocreatine recovery kinetics following moderate-intensity exercise in humans. *Appl Physiol Nutr Metab.* 2008;33(6):1124-31.
54. Richards JC, Johnson TK, Kuzma JN, et al. Short-term sprint interval training increases insulin sensitivity in healthy adults but does not affect the thermogenic response to beta-adrenergic stimulation. *J Physiol.* 2010;588:2961-72.
55. Esbjörnsson Liljedahl M, Holm I, Sylvén C, et al. Different responses of skeletal muscle following sprint training in men and women. *Eur J Appl Physiol Occup Physiol.* 1996;74(4):375-83.
56. Jansson E, Esbjörnsson M, Holm I, et al. Increase in the proportion of fast-twitch muscle fibres by sprint training. *Acta Physiol Scand.* 1990;140:359-63
57. McKenna MJ, Schmidt TA, Hargreaves M, et al. Sprint training increases human skeletal muscle $\text{Na}^+\text{-K}^+\text{-ATPase}$ concentration and improves K^+ regulation. *J Appl Physiol.* 1993;75(1):173-80.
58. Gibala MJ, Little JP, van Essen M, et al. Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *J Physiol.* 2006;575:901-11.
59. Wilmore JH, Costill DL. *Physiology of sport and exercise.* 2nd ed. USA: Human Kinetics, 1999.

60. Buchheit M. Should we be recommending repeated sprints to improve repeated-sprint performance? *Sports Med.* 2012;42(2):169-72.
61. Bogdanis GC, Nevill ME, Boobis LH, et al. Contribution of phosphocreatine and aerobic metabolism to energy supply during repeated sprint exercise. *J Appl Physiol.* 1996; 80: 876-84.
62. Bishop D, Girard O, Mendez-Villanueva A. Repeated-sprint ability - part II: recommendations for training. *Sports Med.* 2011;41(9):741-56.
63. Petkova E, Tarpey T, Huang L, et al. Interpreting meta-regression: application to recent controversies in antidepressants' efficacy. *Stat Med.* 2013;32(17):2875-92.
64. Hawley JA, Gibala MJ. What's new since Hippocrates? Preventing type 2 diabetes by physical exercise and diet. *Diabetologia.* 2012;55(3):535-9.
65. Mujika I. The alphabet of sport science research starts with q. *Int J Sports Physiol Perform.* 2013;8:465-6.

Table 1. Study and subject characteristics for VO_{2max} estimates included in the meta-analysis. Studies are sorted from the largest to the smallest effects on VO_{2max} in the intervention group.

Study (year)	Study design	Subjects	Age (y [mean])	Sample size	Proportion of males	Duration (weeks)	Total sessions	Group	Exercise intensity ^a	No. of reps			Rep duration (s)	Work / rest ratio	Effect on VO_{2max} (%)	
										Start	End	Total reps			Mean	SE
Rakobowchuk et al. [20]	NC	Non-Ath	23.1	11	0.36	6	18	HIT	100 % P_{max}	20	27	207	30	0.50	15.4	2.8
Siahkouhian et al. [21]	NC	Non-Ath	19.1	12	1.00	8	24	HIT	All-out	6	7	191	30	0.125	13.9	3.0
Siahkouhian et al. [21]	NC	Ath	19.4	12	1.00	8	24	HIT	All-out	6	7	191	30	0.125	7.3	3.0
Trilk et al. [22]	C	Sed	30.1	14	0.00	4	12	HIT	All-out	4	7	66	30	0.13	13.4	2.5
Trilk et al. [22]	C	Sed	31.4	14	0.00	4	-	CON	-	-	-	-	-	-	-0.5	2.5
Allemeier et al. [23]	C	Non-Ath	22.7	11	1.00	6	15	HIT	All-out	3	3	45	30	0.03	12.5	2.8
Allemeier et al. [23]	C	Non-Ath	24.0	6	1.00	6	-	CON	-	-	-	-	-	-	-0.7	3.8
McKenna et al. [24]	NC	Non-Ath	20.9	8	1.00	7	21	HIT	All-out	4	10	174	30	0.14	12.5	3.3
MacPherson et al. [25]	NC	Non-Ath	24.3	10	0.60	6	18	HIT	All-out	4	6	90	30	0.13	11.5	2.9
MacPherson et al. [25]	NC	Non-Ath	22.8	10	0.60	6	18	END	65 % VO_{2max}	30	60	-	2700	-	12.5	2.9
Esfandiari et al. [26]	NC	Non-Ath	24.5	8	1.00	2	6	HIT	95-100 % VO_{2max}	8	12	60	60	0.8	11.1	3.7
Esfandiari et al. [26]	NC	Non-Ath	25.6	8	1.00	2	6	END	65 % VO_{2max}	-	-	-	6300	-	4.5	3.7
Dunham et al. [27]	NC	Non-Ath	20.2	8	?	4	12	HIT	90 % P_{max}	5	5	60	60	0.333	9.6	3.7
Dunham et al. [27]	NC	Non-Ath	21.3	7	?	4	12	END	60-70 % P_{max}	-	-	-	2700	-	5.5	4.0

The effect of HIT on fitness

Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	12	HIT	All-out	3	4	48	30	0.13	9.6	4.0
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	12	HIT	125 %P _{max}	6	8	96	30	0.25	9.7	4.1
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	-	CON	-	-	-	-	-	-	-0.9	5.5
Whyte et al. [29]	NC	Sed	32.1	10	1.00	2	6	HIT	All-out	4	6	30	30	0.11	9.5	3.2
Hazell et al. [30]	C	Non-Ath	24.0	12	0.73	2	6	HIT	All-out	4	6	30	30	0.13	9.3	2.7
Hazell et al. [30]	C	Non-Ath	24.0	12	?	2	-	CON	-	-	-	-	-	-	0.2	1.7
Jacobs et al. [31]	NC	Non-athletes	27.0	16	1.00	2	6	HIT	100 %P _{max}	8	12	60	60	0.8	8.9	3.6
Sharp et al. [32]	NC	Non-Ath	25.5	8	1.00	8	32	HIT	All-out	8	8	256	30	0.13	8.3	3.3
Barnett et al. [33]	C	Non-Ath	21.2	8	1.00	8	24	HIT	All-out	3	6	108	30	0.17	8.2	1.5
Barnett et al. [33]	C	Non-Ath	21.2	8	1.00	8	-	CON	-	-	-	-	-	-	4.1	1.4
Harmer et al. [34]	C	Non-Ath	24.0	7	0.57	7	21	HIT	All-out	-	-	174	30	0.14	8.2	3.5
Harmer et al. [34]	C	Non-Ath	25.0	8	0.63	7	21	HIT	All-out	4	10	174	30	0.14	2.4	3.3
Sheperd et al. [35]	NC	Non-Ath	22.0	8	1.00	6	18	HIT	All-out	4	6	90	30	0.11	7.6	3.7
Sheperd et al. [36]	NC	Non-Ath	21.0	8	1.00	6	30	END	65 %VO _{2max}	-	-	-	3000	-	15.6	3.7
Burgomaster et al. [36]	NC	Non-Ath	24.0	10	0.50	6	18	HIT	All-out	4	6	90	30	0.11	7.3	2.9

Burgomaster et al. [36]	NC	Non-Ath	23.0	10	0.50	6	30	END	65 %VO _{2max}	-	-	-	3000	-	9.8	2.9
Bailey et al. [37]	C	Non-Ath	21.0	8	0.63	2	6	HIT	All-out	4	7	35	30	0.13	7.1	3.3
Bailey et al. [37]	C	Non-Ath	21.0	8	0.63	2	6	END	90 %GET	-	-	-	1200	-	0.0	3.3
Bailey et al. [37]	C	Non-Ath	21.0	8	0.63	2	-	CON	-	-	-	-	-	-	-2.1	3.3
MacDougall et al. [38]	NC	Non-Ath	22.7	12	1.00	7	21	HIT	All-out	4	10	174	30	0.15	6.9	2.7
Tong et al. [39]	NC	Sed	22.9	8	0.25	6	18	HIT	120 %P _{max}	20	20	360	30	0.50	6.6	3.3
Esfarnjani et al. [40]	NC	Ath	19.0	6	1.00	10	20	HIT	130 %vVO _{2max}	7	12	190	30	0.11	6.2	3.8
Esfarnjani et al. [40]	NC	Ath	19.0	5	1.00	10	40	CON	75 %vVO _{2max}	-	-	-	3600	-	2.1	4.1
Burgomaster et al. [41]	NC	Non-Ath	22.0	8	1.00	2	6	HIT	All-out	4	7	30	30	0.13	5.5	3.2
Astorino et al. [42]	C	Non-Ath	25.3	20	0.55	2	6	HIT	All-out	4	6	30	30	0.10	5.5	2.1
Astorino et al. [42]	C	Non-Ath	22.8	9	0.55	2	-	CON	-	-	-	-	-	-	1.6	3.1
Sandvei et al. [43]	NC	Non-Ath	25.2	11	0.36	8	24	HIT	All-out	5	10	189	30	0.17	5.1	2.8
Sandvei et al. [43]	NC	Non-Ath	25.2	12	0.33	8	24	END	70-80 %HR _{max}	-	-	-	2700	-	3.8	2.7
Rowan et al. [44]	NC	Ath	19.5	7	0.00	5	10	HIT	All-out	5	5	25	30	0.125	4.7	4.0
Rowan et al. [44]	NC	Ath	19.5	6	0.00	5	10	END	80 %VO _{2max}	-	-	-	2400	-	3.3	4.3
Stathis et al. [45]	NC	Non-Ath	22.1	8	0.75	7	21	HIT	All-out	3	10	153	30	0.13	4.2	3.3
Barnes et al. [46]	NC	Ath	24.9	5	1.00	6	12	HIT	110 %vVO _{2max}	8	16	74	40.5	0.333	3.6	4.7

The effect of HIT on fitness

Laursen et al. [47]	C	Ath	25.0	10	1.00	4	8	HIT	175 %P _{max}	12	12	96	30	0.11	3.1	2.9
Laursen et al. [47]	C	Ath	25.0	10	1.00	4	-	CON	-	-	-	-	-	-	0.8	2.9
Harmer et al. [48]	NC	Non-Ath	22.0	7	1.00	7	21	HIT	All-out	4	10	174	30	0.13	2.9	1.3
Laursen et al. [49]	C	Ath	23.5	7	1.00	2	4	HIT	100 %P _{max}	20	20	80	60	0.25	2.3	3.5
Laursen et al. [49]	C	Ath	23.5	7	1.00	2	-	CON	-	-	-	-	-	-	-0.8	3.5
Dalleck et al. [50]	NC	Non-Ath	21.1	10	0.45	6	6	HIT	110-120 %P _{max}	6	8	42	30	0.14	-0.7	2.9
Dalleck et al. [50]	NC	Non-Ath	21.1	10	0.45	6	12	HIT	110-120 %P _{max}	6	8	84	30	0.14	-0.6	2.9
Iaia et al. [51]	C	Ath	33.9	9	1.00	4	13.6	HIT	93 %V _{max}	8	12	124	30	0.17	-2.4	3.1
Iaia et al. [51]	C	Ath	33.9	8	1.00	4	16	CON	-	-	-	-	3138	-	0.5	3.3

^aAll-out: Encompasses intensities described by the authors as “maximal” [24]; “near maximal” [43]; “sprints” [33]; “maximum efforts” [38,44]; “supramaximal” [23]; “sprint training at the highest resistance maintained for 90 rpm” [32]; or “all-out” [21,22,25,28-30,34-37,41,42,45,48].

%P_{max}: Encompasses intensities described as either a “percentage of peak watt workload” [50], a “percentage of the highest 30 s power output completed” [47], a “percentage of peak work rate” [20,39], a “percentage of final completed work rate maintained for 10 s” [28,49]; a “percentage of peak power output” [31], a “percentage of their final workload” [27]; all determined via a pre-training incremental test.

%HR_{max}: Intensity corresponding to a percentage of maximal heart rate, determined via a pre-training incremental test.

%VO_{2max}: Intensity predicted to elicit a percentage of VO_{2max} on a treadmill [25], and of VO_{2max} on a cycle ergometer [26,36], determined via a pre-training incremental test.

%V_{max}: Intensity corresponding to a percentage of maximum running velocity, determined via a pre-training 30-s all-out sprint run.

%vVO_{2max}: Percentage of the running speed predicted to elicit VO_{2max}, determined during a pre-training incremental test [46].

GET: Gas exchange threshold as determined from a cluster of ventilatory measurements taken during a pre-training incremental test.

NC: Non-controlled study.

C: Controlled study.

Non-Ath: Non-athlete.

Sed: Sedentary.

Ath: Athlete.

HIT: Low-volume high-intensity interval training group.

CON: Control group.

END: Endurance training group.

VO_{2max}: Maximal oxygen uptake.

Rep: Repetition.

-: Not applicable.

?: data not available.

SE: Standard error.

Table 2. Study and subject characteristics for peak power output included in the meta-analysis. Studies are sorted from the largest to the smallest effects (intervention group) on peak power output.

Study (year)	Study design	Subjects	Age (y [mean])	Sample size	Proportion of males	Duration (weeks)	Total sessions	Group	Exercise intensity ^a	No. of reps			Rep duration (s)	Work/rest ratio	Effect on peak power (%)	
										Start	End	Total reps			Mean	SE
Stathis et al. [45]	NC	Non-Ath	22.1	8	0.75	7	21	HIT	All-out	3	10	153	30	0.13	17.1	2.9
Burgomaster et al. [52]	NC	Non-Ath	22.0	8	0.75	2	6	HIT	All-out	4	7	30	30	0.25	15.7	2.9
Siahkoughian et al. [21]	NC	Non-Ath	19.1	12	1.00	8	24	HIT	All-out	6	10	191	30	0.125	14.4	2.4
Forbes et al. [53]	NC	Non-Ath	21.0	7	0.57	2	6	HIT	All-out	4	6	30	30	0.13	11.4	3.1
MacDougall et al. [38]	NC	Non-Ath	22.7	12	1.00	7	21	HIT	All-out	4	10	174	30	0.13	10.6	2.4
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	12	HIT	All-out	3	4	48	30	0.13	10.3	2.7
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	12	HIT	125 %P _{max}	6	8	96	30	0.25	7.5	3.1
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	-	CON	-	-	-	-	-	-	-0.4	3.0
Harmer et al. [48]	NC	Non-Ath	22.0	7	1.00	7	21	HIT	All-out	4	10	174	30	0.14	9.4	3.1
Barnett et al. [33]	C	Non-Ath	21.2	8	1.00	8	24	HIT	All-out	3	6	108	30	0.17	7.7	2.9
Barnett et al. [33]	C	Non-Ath	21.2	8	1.00	8	-	CON	-	-	-	-	-	-	8.9	2.0
Siahkoughian et al. [21]	NC	Ath	19.4	12	1.00	8	24	HIT	All-out	6	10	191	30	0.125	7.9	2.4

Richards et al. [54]	NC	Non-Ath	29.0	12	0.42	2	6	HIT	All-out	4	7	30	30	0.13	6.9	2.4
McKenna et al. [24]	NC	Non-Ath	20.9	8	1.00	7	21	HIT	All-out	4	10	174	30	0.14	5.8	2.9
Esbjörnsson Liljedahl et al. [55]	NC	Non-Ath	25.0	10	0.00	4	12	HIT	All-out	3	3	27	30	0.03	5.6	2.6
Richards et al. [54]	NC	Non-Ath	25.0	11	0.27	2	6	HIT	All-out	4	7	30	30	0.13	5.5	2.5
Burgomaster et al. [41]	NC	Non-Ath	21.0	8	1.00	2	6	HIT	All-out	4	7	30	30	0.13	5.4	2.1
Whyte et al. [29]	NC	Sed	32.1	10	1.00	2	6	HIT	All-out	4	6	26	30	0.11	4.7	3.3
Allemeier et al. [23]	C	Non-Ath	22.7	11	1.00	6	15	HIT	All-out	3	3	45	30	0.03	3.0	2.5
Allemeier et al. [23]	C	Non-Ath	24.0	6	1.00	6	-	CON	-	-	-	-	-	-	3.0	3.3
Jansson et al. [56]	NC	Non-Ath	27.0	7	1.00	4	10	HIT	All-out	3	3	30	30	0.03	2.4	3.1
McKenna et al. [57]	NC	Non-Ath	18.8	6	1.00	7	21	HIT	All-out	4	10	174	30	0.13	2.3	3.3
Esbjörnsson Liljedahl et al. [55]	NC	Non-Ath	26.0	6	1.00	4	12	HIT	All-out	3	3	27	30	0.03	1.4	3.3

^a All-out: Encompasses intensities described as either “maximal” [53,24,55,57]; “sprints” [33]; “maximum efforts” [38]; “supramaximal” [23]; and “all-out” [21,28,29,41,45,48,52,54,56].

%P_{max}: Percentage of the final completed work rate maintained for 10 s during a pre-training incremental test.

NC: Non-controlled study.

C: Controlled study.

Non-Ath: Non-athlete.

Ath: Athlete.

Sed: Sedentary.

HIT: Low-volume high-intensity interval training group.

CON: Control group.

Rep: Repetition.

-: Not applicable.

SE: Standard error.

Table 3. Study and subject characteristics for mean power output included in the meta-analysis. Studies are sorted from the largest to the smallest effects (intervention group) on mean power output.

Study (year)	Study design	Subjects	Age (y [mean])	Sample size	Proportion of males	Duration (weeks)	Total sessions	Treatment group	Exercise intensity ^a	No. of reps			Rep duration (s)	Work/rest ratio	Effect on mean power (%)	
										Start	End	Total reps			Mean	SE
Siahkouhian et al. [21]	NC	Non-Ath	19.1	12	1.00	8	24	HIT	All-out	6	10	191	30	0.13	17.5	3.1
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	12	HIT	All-out	3	4	48	30	0.13	17.2	6.1
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	4	12	HIT	125 %P _{max}	6	8	96	30	0.25	11.2	6.7
Bayati et al. [28]	C	Non-Ath	25.0	8	1.00	-	-	CON	-	-	-	-	-	-	0.3	4.4
Siahkouhian et al. [21]	NC	Ath	19.4	12	1.00	8	24	HIT	All-out	6	10	191	30	0.13	11.1	3.1
Stathis et al. [45]	NC	Non-Ath	22.1	8	0.75	7	21	HIT	All-out	3	10	153	30	0.13	10.7	3.8
Esbjörnsson Liljedahl et al. [55]	NC	Non-Ath	25.0	10	0.00	4	12	HIT	All-out	3	3	27	30	0.03	10.1	2.9
Burgomaster et al. [41]	NC	Non-Ath	21.0	8	1.00	2	6	HIT	All-out	4	7	30	30	0.13	8.7	2.9
Barnett et al. [33]	C	Non-Ath	21.2	8	1.00	8	24	HIT	All-out	3	6	108	30	0.17	7.1	1.6
Barnett et al. [33]	C	Non-Ath	21.2	8	1.00	-	-	CON	-	-	-	-	-	-	2.8	1.0
McKenna et al. [24]	NC	Non-Ath	20.9	8	1.00	7	21	HIT	All-out	4	10	174	30	0.14	6.1	3.8
Allemeier et	C	Non-Ath	22.7	11	1.00	6	15	HIT	All-out	3	3	45	30	0.03	5.0	3.2

al. [23]																	
Allemeier et al. [23]	C	Non-Ath	24.0	6	1.00	6	-	CON	-	-	-	-	-	-	5.0	4.4	
Forbes et al. [53]	NC	Non-Ath	21.0	7	0.57	2	6	HIT	All-out	4	6	30	30	0.13	4.5	1.3	
Richards et al. [54]	NC	Non-Ath	25.0	11	0.27	2	6	HIT	All-out	4	7	30	30	0.13	3.9	3.2	
Whyte et al. [29]	NC	Sed	32.1	10	1.00	2	6	HIT	All-out	4	6	26	30	0.11	3.6	1.5	
Jansson et al. [56]	NC	Non-Ath	27.0	7	1.00	4	10	HIT	All-out	3	3	30	30	0.03	2.8	4.1	
Esbjörnsson Liljedahl et al. [55]	NC	Non-Ath	26.0	6	1.00	4	12	HIT	All-out	3	3	27	30	0.03	1.4	3.7	
Richards et al. [54]	NC	Non-Ath	29.0	12	0.42	2	6	HIT	All-out	4	7	30	30	0.13	1.1	3.1	

^a All-out: Encompasses intensities described as either “maximal” [24,53,55]; “sprints” [33]; “supramaximal” [23]; and “all-out” [21,28,29,41,45,54,55].

%P_{max}: Percentage of the final completed work rate maintained for 10 s during a pre-training incremental test.

NC: Non-controlled study.

C: Controlled study.

Non-Ath: Non-athlete.

Ath: Athlete.

Sed: Sedentary.

HIT: Low-volume high-intensity interval training group.

CON: Control group.

Rep: Repetition.

-: Not applicable.

SE: Standard error.

Table 4. Effects of low-volume HIT on VO_{2max} following reference training,^a with modifying effects for study characteristics, subject characteristics and training parameters.

	Effect on VO_{2max}		Inference
	(%)		
	Mean	±90% CL	
<i>Effect on treatment groups^b</i>			
Sedentary males	10.0	±5.1	Possibly moderate ↑
Sedentary females	7.3	±4.8	Likely small ↑
Active non-athletic males	6.2	±3.1	Likely moderate ↑
Active non-athletic females	3.6	±4.3	Possibly moderate ↑
Athletic males	2.7	±4.6	Unclear
Controls	1.2	±2.0	Unclear
<i>Modifying effects</i>			
Baseline VO_{2max} lower by 10 mL•kg ⁻¹ •min ⁻¹	3.8	±2.5	Possibly moderate ↑
Athlete vs active non-athlete	2.4	±5.7	Unclear
3-fold increase in work/rest ratio	-0.3	±2.0	Unclear
3-fold increase in no. of sessions	-0.3	±2.0	Unclear
Uncontrolled study	-0.7	±2.3	Unclear
Sedentary vs active non-athlete	-2.2	±5.7	Unclear
Females	-2.5	±4.1	Unclear
Replacement of training (male athletes only)	-2.9	±5.3	Unclear

Effects on treatment groups are presented as intervention minus control.

^a Reference training: a controlled study of 13 low-volume HIT sessions with a work/rest ratio of 0.16. (0.14 for sedentary females)

^b Active non-athletic males: baseline VO_{2max} adjusted to 45 mL•kg⁻¹•min⁻¹.

Sedentary males: baseline VO_{2max} adjusted to 30 mL•kg⁻¹•min⁻¹.

Active non-athletic females: baseline VO_{2max} adjusted to 45 mL•kg⁻¹•min⁻¹.

Sedentary females: baseline VO_{2max} adjusted to 30 mL•kg⁻¹•min⁻¹.

Athletic males: baseline VO_{2max} adjusted to 60 mL•kg⁻¹•min⁻¹.

HIT: low-volume high-intensity interval training.

VO_{2max} : maximal oxygen uptake.

CL: confidence limits.

↑: increase.

Table 5. Effects of low-volume HIT on 30-s Wingate peak sprint power following reference training,^a with modifying effects for study characteristics, subject characteristics and training parameters.

	Effect on peak power (%)		Inference
	Mean	±90% CL	
<i>Effect on treatment groups^b</i>			
Males	1.8	±5.0	Unclear
Controls	4.5	±3.8	Possibly moderate ↑
<i>Modifying effects</i>			
5-fold increase in work/rest ratio	5.7	±3.5	Possibly moderate ↑
3-fold increase in sessions	2.9	±3.5	Likely small ↑
Females	2.0	±6.3	Unclear
Baseline peak power lower by 5 W/kg	1.6	±3.2	Unclear
Uncontrolled study	-1.0	±3.7	Unclear

Effects on treatment groups are presented as intervention minus control.

^a Reference training: a controlled study of 12 low-volume HIT sessions with a work/rest ratio of 0.10.

^b Males, with baseline peak power output adjusted to 11.5 W/kg.

HIT: low-volume high-intensity interval training.

CL: confidence limits.

↑: increase

Table 6. Effects of low-volume HIT on 30-s Wingate mean sprint power following reference training,^a with modifying effects for study characteristics, subject characteristics and training parameters.

	Effect on mean power (%)		Inference
	Mean	±90% CL	
<i>Effect on treatment groups^b</i>			
Males	2.2	±10.3	Unclear
Controls	2.8	±8.2	Unclear
<i>Modifying effects</i>			
3-fold increase in sessions	6.2	±3.9	Possibly moderate ↑
Baseline mean power lower by 4 W/kg	2.3	±3.7	Possibly small ↑
5-fold increase in work/rest ratio	1.5	±3.7	Unclear
Females	-0.1	±6.9	Unclear
Uncontrolled study	-2.3	±4.3	Unclear

Effects on treatment groups are presented as intervention minus control.

^a Reference training: a controlled study of 14 low-volume HIT sessions with a work/rest ratio of 0.09.

^b Males, with baseline mean power output adjusted to 7.6 W/kg.

HIT: low-volume high-intensity interval training.

CL: confidence limits.

↑: increase.

