

# Microbial Aspects of Atrazine Degradation in Natural Environments

T. Komang Ralebitso,<sup>1\*</sup> Eric Senior<sup>1,3</sup> and Henk W. van Verseveld<sup>2</sup>

<sup>1</sup>*International Centre for Waste Technology (Africa), School of Applied Environmental Sciences, University of Natal, P/B X01, Scottsville, 3209, South Africa*

<sup>2</sup>*Department of Molecular Cell Physiology, Vrije Universiteit, De Boelelaan 1087, NL-1081 HV Amsterdam, The Netherlands*

<sup>3</sup>*Current address: Environmental Technology Institute, Innovation Centre (NTU), 18 Nanyang Drive, Singapore 637723*

---

\* *Author for correspondence:* Environmental Technology Institute, Innovation Centre (NTU), 18 Nanyang Drive, Singapore 637723. E-mail: [tralebitso@yahoo.com](mailto:tralebitso@yahoo.com)

## **Abstract**

The potential toxicity of the *s*-triazine herbicide atrazine motivates continuous bioremediation-directed research. Several indigenous soil atrazine-catabolizing microbial associations and monocultures have been enriched/isolated from compromised sites. Of these, *Pseudomonas* sp. strain ADP has become a reference strain and has been used to elucidate sequences of the catabolic enzymes *atzA*, *atzB*, *atzC* and *atzD* involved in one aerobic degradation pathway and develop probes for the genes which encode these enzymes. Despite this, hitherto unknown or novel microorganisms, with unique sequences and different enzyme-mediated operative pathways, warrant continued investigations for effective site bioremediation. Also, the sustained effectiveness of natural attenuation must be demonstrated continually so regular site evaluations and results analyses, despite the limitations of chemical extraction methodologies, are crucial practices. For both directed and intrinsic bioremediation monitoring, traditional microbial association studies must be complemented by more advanced physiological and molecular approaches. The occurrence of catabolic plasmids, in particular, should be probed with DNA hybridization techniques. Also, PCR-DGGE and subsequent new sequence elucidation should be used prior to developing new primers for DNA sequences encoding novel catabolic enzymes, and for hybridization probe development, to establish the degradative potential of a compromised site, or adoption of FISH to, for example, monitor bioaugmented remediation.

*Key words:* atrazine, catabolic monocultures/microbial associations, molecular techniques

## Introduction

Although atrazine is an effective herbicide, extensive toxicological investigations (Biradar & Rayburn 1995; Allran & Karasov 2000) have motivated and continue to motivate bioremediation-directed research. Successful enrichment/isolation and characterization of catabolic microbial associations and monocultures have resulted and *Pseudomonas* sp. strain ADP has been used both to study atrazine catabolism under different conditions and elucidate the enzymes involved in one of the aerobic catabolic pathways.

Until recently, due to technique limitations, an understanding of microbial community development in natural ecosystems exposed to different inorganic and organic pollutants was restricted to culturable microorganisms because of technique limitations. However, by complementing traditional microbial association studies with more advanced physiological and molecular approaches, such as community-level physiological profiling (CLPP) (Garland 1997), phospholipid fatty acids analysis (PLFA) (Green & Scow 2000), denaturing- and temperature-gradient gel electrophoresis (DGGE/TGGE) (Heuer and Smalla, 1997) and fluorescent *in situ* hybridization (FISH) (Amann *et al.* 1995), efficient microbial species monitoring in both directed and intrinsic bioremediation should result. Currently, primers for the *atzABC* enzymes elucidated in *Pseudomonas* sp. strain ADP are used to establish the degradative potential of atrazine compromised sites (Shapir *et al.* 2000). However, hitherto unknown or novel microorganisms, with possible unique sequences and different enzyme-mediated operative pathways, warrant continued investigation for effective site bioremediation.

The marked differences of soils prevent generalizations on the rate and extent of sequestration and, thus, the bioavailability of organic compounds *in situ* (Chung and Alexander 1998). It is essential, therefore, to elucidate the adsorption/desorption characteristics of contaminant molecules in the specific soil to be remediated in relation to the enriched catabolic microbial associations in particular. Also, the sustained effectiveness of intrinsic attenuation must be demonstrated continually to satisfy increasingly stringent regulations (Rittmann 2000). Regular site evaluations and results analyses, despite the limitations of chemical extraction methodologies (Shows & Olesik 2000), are, therefore, crucial.

This review, while not exhaustive due to space limitations, discusses atrazine occurrence and availability in natural environments, particularly soil, and microbial catabolic

potential distribution and its elucidation by molecular techniques as a prerequisite for field-scale bioremediation.

### **The atrazine molecule: Impacts on health and the environment**

Atrazine [2-chloro-4-(ethylamino)-6-(*isopropylamino*)-s-triazine] is one of the most environmentally-prevalent s-triazine-ring herbicides through its global use to control, through photosystem II inhibition, pre- and post-emergence broadleaf and grassy weeds in major crops such as maize (*Zea mays*), sorghum (*Sorghum* spp) and sugarcane (*Saccharum officinarum*) (Pick *et al.* 1992; Seiler *et al.* 1992; Mandelbaum *et al.* 1993; Yanze-Kontchou & Gschwind 1994; Sparling *et al.* 1998). The United States Environmental Protection Agency (EPA) (1991) reported that since the 1980s, the annual use of all pesticides had been holding steady at 500 million kg of active ingredients. More specifically, 363 million kg of atrazine were applied between 1980 and 1990 in the USA alone (Yanze-Kontchou & Gschwind 1995).

The herbicide has been classified a Class C/possible human carcinogen (Loprieno *et al.* 1980). This classification was confirmed by Biradar & Rayburn (1995) who observed chromosomal damage of Chinese hamster ovary cells exposed to atrazine (0.005 - 0.080  $\mu\text{M}$ ) for two days. In the same study, analysis of Illinois public water supplies revealed that EPA “safe” concentrations demonstrated the potential of the pesticide as a clastogen. Atrazine dealkylation metabolites, such as deethylatrazine [2-chloro-4-amino-6-*isopropylamine*-1,3,5-triazine] and *deisopropylatrazine* [2-chloro-4-ethylamino-6-amino-1,3,5-triazine], are also regulated compounds and may pose health risks thus warranting investigation of their fates (Brouwer *et al.* 1990; Kolpin *et al.* 1998).

The herbicide has a solubility of approximately 30  $\text{mg l}^{-1}$  and a half-life in soil of between 15 and 100 days (Protzman *et al.* 1999). Atrazine decomposition products are relatively persistent in soil and their detection in surface- and ground-water has prompted some environmental concerns (Sparling *et al.* 1998; Gebendinger & Radosevich 1999).

Through a combination of proven catabolic potential, relatively low cost and minimal environmental impact, bioremediation has been espoused largely as the technology of choice for atrazine-compromised sites despite the availability of various alternative treatment and containment methods such as: ozonation (Ma & Graham 2000); photochemical degradation (Konstantinou *et al.* 2001); diatomaceous earth remediation (Agdi *et al.* 2000); and powdered activated carbon adsorption (Campos *et al.* 2000).

## **Pesticide soil sorption properties and fates**

Pesticide fates in soil, and surface- and ground-water, including irreversible binding and persistence, are governed by interactions between retention, transformation and transport processes. Sorption affects these processes and determines, ultimately, pesticide volatilization, bioavailability, degradation and bioaccumulation. Desorption processes are equally important particularly for quantifying pesticide transport and, therefore, establishing a mass balance (Gao *et al.* 1998; Mata-Sandoval *et al.* 2000).

Several researchers have explored the importance of soil organic matter modification, via, for example, treated sewage sludge application (Masaphy & Mandelbaum 1997; Slusznny *et al.* 1999), colloids (Sprague *et al.* 2000), van der Waals forces, hydrogen bonds, charge transfer and triazine protonation (Moreau-Kervévan & Mouvet 1998), particle size, “ageing” and pH (Huang *et al.* 1984; Roy & Krapac 1994) on the sorption and, thus, microbial catabolism of atrazine. Erickson & Lee (1989) and Moreau-Kervévan & Mouvet (1998) ascribed atrazine adsorption to humic substances, especially humic acids, clays and oxyhydroxides, and to the various interrelated physical and chemical mechanisms of soil which are dependent upon the solute and key environmental determinants such as pH and temperature.

Low atrazine biodegradation rates are effected by its low water solubility ( $\pm 0.15\text{mM}$ , Erickson & Lee 1989) and sorption to regions of the soil particles, including the pores which are inaccessible to microbial species. Enhanced water solubility, increased mass transfer to the water phase and, thus, increased catabolic rates of the herbicide may be facilitated by the addition of surfactants, such as sodium dodecyl sulphate (Mata-Sandoval *et al.* 2000; Sanchez-Camazano *et al.* 2000), and/or inorganic nutrients, via percolation. Although effective, synthetic surfactants have some limitations such as: inhibition of microbial activity on micellar phase-solubilized pollutants (Mata-Sandoval *et al.* 2000); bactericidal activity; and considerable affinity to soil surfaces thus accumulating in and polluting subsurface environments (Chin *et al.* 1996). To circumvent these, biosurfactants, such as atrazine-solubilizing rhamnolipids, may be considered (Mata-Sandoval *et al.* 2000) since they have been shown to effect slightly increased atrazine solubility compared with the synthetic surfactant Triton X-100. The potential impacts of surfactants (and solvents such as methanol) on atrazine catabolism must, however, be evaluated particularly in relation to their preferential use as carbon sources.

## Atrazine biodegradative capacity and distribution

### *Catabolism under carbon- and nitrogen-limited conditions*

As a relatively new and complex xenobiotic molecule, atrazine represents a considerable challenge for microbial catabolism *in situ*. Its worldwide use may, however, have resulted in the global distribution of known microorganisms, with newly-induced catabolic enzymes, or novel microorganisms, with atrazine-catabolic capacities. Also, despite its limitations as an energy source, due to full oxidation of the ring carbons (Radosevich *et al.* 1995), its susceptibility to catabolism is improved by the presence of both carbon and nitrogen with the herbicide targeted as a nitrogen or carbon source in N- and C-limited conditions, respectively.

Laboratory and field studies by Mandelbaum *et al.* (1995), Shapir *et al.* (1998) and Katz *et al.* (2000) showed increased population sizes and herbicide degradation with the provision of tri-sodium citrate as a supplementary carbon source. However, several authors (Assaf & Turco 1994b; Grigg *et al.* 1997; Ames & Hoyle 1999; Gebendinger & Radosevich 1999) have reported that the concentrations and forms of C and N are major determinants of atrazine mineralization. For example, inorganic nitrogen rather than organic nitrogen (plant debris/*in situ* soil organic matter) stimulated pesticide catabolism (Alvey & Crowley 1995) and glucose addition of  $\leq 16\text{gCkg}^{-1}$  soil facilitated the formation of bound atrazine residues and, thus, reduced its bioavailability while  $\geq 16\text{gCkg}^{-1}$  soil enhanced dealkylation (Abdelhafid *et al.* 2000). Further work is, therefore, required before nitrogen and/or carbon supplementations can be introduced on a routine basis into site bioremediation strategies.

### *Enrichment and isolation of catabolic microorganisms*

Several researchers have reported the enrichment and isolation from compromised sites in different geographical regions (Radosevich *et al.* 1995; Bouquard *et al.* 1997; Sparling *et al.* 1998; Topp *et al.* 2000b) of microorganisms which are able to: dealkylate atrazine in a carbon-limited medium (Bheki & Kahn 1986); mineralize and use atrazine as a sole carbon and energy source (Mandelbaum *et al.* 1995; Yanze-Kontchou & Gschwind 1995; Topp *et al.* 2000b); utilize the heterocyclic nitrogen (Bichat *et al.* 1999); and, in the presence of supplemental carbon, mineralize atrazine and its metabolites as a source of nitrogen (Mandelbaum *et al.* 1993; Topp *et al.* 2000a). Successful mineralization of  $>94\%$  of atrazine-C ( $50\ \mu\text{gml}^{-1}$ ) and the isolation of a microorganism, *Agrobacterium radiobacter* strain J14a,

which was able to dealkylate, dehalogenate and mineralize the s-triazine ring when the molecule was used as a nitrogen source, was reported by Struthers *et al.* (1998). By screening some *Rhodococcus* strains which were known to be ubiquitous in soil and had diverse biodegradative capabilities, Behki *et al.* (1993) identified the strain TE1 with a 77-kb plasmid with atrazine-degrading capacity. Other workers reported microbial associations (Mandelbaum *et al.* 1993; Assaf & Turco 1994a; b; Alvey & Crowley 1996; de Souza *et al.* 1998a; Ralebitso *et al.* 1999) and fungal species (Donnelly *et al.* 1993; Mougín *et al.* 1994) which mineralized atrazine.

Culturable soil atrazine-catabolizing microbial associations/monocultures show a preponderance of rod-shaped bacteria (Table 1). Of these, *Pseudomonas* sp. strain ADP, which is capable of catabolizing the herbicide at concentrations  $>1\ 000\text{mg l}^{-1}$ , has become a reference strain and has been used extensively to: study atrazine catabolism under anoxic or denitrifying conditions (Shapir *et al.* 1998; Katz *et al.* 2000); demonstrate efficient bioaugmentation of an atrazine-contaminated soil (Newcombe & Crowley 1999); examine a method for aquifer amelioration *in situ* (Shati *et al.* 1996); explore atrazine degradation in soil, in comparison to other monocultures (Topp 2001); elucidate sequences of the aerobic catabolic enzymes *atzA* (de Souza *et al.* 1996), *atzB* (Boundy-Mills *et al.* 1997) and *atzC* (Sadowsky *et al.* 1998) and develop probes for their encoding genes (de Souza *et al.* 1998b).

Table 1: Some isolated atrazine-catabolizing bacterial strains

Strain	End products	Mineralization	Reference
<i>Pseudomonas</i> sp.	Deethylatrazine deisopropylatrazine	+ -	Behki & Khan (1986)
<i>Rhodococcus</i> sp. strain B-30		-	Behki & Khan (1994)
<i>Pseudomonas</i> sp. strain YAYA6	CO <sub>2</sub>	+	Yanze-Kontchou & Gschwind (1994)
<i>Pseudomonas</i> sp. strain ADP	CO <sub>2</sub> + NH <sub>4</sub> <sup>+</sup>	+	Mandelbaum <i>et al.</i> (1995)
<i>Rhodococcus</i> sp. strain N186/21	Hydroxyisopropylatrazine	-	Nagy <i>et al.</i> (1995)
<i>Ralstonia</i> M91-3	Biuret + CO <sub>2</sub> + NH <sub>4</sub> <sup>+</sup>	+	Radosevich <i>et al.</i> (1995)
<i>Rhizobium</i> sp.	Hydroxyatrazine	-	Bouquard <i>et al.</i> (1997)
<i>Agrobacterium radiobacter</i> J14A	CO <sub>2</sub>	+	Struthers <i>et al.</i> (1998)
<i>Clavibacter michiganese</i> ATZ1	Hydroxyatrazine + N-ethylammelide	-	de Souza <i>et al.</i> (1998a)
<i>Pseudomonas</i> sp. strain CN1	Cyanuric acid + CO <sub>2</sub>	+	de Souza <i>et al.</i> (1998a)
<i>Nocardioides</i> sp.	Hydroxyatrazine + N-ethylammelide	-	Topp <i>et al.</i> (2000a)
<i>Pseudoaminobacter</i> sp.	CO <sub>2</sub>	+	Topp <i>et al.</i> (2000b)

## Molecular microbial ecology of atrazine catabolism

Herbicide bioremediation has been researched extensively (Senior *et al.* 1976; de Souza *et al.* 1998a) although new microbial associations with high specific growth rates and/or high critical substrate concentrations may still be isolated. Characterization of such associations in relation to key environmental and physiological determinants will be needed then to facilitate the implementation of effective bioremediation protocols.

The study of nucleic acids extracted from environmental samples has allowed analysis independent of the problems which arise from laboratory degradative potential assays, laboratory culture-based enumerations, and our inability to culture the majority of species (Brockman 1995). Therefore, the need for broad-based, non-selective DNA extraction procedures or molecular techniques (*e.g.* DGGE/TGGE) and terminal-restriction fragment length polymorphism (T-RLFP)) which explore the complexities of microbial associations and their dynamics has been highlighted (Muyzer & Smalla 1998).

Ames & Hoyle (1999) concluded that chemical parameters, such as (sediment) N content or atrazine residues, could not be used alone as good primary predictors of biodegradative potential and/or activity. Therefore, understanding the distribution of contaminant-catabolizing microorganisms, genes or enzymes is of paramount importance (Mason *et al.* 1998). Thus, with the use of custom-designed primers, de Souza *et al.* (1998b) identified homology to the *Pseudomonas* sp. strain ADP *atzABC* genes in five atrazine-catabolizing microbial isolates and suggested that the genes were widespread. Shapir *et al.* (1998; 2000) used magnetic capture hybridization-polymerase chain reaction followed by nested PCR to detect atrazine-catabolizing genotypes and ascertain the expressed level of biodegradative activity *in situ* without the need to cultivate or activate the indigenous catabolic species. Generally, the mineralization rate and proliferation of catabolic microorganisms was correlated positively to the presence and copy number of the atrazine-catabolic genes. Therefore, the absence of the atrazine-dechlorinating gene *atzA* limited, possibly, intrinsic bioremediation (Shapir *et al.* 1998) while its presence (100 copies g<sup>-1</sup> soil) in soils with histories of atrazine contamination conferred short (4-5 day) lag phases before 40-54% mineralization of the herbicide (Shapir *et al.* 2000).

To date, several summaries of suggested catabolic pathways of atrazine and related s-triazines have been published (Cook 1987; Erickson & Lee 1989; [http://www.labmed.umn.edu/umbbd/atr/atr\\_map.html](http://www.labmed.umn.edu/umbbd/atr/atr_map.html) 20-12-2001). Of the pathways, one has been elucidated fully in *Pseudomonas* sp. strain ADP (de Souza *et al.* 1998a) and is known to

be hydrolytic rather than oxidative and consists of four main steps: dehalogenation; N-dealkylation; deamination; and ring cleavage (Figure 1). The catabolic enzymes were identified by de Souza *et al.* (1996), Boundy-Mills *et al.* (1997) and Sadowsky *et al.* (1998). de Souza *et al.* (1998c) then identified the location of the genes *atzABC* which encode these enzymes on a self-transmissible plasmid. Topp *et al.* (2000a) reported an alternative pathway with a new hydrolase for the degradation of a range of *s*-triazines by a *Nocardioides* sp. Also,




Figure 1: Aerobic atrazine catabolic pathway elucidated with *Pseudomonas* sp. strain ADP (de Souza *et al.* 1998a).

a dechlorinating enzyme, *s*-triazine hydrolase (*trzA*), and a gene controlling N-dealkylation have been characterized for *Rhodococcus corallinus* (Mulbry, 1994; Shao *et al.* 1995) and the involvement of the cytochrome P-450 system in atrazine N-dealkylation in *Rhodococcus* sp. strain N186/21 explored (Nagy *et al.* 1995). Nonetheless, new species and novel catabolic mechanisms may await identification and elucidation.

Together with methods to promote indigenous microbial activity, the introduction of genetically-engineered strains (bioaugmentation), which must be critically monitored/tracked, may be considered. Shao *et al.* (1995) and Strong *et al.* (2000), for example, conferred catabolic capacity by constructing recombinant *Rhodococcus* strains and *Escherichia coli* which were capable of dealkylating and dechlorinating atrazine. While the longevity and survival of the genetically-modified strains may be limited, transfer of the catabolic genes, particularly if located on plasmids, to robust and acclimated indigenous microorganisms may facilitate atrazine catabolism. Central to the adoption of this practice would be consideration of critical public and ethical concerns.

### Potential applications: Adopting a multi-disciplinary approach

Despite recommendations for controlled and managed herbicide applications (Gerstl *et al.* 1998), atrazine usage will probably continue and will result in further compromised soil and surface- and ground-water. While widely accepted that *in situ* treatments rarely yield

undesirable by-products, precautions and preliminary baseline tests (detailed contaminated site characterization in relation to the pollutant, hydrogeochemistry and microbiology) are always recommended for successful implementation (Mason *et al.* 1998; Haack & Bekins 2000). Also, species monitoring is necessary for public health considerations and for following the activities of microorganisms with genes encoding specific catabolic activities. Therefore, since DGGE/TGGE and PLFA patterns lend themselves to artificial neural network scrutiny (Almeida *et al.* 1999; Noble *et al.* 2000), they can be used, following specific catabolic association profile characterization, with different bioremediation strategies.

DNA hybridization techniques may be used to elucidate plasmid-borne catabolic sequences. Also, PCR-DGGE and subsequent new sequence elucidation may be used prior to developing new primers for chromosomal sequences encoding novel catabolic enzymes, and for hybridization probe development to establish the degradative potential of a compromised site or, by use of FISH, to monitor bioaugmented remediation. Direct hybridization avoids the possible biases introduced by DNA extraction and cleanup methods, which can distort the visualized community structure and component species dominance (Martin-Laurent *et al.* 2001), and preferential PCR amplification of the more abundant species (Head *et al.* 1998). Instead, whole cell *in situ* hybridization techniques facilitate quantitative estimation. This can give absolute enumeration and identify spatial localization and distribution of the metabolic activities of specific microorganisms in natural environments (Amann *et al.* 1995).

Together with measuring the occurrence of genes which encode, for example, atrazine catabolic enzymes, gene probe technology may be used to identify the concentration of the herbicide which effects changes in a microbial association and soils that have been affected by exposure to the molecule (Guo *et al.* 1997).

Techniques such as PLFA may be used to both estimate the cellular viability, nutritional and physiological status (assuming rapid degradation of intact phospholipids upon cell death) and fingerprint the microbial community configurations *in situ* (Parkes 1987; Green & Scow 2000). However, the constraint associated with Gram-negative bacteria community structure characterization, where the dominance and broad distribution of monoenoic, saturated and cyclopropane fatty acids limit differentiation or species subdivision (Zelles 1997), should be considered. Also, as exemplified by El Fantroussi *et al.* (1999), bacterial diversity in herbicide-contaminated soils might be studied by the combined approach of molecular (DGGE analysis) and culture-based methods (Biolog assay) despite the well-documented limitations of each technique.

The sustained effectiveness of intrinsic and enhanced (bioaugmented/biosupplemented) remediations must be demonstrated continually so regular site evaluations and results analyses, despite the limitations of chemical extraction methodologies, are crucial. This is particularly important since the contaminant concentration limits of current regulations are becoming increasingly stringent as exemplified by the  $0.1 \mu\text{g l}^{-1}$  allowed in Europe for a single pesticide in drinking water (Martín-Esteban *et al.* 1997). In contrast, the South African Water Quality Guidelines (1996) stipulated an atrazine concentration of  $\leq 2 \mu\text{g l}^{-1}$  in domestic water (Anon 1996). The routine use of sensitive, rapid and inexpensive detection and screening methods, such as immunoassays/immunosensors (Turiel *et al.* 1999), improved and more sophisticated and upscaled pollution (water) analytical techniques, such as immunoaffinity chromatography (Martín-Esteban *et al.* 1997), and extraction methods to predict bioavailability (Chung & Alexander 1998) are, therefore, central to environmental monitoring/protection.

For atrazine-contaminated surface- and ground-water, novel enzymes of inducible putative (Cook 1987) or alternative constitutive catabolic pathways could be over-expressed in killed whole-cell suspensions of recombinant microorganisms (Strong *et al.* 2000) and/or isolated and exploited via immobilization on an insoluble cellulose matrix (Kauffmann *et al.* 2000), as was done with atrazine chlorohydrolase (AtzA).

In general, a combination of conventional culture-dependent microbial techniques, various relevant molecular microbial ecology approaches and improved pollutant extraction and analysis methods should underpin all bioremediation protocols and so facilitate a comprehensive Code of Practice for the bioremediation of atrazine-compromised soil, surface- and ground-water.

## References

- Abdelhafid R, Houot S & Barriuso E (2000) How increasing availabilities of carbon and nitrogen affect atrazine behaviour in soils. *Biology and Fertility of Soils* 30: 333-340
- Agdi K, Bouaid A, Esteban AM, Hernando PF, Azmani A & Camara C (2000) Removal of atrazine and four organophosphorus pesticides from environmental waters by diatomaceous earth-remediation method. *Journal of Environmental Monitoring* 2: 420-423

- Allran JW & Karasov WH (2000) Effects of atrazine and nitrate on northern leopard frog (*Rana pipiens*) larvae exposed in the laboratory from posthatch through metamorphosis. *Environmental Toxicology and Chemistry* 19: 2850-2855
- Almeida JS, Barreto CMT, Figueiredo MJJ, Noble PA, MacNaughton SJ, Stephen JR, White DC and Carrondo MJT (1999) Microbial typing for management of remediation in contaminated soils. Proceedings of the 1999 African International Environmental Protection Symposium, (CD-ROM), Pietermaritzburg
- Alvey S & Crowley DE (1995) Influence of organic amendments on biodegradation of atrazine as a nitrogen source. *Journal of Environmental Quality* 24: 1156-1162
- Alvey S & Crowley DE (1996) Survival and activity of an atrazine-mineralizing bacterial consortium in rhizosphere soil. *Environmental Science and Technology* 30: 1596-1603
- Amann RI, Ludwig W & Schleifer KH (1995) Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiological Reviews* 59: 143-169
- Ames RA & Hoyle BL (1999) Biodegradation and mineralization of atrazine in shallow subsurface sediments from Illinois. *Journal of Environmental Quality* 28: 1674-1681
- Anon (1996) South African Water Quality Guidelines, Domestic Water Use, Department of Water Affairs and Forestry, Second Edition
- Assaf NA & Turco RF (1994a) Accelerated biodegradation of atrazine by a microbial consortium is possible in culture and soil. *Biodegradation* 5: 29-35
- Assaf NA & Turco RF (1994b) Influence of carbon and nitrogen application on the mineralization of atrazine and its metabolites. *Pesticide Science* 41: 41-47
- Behki RM & Khan SU (1986) Degradation by *Pseudomonas*: n-Dealkylation and dehalogenation of atrazine and its metabolites. *Journal of Agriculture and Food Chemistry* 34: 746- 749
- Behki RM & Khan SU (1994) Degradation of atrazine, propazine and simazine by *Rhodococcus* strain B-30. *Journal of Agriculture and Food Chemistry* 42: 1237-1241
- Behki R, Topp E, Dick W & Germon P (1993) Metabolism of the herbicide atrazine by *Rhodococcus* strains. *Applied and Environmental Microbiology* 59: 1955-1959
- Bichat F, Sims GK & Mulvaney RL (1999) Microbial utilization of heterocyclic nitrogen from atrazine. *Soil Science Society of America Journal* 63: 100-110

- Biradar DP & Rayburn AL (1995) Chromosomal damage induced by herbicide contamination at concentrations observed in public water supplies. *Journal of Environmental Quality* 24: 1222-1225
- Boundy-Mills K, de Souza ML, Mandelbaum RM, Wackett LP & Sadowsky MJ (1997) The *atzB* gene of *Pseudomonas* sp. strain ADP encodes the second enzyme of a novel atrazine degradation pathway. *Applied and Environmental Microbiology* 63: 916-923
- Bouquard C, Ouazzani J, Promé J-C, Michel-Briand Y & Plésiat P (1997) Dechlorination of atrazine by a *Rhizobium* sp. isolate. *Applied and Environmental Microbiology* 63: 862-866
- Brockman FJ (1995) Nucleic acid-based methods for monitoring the performance of *in situ* bioremediation. *Molecular Ecology* 4: 567-578
- Brouwer WWM, Boesten JJTI & Siegers WG (1990) Adsorption of transformation products of atrazine by soil. *Weed Research* 30: 123-128
- Campos C, Snoeyink VL, Marinas B, Baudin I & Laine JM (2000) Atrazine removal by powdered activated carbon in floc blanket reactors. *Water Research* 34: 4070-4080
- Chin Y-P, Kimble KD & Swank CR (1996) The sorption of 2-methylnaphthalene by Rossburg soil in the absence and presence of a nonionic surfactant. *Journal of Contaminant Hydrology* 22: 83-94
- Chung N & Alexander M (1998) Differences in sequestration and bioavailability of organic compounds aged in dissimilar soils. *Environmental Science and Technology* 32: 855-860
- Cook AM (1987) Biodegradation of *s*-triazine xenobiotics. *FEMS Microbiology Review* 46: 93-116
- de Souza ML, Newcombe D, Alvey S, Crowley DE, Hay A, Sadowsky MJ & Wackett LP (1998a) Molecular basis of a bacterial consortium: Interspecies catabolism of atrazine. *Applied and Environmental Microbiology* 64: 178-184
- de Souza ML, Sadowsky MJ & Wackett LP (1996) Atrazine chlorohydrolase from *Pseudomonas* sp. strain ADP: Gene sequences, enzyme purification, and protein characterization. *Journal of Bacteriology* 178: 4894-4900
- de Souza ML, Seffernick J, Martinez B, Sadowsky MJ & Wackett LP (1998b) The atrazine catabolism genes *atzABC* are widespread and highly conserved. *Journal of Bacteriology* 180:1951-1954

- de Souza ML, Wackett LP & Sadowsky MJ (1998c) The *atzABC* genes encoding atrazine catabolism are located on a self-transmissible plasmid in *Pseudomonas* sp. strain ADP. *Applied and Environmental Microbiology* 64: 2323-2326
- Donnelly PK, Entry JA & Crawford DL (1993) Degradation of atrazine and 2,4-dichlorophenoxyacetic acid by mycorrhizal fungi at three nitrogen concentrations *in vitro*. *Applied and Environmental Microbiology* 59: 2642-2647
- El Fantroussi S, Verschuere L, Verstraete W & Top EM (1999) Effect of phenylurea herbicides on soil microbial communities estimated by analysis of 16S rRNA gene fingerprints and community-level physiological profiles. *Applied and Environmental Microbiology* 65: 982-988
- Erickson EL & Lee KH (1989) Degradation of atrazine and related *s*-triazines. *Critical Reviews in Environmental Control* 19: 1-14
- Gao JP, Maguhn J, Spitzauer P & Kettrup A (1998) Sorption of pesticides in the sediment of the Teufelsweiher pond (Southern Germany). I: Equilibrium assessments, effect of organic carbon content and pH. *Water Research* 32: 1662-1672
- Garland JL (1997) Analysis and interpretation of community-level physiology profiles in microbial ecology. *FEMS Microbial Ecology* 24: 289-300
- Gebendinger N & Radosevich M (1999) Inhibition of atrazine degradation by cyanazine and exogenous nitrogen in bacterial isolate M91-3. *Applied Microbiology and Biotechnology* 51: 375-381
- Gerstl Z, Nasser A & Mingelgrin U (1998) Controlled release of pesticides into water from clay-polymer formulations. *Journal of Agriculture and Food Chemistry* 46: 3803-3809
- Green CT & Scow KM (2000) Analysis of phospholipid fatty acids (PLFA) to characterize microbial communities in aquifers. *Hydrogeology Journal* 8: 126-141
- Grigg BC, Bischoff M & Turco RF (1997) Cocontaminant effects on degradation of triazine herbicides by a mixed microbial culture. *Journal of Agriculture and Food Chemistry* 45: 995-1000
- Guo C, Sun W, Harsh JB & Ogram A (1997) Hybridization analysis of microbial DNA from fuel oil-contaminated and noncontaminated soil. *Microbial Ecology* 34: 178-187
- Haack SK & Bekins BA (2000) Microbial populations in contaminant plumes. *Hydrogeology Journal* 8: 63-76
- Head IM, Saunders JR & Pickup RW (1998) Microbial evolution, diversity and ecology: A decade of ribosomal analysis of uncultivated microorganisms. *Microbial Ecology* 35: 1-21

- Heuer H & Smalla K (1997) Application of denaturing gradient gel electrophoresis and temperature gradient gel electrophoresis for studying soil microbial communities. In: van Elsas JD, Trevor JT & Wellington EMH (Eds) *Modern Soil Microbiology*, (pp. 353-373). Marcel Dekker Inc., New York
- Huang PM, Grover R & McKercher RB (1984) Components and particle size fractions involved in atrazine adsorption by soils. *Soil Science* 28: 20-24
- Katz I, Green M, Ruskol Y & Dosoretz CG (2000) Characterization of atrazine degradation and nitrate reduction by *Pseudomonas* sp. strain ADP. *Advances in Environmental Research* 4: 219-224
- Kauffmann C, Shoseyov O, Shpigel E, Bayer EA, Lamed R, Shoham Y & Mandelbaum RT (2000) Novel methodology for enzymatic removal of atrazine from water by CBD-fusion protein immobilized on cellulose. *Environmental Science and Technology* 34: 1292-1296
- Kolpin DW, Thurman EM & Linhart SM (1998) The environmental occurrence of herbicides: The importance of degradates in ground water. *Archive of Environmental Contamination and Toxicology* 35: 385-390
- Konstantinou IK, AK Zarkadis & TA Albanis (2001) Photodegradation of selected herbicides in various natural waters and soils under environmental conditions. *Journal of Environmental Quality* 30: 121-130
- Loprieno N, Barale R, Mariani L, Presciuttini S, Rossi AM, Sbrana I, Zaccaro L, Abbondandolo A & Bonatti S (1980) Results of mutagenicity tests on the herbicide atrazine. *Mutation Research* 74: 250
- Ma J & Graham NJD (2000) Degradation of atrazine by manganese-catalysed ozonation – Influence of radical scavengers. *Water Research* 34: 3822-3828
- Mandelbaum RT, Allan BH & Wackett LP (1995) Isolation of a *Pseudomonas* sp. that mineralizes the *s*-triazine herbicide atrazine. *Applied and Environmental Microbiology* 61: 1451- 1457
- Mandelbaum RT, Wackett LP & Allan DL (1993) Mineralization of the *s*-triazine ring of atrazine by stable bacterial mixed cultures. *Applied and Environmental Microbiology* 59: 1695-1701
- Martín-Esteban A, Fernández P & Cámara C (1997) Immunosorbents: A new tool for pesticide sample handling in environmental analysis. *Fresenius' Journal of Analytical Chemistry* 357: 927-933

- Martin-Laurent F, Philippot L, Hallet S, Chaussod R, Germon JC, Soulas G & Catroux G (2001) DNA extraction from soils: Old bias for new microbial diversity analysis methods. *Applied and Environmental Microbiology* 67: 2354-2359
- Masaphy S & Mandelbaum RT (1997) Atrazine mineralization in slurries from soils irrigated with treated waste water. *Applied Soil Ecology* 6: 283-291
- Mason JR, Haider I, Henkler C & Fenwick C (1998) Assessment of biodegradative potential and activity in contaminated sites: Molecular approaches to environmental monitoring. *Biochemical Society Transactions* 26: 694-697
- Mata-Sandoval JC, Karns J & Torrents A (2000) Effects of rhamnolipids produced by *Pseudomonas aeruginosa* UG2 on the solubilization of pesticides. *Environmental Science and Technology* 34: 4923-4930
- Moreau-Kervévan C & Mouvet C (1998) Adsorption and desorption of atrazine, deethylatrazine and hydroxyatrazine by soil components. *Journal of Environmental Quality* 27: 46-53
- Mougin C, Laugero C, Asther M, Dubroca J, Frasse P & Asther M (1994) Biotransformation of the herbicide atrazine by the white rot fungus *Phanerochaete chrysosporium*. *Applied and Environmental Microbiology* 60: 705-708
- Mulbry WW (1994) Purification and characterization of an inducible *s*-triazine hydrolase from *Rhodococcus corallinus* NRRL B-15444R. *Applied and Environmental Microbiology* 60: 613-618
- Muyzer G & Smalla K (1998) Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie van Leeuwenhoek* 73: 127-141
- Nagy I, Compernelle F, Ghys K, Vanderleyden J & de Mot R (1995) A single cytochrome P-450 system is involved in degradation of the herbicides EPTC (*s*-ethyl dipropylthiocarbamate) and atrazine by *Rhodococcus* sp. strain N186/21. *Applied and Environmental Microbiology* 61: 2056-2060
- Newcombe DA & Crowley DE (1999) Bioremediation of atrazine-contaminated soil by repeated applications of atrazine-degrading bacteria. *Applied Microbiology and Biotechnology* 51: 877-882
- Noble PA, Almeida JS & Lovell CR (2000) Application of neural computing methods for interpreting phospholipid fatty acid profiles of natural microbial communities. *Applied and Environmental Microbiology* 66: 694-699

- Parkes RJ (1987) Analysis of microbial communities within sediments using biomarkers. In: Fletcher M, Gray TRG and Jones JG (Eds) Ecology of Microbial Communities, (pp. 147-177). Cambridge University Press, Cambridge
- Pick FE, van Dyk LP & Botha E (1992) Atrazine in ground and surface water in maize production areas of Transvaal, South Africa. *Chemosphere* 25: 335-341
- Protzman RS, Lee PH, Ong SK & Moorman TB (1999) Treatment of formulated atrazine rinsate by *Agrobacterium radiobacter* strain J14A in a sequencing batch biofilm reactor. *Water Research* 33: 1399-1404
- Radosevich M, Hao Y-L, Traina SJ & Tuovinen OH (1995) Degradation and mineralization of atrazine by a soil bacterial isolate. *Applied and Environmental Microbiology* 61: 297-302
- Ralebitso TK, Röling WFM, van Verseveld HW & Senior E (1999) A molecular approach to characterizing atrazine- and selected petroleum hydrocarbon-catabolizing microbial associations isolated from soil. Proceedings of the 1999 African International Environmental Protection Symposium, (CD-ROM), Pietermaritzburg
- Rittmann BE (2000) Natural attenuation's promise and application. *Water* 21 8: 20-22
- Roy WR & Krapac IG (1994) Adsorption and desorption of atrazine and deethylatrazine by 17 low organic carbon geologic materials. *Journal of Environmental Quality* 23: 549-556
- Sadowsky MJ, Tong Z, de Souza ML & Wackett LP (1998) AtzC is a new member of the amidohydrolase protein superfamily strain and is homologous to other atrazine-metabolizing enzymes. *Journal of Bacteriology* 180: 152-158
- Sanchez-Camazano M, Sanchez-Martin MJ & Rodriguez-Cruz MS (2000) Sodium dodecyl sulphate-enhanced desorption of atrazine: Effect of surfactant concentration and of organic matter content of soils. *Chemosphere* 41: 1301-1305
- Seiler A, Brenneisen P & Green DH (1992) Benefits and risks of plant protection products - possibilities of protecting drinking water: Case atrazine. *Water Supply* 10: 31-42
- Senior E, Bull AT & Slater JH (1976) Enzyme evolution in a microbial community growing on the herbicide Dalapon. *Nature (London)* 263: 476-479
- Shao ZQ, Seffens W, Mulbry W & Behki RM (1995) Cloning and expression of the *s*-triazine hydrolase gene (*trzA*) from *Rhodococcus corallinus* and development of *Rhodococcus* recombinant strains capable of dealkylating and dechlorinating the herbicide atrazine. *Journal of Bacteriology* 177: 5748-5755

- Shapir N, Goux S, Mandelbaum RT & Pussemier L (2000) The potential of soil microorganisms to mineralize atrazine as predicted by MCH-PCR followed by nested PCR. *Canadian Journal of Microbiology* 46: 425-432
- Shapir N, Mandelbaum RT and Jacobsen CS (1998) Rapid atrazine mineralization under denitrifying conditions by *Pseudomonas* sp. strain ADP in aquifer sediments. *Environmental Science and Technology* 32: 3789-3792
- Shati MR, Rönen D & Mandelbaum R (1996) Method for *in situ* study of bacterial activity in aquifers. *Environmental Science and Technology* 30: 2646-2653
- Shows ME & Olesik SV (2000) Extraction of atrazine and its metabolites using supercritical fluids and enhanced-fluidity liquids. *Journal of Chromatographic Science* 38: 399-408
- Sluszny C, Graber ER & Gerstl Z (1999) Sorption of *s*-triazine herbicides in organic matter amended soils: Fresh and incubated systems. *Water, Air, and Soil Pollution* 115: 395-410
- Sparling G, Dragten R, Aislabie J & Fraser R (1998) Atrazine mineralization in New Zealand topsoils and subsoils: Influence of edaphic factors and numbers of atrazine-degrading microbes. *Australian Journal of Soil Research* 36: 557-570
- Sprague LA, Herman JS, Hornberger GM & Mills AL (2000) Atrazine adsorption and colloid-facilitated transport through the unsaturated zone. *Journal of Environmental Quality* 29: 1632-1641
- Strong LC, McTavish H, Sadowsky MJ & Wackett LP (2000) Field-scale remediation of atrazine-contaminated soil using recombinant *Escherichia coli* expressing atrazine chlorohydrolase. *Environmental Microbiology* 2: 91-98
- Struthers JK, Jayachandran K & Moorman TB (1998) Biodegradation of atrazine by *Agrobacterium radiobacter* J14A and use of this strain in bioremediation of contaminated soil. *Applied and Environmental Microbiology* 64: 3368-3375
- Topp E (2001) A comparison of three atrazine-degrading bacteria for soil bioremediation. *Biology and Fertility of Soils* 33: 529-534
- Topp E, Mulbry WM, Zhu H, Nour SM & Cuppels D (2000a) Characterization of *s*-triazine herbicide metabolism by a *Nocardioides* sp. isolated from agricultural soils. *Applied and Environmental Microbiology* 66: 3134-3141
- Topp E, Zhu H, Nour SM, Houot S, Lewis M & Cuppels D (2000b) Characterization of an atrazine-degrading *Pseudaminobacter* sp. isolated from Canadian and French agricultural soils. *Applied and Environmental Microbiology* 66: 2773-2782

- Turiel E, Fernandez P, Perez-Conde C, Gutierrez AM & Camara C (1999) Oriented antibody immobilization for atrazine determination by flow-through fluoroimmunosensor. *Fresenius' Journal of Analytical Chemistry* 365: 658-662
- Yanze-Kontchou C & Gschwind N (1994) Mineralization of the herbicide atrazine as the carbon source by a *Pseudomonas* strain. *Applied and Environmental Microbiology* 60: 4297-4302
- Yanze-Kontchou C & Gschwind N (1995) Mineralization of the herbicide atrazine in soil inoculated with a *Pseudomonas* strain. *Journal of Agriculture and Food Chemistry* 43: 2291-2294
- Zelles L (1997) Phospholipid fatty acid profiles in selected members of soil microbial communities. *Chemosphere* 35: 275-294