

# HOSPITAL FOOD SERVICE QUALITY IMPROVEMENT QUESTIONNAIRE (HFSQIQ): DEVELOPMENT, TRANSLATION AND VALIDATION OF A QUESTIONNAIRE

**Osman NS<sup>1,2</sup>, Md Nor N<sup>2,3</sup>, Mustafa N<sup>1</sup>, Rusali R<sup>1</sup>, and Ruzana I<sup>4</sup>.**

<sup>1</sup>Department of Nutrition Sciences, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, 25200 Kuantan, Malaysia

<sup>2</sup>Centre for Nutrition and Dietetics Studies, Faculty of Health Sciences, Universiti Teknologi MARA, Puncak Alam Campus, 42300 Puncak Alam, Malaysia

<sup>3</sup>Integrative Pharmacogenomics Institute (iPROMISE), Universiti Teknologi MARA, Puncak Alam Campus, 42300 Puncak Alam, Malaysia

<sup>4</sup>School of Health and Life Sciences, Teesside University, TS1 3BX Middlesbrough, England, United Kingdom

## Correspondence:

Norazmir Md Nor,

Integrative Pharmacogenomics Institute (iPROMISE),

Universiti Teknologi MARA, Puncak Alam Campus,

42300 Puncak Alam, Malaysia.

Email: azmir2790@uitm.edu.my

## Abstract

The hospital food service department provides meals to staff, patients, and their caregivers while adhering to dietary therapy guidelines and promoting nutritional wellness. High-quality food service plays a pivotal role in offering inpatients nourishing meals that promote physical and mental well-being, aiding their recovery and overall health during their hospitalisation. This study aimed to develop and validate a tool for measuring and evaluating hospital food operations using the Total Quality Management approach. A literature review, in-depth interviews with food service employees, and a peer-review process were conducted to identify the domains and items for the questionnaire. A "Hospital Food Service Quality Improvement Questionnaire" (HFSQIQ) with 61 items in six domains was developed and the content validation was performed by seven experts. The questionnaire was translated into Malay, and the internal consistency of the HFSQIQ was examined using Cronbach's alpha. Resultantly, the HFSQIQ depicted high validity and reliability, with a high I-CVI and Kappa index rating for most items and a Cronbach alpha value of 0.97 and 0.98 for the importance and performance scales, respectively. In conclusion, the HFSQIQ is a useful tool for evaluating and improving the quality of hospital food service operations.

**Keywords:** Questionnaire Development, Questionnaire Validation, Reliability, Hospital Food Service, Quality Improvement, Total Quality Management

## Introduction

Hospitals in the healthcare sector all provide the same services, but the service quality differs. Given the growing competition, healthcare organisations have allocated top priority to service quality. A hospital food service is a department that caters for the food of staff, patients, and their caregivers by attending to a diet therapy regime and enhancing nutrition application (1). The quality of food service plays a significant role by offering inpatients nourishing meals that are beneficial physically and mentally to their health and recovery during hospital stays. The primary objective of a hospital food service is met when meals are meticulously planned and tailored to satisfy the patient's specific dietary requirements (2). Patients who are well-nourished upon admission have the right to maintain their current nutritional condition after discharge (3). Failure to provide acceptable quality food services may

result in poor food intake, thereby prolonging the recovery time, complication rates, and length of stay. These events may culminate in increased healthcare expenses, especially among the elderly (4).

Patient satisfaction is a widely accepted measure of food service quality, a key indicator. Food quality has been demonstrated as one of the most significant predictors of overall hospital stay satisfaction (5). Patients' nutritional status commonly deteriorates during hospitalisation (6, 7). The hospital food is often negatively perceived as cold, tasteless, poorly presented, and badly served (8). Besides, high patient satisfaction is influenced by several factors including appreciation towards meal services, staff interactions, and eating and physical eating environments (9). Patient satisfaction was associated with emotions, morals, medical discourse, and cultures of gratitude (10).

However, satisfaction assessment is typically limited to a few general questions about food service, which are insufficient to elicit information from patients about objectives and interpersonal aspects, and to examine patients' wishes for personalised service (11).

In Malaysia, most studies used instruments, such as surveys and measures of plate or food waste, adapted from earlier studies conducted in other countries to measure or assess patients' food intake and satisfaction with hospital foods (9, 12). A qualitative study found that emotions indirectly influenced patients' meal experience and food intake in hospitals (13). In addition, the staff play a primary role in providing a positive meal experience among patients during hospitalisation (14).

Various models have been developed to measure service quality. Total Quality Management (TQM) is one approach for fundamental measurement and continuous improvement. TQM is a concept rooted in the Japanese management style (15), which assists to improve the quality of services and goods through a collaborative approach and standardised performance. A previous study highlighted that all departments and individuals contribute to TQM in attaining standards on customer service and end-user satisfaction, which brings excellence to business (16). Nevertheless, there is insufficient information in the questionnaire designed for the Malaysian population. It is challenging to use the instrument in measuring food preferences due to differences in food culture and practices, which are linked to multi-ethnicity and diverse religions and socio-demographic backgrounds (17). Thus, this study aims to develop and validate a measurement tool using the TQM approach for monitoring and evaluating hospital food service operations.

## **Materials and Methods**

### **Questionnaire development**

In this study, the domain of the newly proposed questionnaire was identified through a literature review, in-depth interview, and peer review process. A literature search related to hospital food services quality improvement topic was used as a guide in generating the questionnaire's items for the measure (18-21). Next, the interview was conducted among 24 food service employees, and the acquired data were thematically analysed. The findings helped to determine which questions should be added to or removed from the initial questionnaire. The interviews also assisted to build and improve the answer options. Details on the thematic analysis of the interview session for the process of item development can be gleaned from a previous study (22). Next, the questionnaire underwent a peer-reviewed process to identify any overlapping and duplication questionnaire. A preliminary Hospital Food Service Quality Improvement Questionnaire (HFSQIQ) comprising 61 items generated from six domains were developed throughout this process.

### **Content validation**

The preliminary HFSQIQ underwent content validation, where a further judgement of the relevancy of items took place. According to Yusoff (23), the minimum number of experts for content validation should be six and the maximum number should not exceed ten. In addition, Polit et al. (24) and Polit and Beck (25) recommended using a CVI cut-off score of at least 0.83 with a minimum of six experts for the content validation procedure. Therefore, seven experts who were chosen from lecturers in food service or food technology, dietitians or catering dietitians, and catering officers working in the food service division of public and private hospitals for this study. The experts were selected based on criteria established by the researchers, such as their in-depth knowledge of scale development and/or the relevant domain, their dissimilarity to the individuals who developed the item pool, and the application of systematic expert judgement to avoid bias in the evaluation of items. The experts were specifically given the definition and the items describing the domain in the content validation form. Each domain and its underlying items were critically evaluated by the experts before scoring each item.

The experts had to provide written or verbal feedback to enhance the item's relevance to the intended domain. All feedback was considered when improving the domain and its item. Then, the experts were required to provide scores for each item individually based on the relevance scale following the domain and items review. Scales for evaluating relevance ranged from 1 = Not relevant to 4 = Highly relevant. To avoid having a neutral and ambiguous at the midpoint, Lynn (26), suggests adopting a 4-point rating scale instead of a 3- or 5-point rating scale (24). Each domain's item level (I-CVI) and scale level (S-CVI) content validity indexes were manually calculated. Two different indices were calculated to determine the S-CVI: 1) the percentage of items on one scale that an expert rated as valid (S-CVI/UA = universal agreement by the expert), and 2) the average percentage of items on one scale rated as valid (S-CVI/Ave = average agreement by experts) (27). Each item rated 3 or 4 was transformed to valid ('1'), whereas items with ratings of 1 or 2 were transformed to nonvalid ('0'). S-CVI/Ave was determined using two formulas:

$$I-CVI = (\text{agreed item}) / (\text{number of raters}); \text{ and}$$

$$S-CVI/Ave = (\text{summation all } I-CVI) / (\text{number of items})$$

All I-CVI values were initially obtained and divided by the number of items. The average percentage of each rater was then obtained using the second formula. Next, the number of items with 100% agreement was divided by the total number of items in that particular domain to determine S-CVI/UA (23). For the S-CVI/UA and S-CVI/Ave, a value of 0.8 was considered acceptable (24, 25). The items were revised or removed based on the panel's recommendations, and experts were contacted for second round of expert reviews to clarify any uncertainties.

### ***The translation process of the pre-final HFSQIQ from English into Malay***

The newly developed questionnaire, prepared in English, was translated into Malay following content validation. The purpose of the translation is to maintain the text's original meaning, style, and impact while translating the genuine context from English to Malay. The forward and backward translation process was adapted from Sousa and Rojjanasrirat guidelines (28).

The English version of the HFSQIQ was translated into Malay by two qualified independent translators. The first translator was familiar with healthcare terminology and the questionnaire's content in both languages. In contrast, the second translator was referred to as a naive translator who needed to be made aware of the objective the questionnaire intended to measure. The translators were instructed to independently prepare a forward translation version conceptually equivalent to the original HFSQIQ. A professional and qualified proof-reader then reviewed the translated questionnaire for inaccuracies. Lastly, the four members of the supervisory research team that are proficient in English and Malay examined and compared the questionnaire items translated into Malay to check for discrepancies between the translated text and the original language (English). To provide a preliminary forward translation of the HFSQIQ, the translations were reconciled to reach a consensus among research team members.

The next stage of the translation process entailed sending the preliminary initial forward translation Malay version of the preliminary HFSQIQ to a third translator. The third translator back-translated the tool from the target language (Malay) into the original language (English). To avoid reference to existing sources of teamwork assessment, the translator was not informed that the tool was being back-translated. After producing the backward translation version, research supervisory team members needed to reconcile the two English versions by comparing the backward translated version and the original version. Members of the supervisory research team were responsible for evaluating, revising, and consolidating the back-translated questionnaire to ensure conceptual, semantic, and content equivalence. They were also in charge of creating the pre-testing final target language questionnaire for pilot and psychometric testing (27). The members of the supervisory research team then discussed any disparities between the back-translated version and the original to guide the selection of phrases and words in the Malay version. All the comments and amendments made for the pre-final version of HFSQIQ were documented.

### ***Face validation of pre-final version in both the English and Malay versions of HFSQIQ***

Upon completing the content validation, 10 volunteer panels who were selected from the academician and healthcare professionals performed the face validation. The aim was to determine the clarity and understandability of

the translated items. Based on the comprehensibility and clarity of the source and translated items in the HFSQIQ questionnaire, the raters were instructed to provide a Likert scale score between 1 (item not clear and understandable) and 4 (item very clear and understandable). The ratings of 1 and 2 were then reclassified as 0 (not clear and understandable). Scores 3 and 4 were concurrently reclassified as 1 (clear and understood). The item-level face validity index (I-FVI) was computed using the raw scores for each item's comprehensibility and clarity in Microsoft Excel. In addition, the average of the I-FVI score (S-FVI/Ave) for all items on the scale or the average of proportional clarity and comprehension as evaluated by all raters was calculated (29). According to Marzuki et al. (30), the minimum acceptable number of raters for an online survey is 10, with acceptable FVI values of at least 0.83. The following formula was used:

$$I-FVI = (\text{agreed item}) / (\text{number of raters})$$

$$S-FVI/Ave = (\text{sum of I-FVI scores}) / (\text{number of items})$$

A modified Kappa index was also generated to estimate the I-CVI. The modified Kappa ( $k^*$ ) is an index of agreement among experts that demonstrates, beyond the possibility of random variation, that the item is relevant, clear, or possesses another quality of relevance. Polit et al. (24) formula was implemented to calculate the modified Kappa index in this study. For each item, the probability of chance agreement ( $P_c$ ) was first computed using the formula below:

$$P_c = [(N!/A!)(N - A)!] * 0.5N$$

Where  $N$  represents the total number of experts, and  $A$  represents the total number of experts or target users who agreed that the item was comprehensible, relevant, and clear. The next step was to calculate the Kappa value using the following formula:

$$k = (\text{item - level content validity index} - P_c) / (1 - P_c)$$

Microsoft Excel was used to calculate the Kappa calculation. Based on the formula above, 0.74 is considered exceptional, 0.60 to 0.74 is acceptable, and 0.54 to 0.59 is fair (23).

### ***Psychometric testing of the pre-final version of the HFSQIQ in a sample of the target population***

This final step was to establish the initial psychometric features of the newly designed questionnaire using a sample of the population of interest. The purpose is to examine the internal structure of the questionnaire; the current study conducts a reliability test that is represented by a high value of the internal consistency and reliability coefficient, often determined by Cronbach's alpha coefficient ( $\alpha$ ) (27). In the present study, the formula presented by Bonett was used (31). The minimal sample size required to determine at least 80.0% power of the test is 22 hospitals based on an alpha value of 0.05. A minimum sample size estimation of 27 respondents from 27 hospitals was employed to determine the internal consistency by conducting a reliability test.

This pilot study’s target population was selected among management representatives from the food service departments of government, private, and teaching hospitals in Malaysia, including the head of the department, catering officer/assistant catering officer, operating manager, and dietitian/catering dietitian. The respondents were chosen based on the researcher’s inclusion and exclusion criteria. They must be a Malaysian citizen, hold a position on the administrative team of the dietetics and food service department, have at least six months of experience working in a hospital food service department, and be proficient in Malay and/or English. To facilitate data collection, the questionnaire was distributed via an online Google Form. A URL link was sent to each respondent by email or WhatsApp. Respondents were requested to evaluate the indicators based on their perceptions of the importance and performance of food service operations.

**Data analysis**

Reliability analysis was performed using the Statistical Package for Social Sciences (SPSS) version 26.0. Cronbach’s alpha ( $\alpha$ ) was used to examine the internal consistency of the HFSQIQ for two subscales measuring the importance and performance of food service indicators. Additionally, Cronbach’s alpha was calculated for the whole HFSQIQ. Internal consistency is acceptable if Cronbach’s alpha value

exceeds 0.70 (29). Furthermore, the corrected item-total correlation and Cronbach’s alpha if an item is deleted were examined for testing the reliability of the newly created HFSQIQ. A good correlation between the item and the total excluding the item is indicated by a corrected item-total correlation ideal value greater than 0.5 and the minimum acceptable value is should not less than 0.30 (32).

**Results**

**Preliminary HFSQIQ questionnaire design**

The literature review identified a few models or methodologies designed to evaluate the performance of hospital food service operations. Additionally, a qualitative content analysis using semi-structured in-depth interviews with 24 staff from the dietetics and food service departments at two hospitals was undertaken to identify the indicators within six domains, including the food service operational management, food production and distribution management, staff management, nutritional management, and patient/customer service management. The domains of each of these indicators were determined conceptually by combining qualitative research with a literature review. The summary of the development HFSQIQ questionnaire is illustrated in Table 1.

**Table 1:** Summary of different version of HFSQIQ from Stage 1 and 2 questionnaire development

Domains	Version 1.0 Prior to the expert evaluation as result of Stage 1, steps 1 & 2.		Version 2.0 after expert evaluation as a result of Stage 2, step 3.		Version 3.0 after evaluation by target population in Phase 2, step 4.	
	Domains	Number of items	Domains	Number of items	Domains	Number of items
Food service operational management	Specification of contract	2	Specification of contract	2	Specification of contract	2
	Diet ordering system	2	Diet ordering system	2	Diet ordering system	2
	Financial/ budget allocation	2	Financial/ budget allocation	2	Financial/ budget allocation	2
	Purchasing and receiving management	4	Purchasing and receiving management	4	Purchasing and receiving management	3
Food production and distribution management	Cooking and ingredients management	5	Cooking and ingredients management	4	Cooking and ingredient management	4
	Hygiene management	4	Hygiene management	2	Hygiene management	3
	Menu planning	5	Menu planning	3	Menu planning	3
	Quality management	2	Quality management	3	Quality management	3
	Distribution management	3	Distribution management	4	Distribution management	4

**Table 1:** Summary of different version of HFSQIQ from Stage 1 and 2 questionnaire development (continued)

Domains	Version 1.0 Prior to the expert evaluation as result of Stage 1, steps 1 & 2.		Version 2.0 after expert evaluation as a result of Stage 2, step 3.		Version 3.0 after evaluation by target population in Phase 2, step 4.	
	Domains	Number of items	Domains	Number of items	Domains	Number of items
Equipment and facility management	Facility management	7	Facility management	6	Facility management	6
	Equipment and facility maintenance	3	Equipment and facility maintenance	2	Equipment and facility maintenance	2
	Upgrade equipment	2	Upgrade equipment	2	Upgrade equipment	2
Staff management	Human resource	4	Human resource	4	Human resource	4
	Job performance	3	Job performance	3	Job performance	2
Nutritional management	Therapeutic diet management	3	Therapeutic diet management	3		
	Nutritional education	4	Nutritional education	4		
Patient/ customer service management	Patient/ customer satisfaction	4	Patient/ customer service	5	Patient/ customer service	5
	Mealtime service	2	Mealtime service	2	Mealtime service	4
<b>Total items</b>		<b>61</b>		<b>57</b>		<b>51</b>

**Expert panel judgement of HFSQIQ’s validity**

The CVI scores’ average and universal agreement is displayed in Table 2. The overall CVI scale (0.88) was evaluated for good Validity, while the S-CVI/Ave ranged from 0.79 and 0.93. The findings revealed that all the items had S-CVI/UA scores less than 0.80 (ranging from 0.17 to 0.50). The I-CVI ratings for 53 items (86.9%) were greater

than or equal to 0.80, and the κ index was greater than or equal to 0.74. In contrast, 8 items (13.1%) had I-CVI ratings below 0.80 and κ index below 0.74. Six items (9.8%) had ratings of I-CVI of 0.71 and κ index of 0.17, one item (1.6%) had I-CVI ratings of 0.57 and κ index of 1.05, and one item (1.6%) had I-CVI ratings of 0.29 and κ index of 1.00.

**Table 2:** The I-CVI and modified kappa index for items for first version of HFSQIQ

Items	Relevant (rating 3 or 4) (n)	Not relevant (rating 1 or 2) (n)	I-CVIs <sup>a</sup>	Pc <sup>b</sup>	κ <sup>c</sup>	Evaluation <sup>d</sup>	Interpretation
<b>Food service operational management</b>							
Item 1	6	1	0.86	0.005	0.85	Excellent	Appropriate
Item 2	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 3	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 4	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 5	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 6	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 7	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 8	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 9	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 10	6	1	0.86	0.055	0.85	Excellent	Eliminate

**Table 2:** The I-CVI and modified kappa index for items for first version of HFSQIQ (continued)

Items	Relevant (rating 3 or 4) (n)	Not relevant (rating 1 or 2) (n)	I-CVIs <sup>a</sup>	Pc <sup>b</sup>	κ <sup>c</sup>	Evaluation <sup>d</sup>	Interpretation
		<b>S-CVI/Ave</b>	0.93				
		<b>S-CVI/UA</b>	0.50				
<b>Food production and distribution management</b>							
Item 11	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 12	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 13	4	3	0.57	9.844	1.05	Fair	Eliminate
Item 14	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 15	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 16	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 17	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 18	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 19	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 20	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 21	5	2	0.71	0.656	0.17	Fair	Eliminate
Item 22	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 23	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 24	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 25	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 26	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 27	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 28	5	2	0.71	0.656	0.17	Fair	Eliminate
Item 29	7	0	1.00	0.008	1.00	Excellent	Appropriate
		<b>S-CVI/Ave</b>	0.83				
		<b>S-CVI/UA</b>	0.26				
<b>Patient/customer service management</b>							
Item 30	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 31	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 32	2	5	0.29	2362.500	1.00	Poor	Eliminate
Item 33	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 34	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 35	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 36	7	0	1.00	0.008	1.00	Excellent	Appropriate
		<b>S-CVI/Ave</b>	0.79				
		<b>S-CVI/UA</b>	0.17				
<b>Equipment and facility management</b>							
Item 37	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 38	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 39	5	2	0.71	0.656	0.17	Fair	Eliminate
Item 40	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 41	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 42	7	0	1.00	0.008	1.00	Excellent	Appropriate

**Table 2:** The I-CVI and modified kappa index for items for first version of HFSQIQ (continued)

Items	Relevant (rating 3 or 4) (n)	Not relevant (rating 1 or 2) (n)	I-CVIs <sup>a</sup>	Pc <sup>b</sup>	κ <sup>c</sup>	Evaluation <sup>d</sup>	Interpretation
Item 43	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 44	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 45	5	2	0.71	0.656	0.17	Fair	Eliminate
Item 46	5	2	0.71	0.656	0.17	Fair	Need for revision
Item 47	5	2	0.71	0.656	0.17	Fair	Eliminate
		<b>S-CVI/Ave</b>	0.88				
		<b>S-CVI/UA</b>	0.50				
<b>Staff management</b>							
Item 48	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 49	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 50	7	0	1.00	0.008	1.00	Excellent	Appropriate
Item 51	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 52	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 53	6	1	0.86	0.055	0.85	Excellent	Appropriate
Item 54	6	1	0.86	0.055	0.85	Excellent	Appropriate
		<b>S-CVI/Ave</b>	0.92				
		<b>S-CVI/UA</b>	0.43				
<b>Nutritional management</b>							
Item 55	6	1	0.86	0.055	0.85	Excellent	Eliminate
Item 56	7	0	1.00	0.008	1.00	Excellent	Eliminate
Item 57	7	0	1.00	0.008	1.00	Excellent	Eliminate
Item 58	6	1	0.86	0.055	0.85	Excellent	Need for revision
Item 59	6	1	0.86	0.055	0.85	Excellent	Need for revision
Item 60	6	1	0.86	0.055	0.85	Excellent	Eliminate
Item 61	7	0	1.00	0.008	1.00	Excellent	Eliminate
		<b>S-CVI/Ave</b>	0.92				
		<b>S-CVI/UA</b>	0.43				

<sup>a</sup>Overall-scale CVI = 0.88

<sup>b</sup>Pc (probability of chance occurrence) =  $[(N!A!)(N - A)!] * 0.5N$ , where N= number of experts, and A=number of experts who agreeing on a rating of 3 and 4.

<sup>c</sup>κ\* =  $(I-CVI - Pc) / (1 - Pc)$

<sup>d</sup>Evaluation criteria for level of content validity: relationship between I-CVI and κ\*: excellent validity =  $0.80 > κ > 0.74$ ; good validity =  $0.60 ≤ κ ≤ 0.74$ ; fair validity =  $0.40 ≤ κ ≤ 0.59$ .

The S-FVI/Ave ranged from 0.99 to 1.00 for the pre-final HFSQIQ English version, whereas the S-FVI/UA ranged from 0.89 to 1.00. Meanwhile, S-FVI/Ave varied from 0.93 to 1.00 and S-FVI/UA ranged from 0.79 to 1.00 for the Malay version of the pre-final HFSQIQ. For the pre-final HFSQIQ English version, the I-FVI ratings for all items were evaluated as excellent for face validity, with values greater than 0.80 (ranging from 0.90 to 1.00). While the Malay version was rated at or above 0.80 and κ index greater than 0.74 (ranging from 0.90 to 1.00) (Table 3). Based on

the findings, the pre-final HFSQIQ for both English and Malay versions was appropriate, as each received excellent validation ratings for face validity.

Five males (18.5%) and twenty-two females (81.5%), representing 27 hospitals, signed the online consent form and completed the questionnaire for the reliability study. The respondents' mean average age ( $±$  standard deviation) was 34.74 ( $±$  7.128) years. Most respondents were dietitians or catering dietitians (59.3%), followed by

**Table 3:** FVI of item understandability and modified kappa agreement index for English and Malay version of pre-final HFSQIQ (N=10)

Items No	English version of HFSQIQ					Malay version of HFSQIQ				
	Understand (rating 3 or 4)	I-FVIs*	Pc <sup>a</sup>	κ <sup>b</sup>	Interpretation <sup>c</sup>	Understand (rating 3 or 4)	I-FVIs**	Pc <sup>a</sup>	κ <sup>b</sup>	Interpretation <sup>c</sup>
<b>Food service operational management</b>										
Item 1	9	0.90	0.010	0.90	Excellent	10	1.00	0.001	1.00	Excellent
Item 2	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 3	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 4	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 5	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 6	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 7	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 8	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 9	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
	<b>S-FVI/Ave</b>	<b>0.99</b>				<b>S-FVI/Ave</b>	<b>1.00</b>			
	<b>S-FVI/UA</b>	<b>0.89</b>				<b>S-FVI/UA</b>	<b>1.00</b>			
<b>Food production and distribution management</b>										
Item 10	10	1.00	0.001	1.00	Excellent	9	0.90	0.010	0.90	Excellent
Item 11	9	0.90	0.010	0.90	Excellent	10	1.00	0.001	1.00	Excellent
Item 12	10	1.00	0.001	1.00	Excellent	9	0.90	0.010	0.90	Excellent
Item 13	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 14	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 15	10	1.00	0.001	1.00	Excellent	9	0.90	0.010	0.90	Excellent
Item 16	9	0.90	0.010	0.90	Excellent	10	1.00	0.001	1.00	Excellent
Item 17	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 18	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 19	10	1.00	0.001	1.00	Excellent	9	0.90	0.010	0.90	Excellent
Item 20	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 21	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 22	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 23	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 24	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 25	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 26	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 27	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
	<b>S-FVI/Ave</b>	<b>0.99</b>				<b>S-FVI/Ave</b>	<b>0.93</b>			
	<b>S-FVI/UA</b>	<b>0.89</b>				<b>S-FVI/UA</b>	<b>0.79</b>			
<b>Patient/customer service management</b>										
Item 28	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 29	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 30	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 31	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 32	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 33	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 34	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 35	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 36	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
	<b>S-FVI/Ave</b>	<b>1.00</b>				<b>S-FVI/Ave</b>	<b>1.00</b>			
	<b>S-FVI/UA</b>	<b>1.00</b>				<b>S-FVI/UA</b>	<b>1.00</b>			
<b>Equipment and facility management</b>										
Item 37	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 38	9	0.90	0.010	0.90	Excellent	10	1.00	0.001	1.00	Excellent



**Table 3:** FVI of item understandability and modified kappa agreement index for English and Malay version of pre-final HFSQIQ (N=10) (continued)

Items No	English version of HFSQIQ					Malay version of HFSQIQ				
	Understand (rating 3 or 4)	I-FVIs*	Pc <sup>a</sup>	κ <sup>b</sup>	Interpretation <sup>c</sup>	Understand (rating 3 or 4)	I-FVIs**	Pc <sup>a</sup>	κ <sup>b</sup>	Interpretation <sup>c</sup>
Item 39	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 40	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 41	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 42	10	1.00	0.001	1.00	Excellent	9	0.90	0.010	0.90	Excellent
Item 43	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 44	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 45	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 46	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
	<b>S-FVI/Ave</b>	<b>0.99</b>				<b>S-FVI/Ave</b>	<b>0.99</b>			
	<b>S-FVI/UA</b>	<b>0.89</b>				<b>S-FVI/UA</b>	<b>0.89</b>			
<b>Staff management</b>										
Item 47	10	1.00	0.001	1.00	Excellent	9	0.90	0.010	0.90	Excellent
Item 48	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 49	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 50	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
Item 51	10	1.00	0.001	1.00	Excellent	10	1.00	0.001	1.00	Excellent
	<b>S-FVI/Ave</b>	<b>1.00</b>				<b>S-FVI/Ave</b>	<b>0.98</b>			
	<b>S-FVI/UA</b>	<b>1.00</b>				<b>S-FVI/UA</b>	<b>0.83</b>			

\*Face validity index average for 51 items in English version of HFSQIQ = 0.99

\*\*Face validity index average for 51 items in Malay version of HFSQIQ = 0.99

<sup>a</sup>Pc (probability of chance occurrence) = [(N!A!)(N - A)!] \* 0.5N, where N = number of experts, and A=number of experts who agreeing on a rating of 3 and 4.

<sup>b</sup>κ = (I-CVI-Pc)(1-Pc)

<sup>c</sup>Evaluation criteria for level of face validity: Excellent = κ > 0.7, Good = 0.6 ≤ κ ≤ 0.74, and Fair = 0.40 ≤ κ ≤ 0.59

directors or heads of departments (25.9%), catering officers or assistant catering officers (11.1%), and only one manager or catering manager (3.7%). The mean average length of service was 9.219 (± 7.702) years. The majority of them have a bachelor’s degree (85.2%), three have a master’s or PhD (11.1%), and only one has a diploma (3.6%). Table 4 provides a summary of the respondent characteristics.

**Table 4:** The characteristics of the respondents for reliability study (N = 27)

Characteristics	n (%) / mean (±SD)
Age (years), mean (±SD)	34.74 (±7.128)
Sex	
Male	5 (18.5)
Female	22 (81.5)
Job position	
Director/ Operation director/ Head of department	7 (25.9)
Manager/ Catering manager/ Outlet manager	1 (3.7)

**Table 4:** The characteristics of the respondents for reliability study (N = 27) (continued)

Characteristics	n (%) / mean (±SD)
Dietitian/ Catering dietitian	16 (59.3)
Catering officer/ Assistant catering officer	3 (11.1)
Length of service, mean (±SD)	9.219 (±7.706)
Academic qualification	
Diploma	1 (3.7)
Degree	23 (85.2)
Master/ PhD	3 (11.1)

Scoring scale analysis was also performed for the final HFSQIQ by assessing the internal consistency and reliability of the importance and performance scales as depicted in Tables 5 and 6, respectively. The Cronbach alpha (α) values for the overall importance and performance scales were 0.97 and 0.98, respectively. On the importance scale, the subscale scores for food service operational management,

food production and distribution management, patient/customer service management, equipment and facility management, and staff management were 0.809, 0.937, 0.900, 0.919, and 0.939, respectively.

**Table 5:** The internal consistency of the item total statistics for importance scale

Item	Scale means if item deleted	Scale variance if item deleted	Corrected item total correlation	Cronbach alpha if item deleted
<b>Food service operational management</b>				
Item 1	37.63	6.934	0.459	.796
Item 2	37.59	7.020	0.557	.785
Item 3	37.96	7.268	0.180	.853
Item 4	37.44	7.564	0.495	.796
Item 5	37.55	7.103	0.556	.786
Item 6	37.67	6.712	0.569	.782
Item 7	37.67	6.538	0.597	.778
Item 8	37.67	5.923	0.855	.740
Item 9	37.56	7.103	0.556	.786
			<b>Cronbach alpha</b>	<b>.809</b>
<b>Food production and distribution management</b>				
Item 10	79.11	41.179	0.658	.933
Item 11	79.26	40.123	0.632	.935
Item 12	79.70	44.678	0.523	.936
Item 13	78.96	42.575	0.654	.933
Item 14	78.89	42.641	0.577	.935
Item 15	78.78	44.103	0.525	.936
Item 16	78.74	44.430	0.509	.936
Item 17	78.96	41.268	0.749	.931
Item 18	78.67	44.769	0.610	.936
Item 19	79.15	39.977	0.819	.929
Item 20	79.07	39.994	0.813	.929
Item 21	79.00	39.692	0.864	.928
Item 22	79.04	41.422	0.708	.932
Item 23	78.85	42.362	0.767	.932
Item 24	78.89	42.362	0.767	.932
Item 25	79.11	43.179	0.498	.936
Item 26	79.04	41.256	0.586	.935
Item 27	79.04	41.729	0.665	.933
			<b>Cronbach alpha</b>	<b>.937</b>
<b>Patient/ customer service management</b>				
Item 28	36.04	12.575	0.629	.891
Item 29	35.96	12.422	0.680	.888
Item 30	36.26	10.892	0.789	.879
Item 31	36.11	11.256	0.723	.885
Item 32	36.04	11.114	0.835	.874
Item 33	35.82	13.157	0.513	.899

**Table 5:** The internal consistency of the item total statistics for importance scale (continued)

Item	Scale means if item deleted	Scale variance if item deleted	Corrected item total correlation	Cronbach alpha if item deleted
Item 34	35.85	12.900	0.572	.895
Item 35	35.96	12.191	0.640	.890
Item 36	36.11	12.256	0.629	.891
			<b>Cronbach alpha</b>	<b>.900</b>
<b>Equipment and facility management</b>				
Item 37	38.26	7.046	0.870	.898
Item 38	38.15	7.593	0.794	.906
Item 39	38.07	8.302	0.608	.918
Item 40	38.30	6.755	0.690	.917
Item 41	38.19	7.695	0.663	.913
Item 42	38.15	7.746	0.710	.910
Item 43	38.30	7.524	0.614	.917
Item 44	38.30	7.046	0.870	.898
Item 45	38.33	7.154	0.748	.907
			<b>Cronbach alpha</b>	<b>.919</b>
<b>Staff management</b>				
Item 46	23.04	5.422	0.795	.931
Item 47	22.96	5.114	0.812	.930
Item 48	22.85	5.746	0.690	.943
Item 49	22.93	5.225	0.914	.917
Item 50	22.93	5.302	0.874	.922
Item 51	22.89	5.103	0.841	.925
			<b>Cronbach alpha</b>	<b>.939</b>

\*Overall Cronbach alpha ( $\alpha$ ) = .970

On the performance scale, the subscale scores for food service operational management, production and distribution management, patient/customer service management, equipment and facility management, and staff management were 0.907, 0.972, 0.933 and 0.914,

respectively. The questionnaire's Cronbach alpha remained consistent with a significant difference if an item was deleted from the importance and importance scales, demonstrating that the newly developed questionnaire has excellent internal reliability.

**Table 6:** The internal consistency of the item total statistics for performance scale

Item	Scale means if item deleted	Scale variance if item deleted	Corrected item total correlation	Cronbach alpha if item deleted
<b>Food service operational management</b>				
Item 1	26.48	28.413	0.735	.895
Item 2	26.33	27.077	0.766	.891
Item 3	27.00	27.462	0.458	.920
Item 4	26.74	27.969	0.473	.915
Item 5	26.41	26.789	0.816	.888
Item 6	26.44	27.179	0.834	.888

**Table 6:** The internal consistency of the item total statistics for performance scale (continued)

Item	Scale means if item deleted	Scale variance if item deleted	Corrected item total correlation	Cronbach alpha if item deleted
Item 7	26.33	26.385	0.804	.888
Item 8	26.33	26.538	0.784	.889
Item 9	26.15	27.516	0.718	.894
			<b>Cronbach alpha</b>	<b>.907</b>
<b>Food production and distribution management</b>				
Item 10	56.96	153.268	0.802	.970
Item 11	56.85	153.285	0.827	.970
Item 12	56.78	150.333	0.919	.969
Item 13	56.74	147.661	0.861	.969
Item 14	56.67	153.077	0.828	.970
Item 15	56.67	149.231	0.872	.969
Item 16	56.70	149.370	0.912	.969
Item 17	56.96	154.575	0.607	.972
Item 18	56.81	149.849	0.845	.969
Item 19	57.00	154.385	0.704	.971
Item 20	56.93	152.148	0.851	.969
Item 21	56.89	151.410	0.741	.971
Item 22	57.04	153.575	0.639	.972
Item 23	56.74	146.969	0.892	.969
Item 24	56.81	147.772	0.860	.969
Item 25	56.89	147.564	0.842	.969
Item 26	57.30	153.293	0.656	.972
Item 27	57.04	150.575	0.778	.970
			<b>Cronbach alpha</b>	<b>.972</b>
<b>Patient/ customer service management</b>				
Item 28	26.56	29.718	0.723	.926
Item 29	26.59	29.635	0.837	.920
Item 30	26.70	29.293	0.849	.919
Item 31	26.67	29.000	0.876	.917
Item 32	26.63	30.396	0.806	.922
Item 33	26.41	30.097	0.727	.926
Item 34	26.41	31.020	0.663	.930
Item 35	27.07	28.764	0.727	.927
Item 36	26.96	31.268	0.580	.935
			<b>Cronbach alpha</b>	<b>.933</b>
<b>Equipment and facility management</b>				
Item 37	25.89	32.718	0.706	.903
Item 38	25.89	36.949	0.260	.933
Item 39	25.89	32.333	0.847	.895
Item 40	25.89	32.333	0.794	.898
Item 41	25.81	31.541	0.919	.890
Item 42	25.70	32.217	0.876	.894

**Table 6:** The internal consistency of the item total statistics for performance scale (continued)

Item	Scale means if item deleted	Scale variance if item deleted	Corrected item total correlation	Cronbach alpha if item deleted
Item 43	26.15	32.054	0.686	.905
Item 44	25.82	31.003	0.832	.894
Item 45	26.15	32.285	0.556	.918
			<b>Cronbach alpha</b>	<b>.914</b>
<b>Staff management</b>				
Item 46	15.63	8.550	0.790	.847
Item 47	15.67	9.154	0.515	.882
Item 48	15.78	8.179	0.642	.865
Item 49	16.22	7.718	0.621	.873
Item 50	15.78	7.256	0.773	.842
Item 51	15.74	7.892	0.867	.830
			<b>Cronbach alpha</b>	<b>.878</b>

\*Overall Cronbach alpha ( $\alpha$ ) = .981

## Discussion

This study aimed to develop a valid and reliable questionnaire for measuring the performance of hospital food service operations in Malaysia. This study described the items' development process, translation, and validation of a newly proposed "Hospital Food Service Quality Improvement Questionnaire" (HFSQIQ). The TQM approach proposed by Balasubramanian for basic measurement and continuous improvement provided a framework for developing the questionnaire domains (15). In addition, a literature search on hospital food service quality improvement served as a guide for developing the questionnaire's items (18-21). Some researchers from prior studies presented the "Importance – performance analysis" (IPA) developed by Martilla and James (33) to quantify quality attributes based on two measurement scales: 1) their value to operations (importance), and 2) their effectiveness of the operations or management (performance) (18, 20, 21). The development of the questionnaire items should start with identifying significant elements of the management of the food service operation from previous research in the same or related areas. Various qualitative research methods, such as focus groups, personal interviews, and managerial discretion, are essential for identifying potentially significant variables that might be overlooked.

This study utilised several approaches, including a literature review, in-depth interviews, peer review, and expert panel judgment, to identify the relevant topics or domains and items for measuring the perceived importance and performance of hospital food service management. The first version of HFSQIQ, which included six domains with 61 items, was generated before the expert evaluation. After expert review and content validation analysis, the number of items in the second version of the HFSQIQ was reduced to 57 while the six domains were maintained.

Item 10 was removed after discussion and agreement with the supervisory team, despite its excellent validity, given that the item was identified as not a crucial component of the measurement scale. Additionally, since staff members other than dietitians cannot evaluate performance in the nutritional management domain and the questionnaire must focus on food service management, the supervisory teams opted to eliminate Items 51, 52, 54, 55, 56, and 57. However, Item 53 was suggested to be added to the patient/customer service management domain for it is pertinent. Finally, after face validity analysis, the third version of the HFSQIQ was revised to five domains and 51 items.

Before the translation process, the first version of the HFSQIQ underwent content validation, which included a detailed evaluation of the items' relevance and clarity, and face validation after the translation process to evaluate the understandability of the questionnaire by the target users or population. The content validity index is easy to use and understandable by providing detailed information on the strengths and weaknesses of each item, leading to the deletion and modification of items for a valid reason (34). Face validity, on the other hand, focused more on the design or structure of the questionnaire and its readability by the targeted users (35). In this study, two different indices were computed for content validation to determine the percentage of items on the S-CVI/UA and S-CVI/Ave by the experts. However, the findings revealed that the average agreement among experts has a high content validity level, although the experts' universal agreement ratings were less than 0.80 for all domains.

Questions were raised regarding the calculation of the agreement indices and the possibility of inaccuracy. Although the agreement indices are just one step in

determining content validity, other factors should be used as well to decide whether to reject or modify items (34). Most scale developers employ the 0.80 criterion set by Davis (36) as the minimum acceptable S-CVI value for new instruments or questionnaires (27). For example, Polit et al. (24) stated that even if the content validity of the scale of the items was insufficient using the S-CVI/UA approach ( $< 0.80$ ), it is still sufficient to employ the S-CVI/Ave approach ( $> 0.80$ ). As an alternative to the content validity index, the modified kappa index was utilised in this study to verify the findings as it considers and incorporates chance agreement. Including the CVI and a multi-rater kappa coefficient in the content validation, as suggested by Wynd et al. (37), is a significant supplement to the CVI since the kappa coefficient offers information regarding the degree of agreement that exceeds chance. Therefore, high content and face validity index scores indicate that the HFSQIQ was established appropriately and is reliable for hospital food service operations in Malaysia.

The translation, adaptation, and validation method used in this study aligned with a thorough and detailed set of guidelines developed by Sousa and Rojjanasrirat (28) to maintain the items' original impact, style, and meaning when they are translated from English into Malay. Since these terminologies are more commonly used in English than in Malay, more efforts were required to translate some technical words. For example, a direct translation for item 30, "providing various food choices for a patient with normal diet", from English to Malay, "*menyediakan pelbagai makanan untuk pesakit diet normal*" was accurate. However, based on review by the certified translator, the translation was harmonised to "*menyediakan pelbagai pilihan makanan untuk pesakit dengan diet normal*". Hence, the participation of professional translators proficient in both the target and source languages is required to ensure that the respondents appropriately translated and understood the items.

One of the important components of test quality is reliability. It involves either an examinee's performance on the test items or the consistency or reproducibility of the results. Reliability is the consistent results of a given measurement. When a measurement is considered reliable if it consistently produces the same results under the same conditions (32). Internal consistency reliability was used in this study to evaluate the consistency of results across the HFSQIQ items. Cronbach's alpha is the most common internal consistency statistic used to identify the relationship between all test items (32). The findings indicate that the HFSQIQ was evaluated among 27 respondents. Cronbach's alpha for the five importance and performance measurement scales subscales ranged from 0.809 to 0.973. This result demonstrated that the newly developed HFSQIQ could rely upon to evaluate the performance of hospital food service operations in Malaysia.

Several limitations were identified during data collection throughout this study. It was not feasible to collect face-to-face data in hospitals due to the COVID-19 pandemic, thus a web-based data collection method was applied in this study. The responses were obtained through phone interviews or virtual meetings using the Google Meet platform. The questionnaires were distributed using the Google Forms application. Although virtual meetings have a few advantages, some technical concerns, such as a sudden internet disruption or slowdown, may have resulted in communication problems. To solve this problem, separate virtual meetings were held with the respondents who experienced issues with their phone or internet connections. Another limitation is that the items derived from research conducted in other countries may not be relevant to Malaysian settings. Thus, the Delphi technique study may be recommended for future research in developing tools to evaluate hospital food service performance. This could be achieved by soliciting the opinions of experts to identify a consensus position and present findings on a specific topic or set of questions based on the knowledge and experience of experts in the field (38). Finally, despite internal consistency reliability being the most basic test used for newly developed instruments, recommending additional reliability measures of re-test reliability would be beneficial for determining the consistency of a set of parameters.

### **Conclusion**

The results of this study demonstrate the high validity and reliability of a newly developed questionnaire entitled the HFSQIQ. HFSQIQ is a tool that can evaluate the importance and performance of hospital food service aspects or components for quality enhancement. Future research should include a criterion validation study, as this would be able to predict the outcome of another measure or domain of the HFSQIQ. This questionnaire can also be used in other businesses or industries that provide food services because it is a straightforward and practical tool for identifying food service-related components for continuous quality improvement of food service operations.

### **Acknowledgement**

We would like to thank Assoc. Prof. Dr. Mohd Shazali Md Sharif, Dr. Syahrul Bariah Abdul Hamid, and Dr. Syafiqah Rahamat for their contribution to the supervisory team review during the validation process. We would like to acknowledge Datin Musalmah Buzri, Mrs. Khor Jo Ann, Dr. Syakirah M Hussein, and Mr. Muhammad Zaki Ramli, for their expert comments in improving the translation process. We also would like to thank those who participated in this study.

### **Competing interests**

The authors declare that they have no competing interests.

### Ethical clearance

The study adhered to the Declaration of Helsinki and was approved by the UiTM Research Ethics Committee (reference number 600-IRMI (5/1/6)) and the Medical Research and Ethics Committee, Ministry of Health, Malaysia (NMRR-19-3262-47594 (IIR)).

### Financial support

This research was financially supported by Universiti Teknologi MARA through Research Incentive Grant (600-RMC/GIP 5/3 (015/2021). The funders were not involved in the study's design, data collection, analysis, interpretation, manuscript writing, or the decision to publish the findings.

### References

- Gonçalves JM, Rodrigues KL, Almeida ÂTS, Pereira GM, Buchweitz MRD. Assessment of good practices in hospital food service by comparing evaluation tools. *Nutricion Hospitalaria*. 2015; 32(4):1796–1801.
- Dall'Oglio I, Nicolò R, Di Ciommo V, Bianchi N, Ciliento G, Gawronski O, *et al*. A systematic review of hospital foodservice patient satisfaction studies. *Journal of the Academy of Nutrition and Dietetics*. 2015; 115(4): 567–584.
- Zahran SE, Bakr ESH. (2021). Suitability of patient's meals for patients' diagnosis and food quality services evaluation in hospitals. *Bioscience Research*. 2021; 18(1):773–781.
- Barker LA, Gout BS, Crowe TC. Hospital malnutrition: prevalence, identification and impact on patients and the healthcare system. *International Journal of Environmental Research and Public Health*. 2011; 8(2):514–527.
- Lai H, Gemming L. Approaches to patient satisfaction measurement of the healthcare food services: A systematic review. *Clinical Nutrition ESPEN*. 2021; 42:61–72.
- Banks M, Hannan-Jones M, Ross L, Buckley A, Ellick J, Young A. Measuring the quality of hospital food services: Development and reliability of a Meal Quality Audit Tool. *Nutrition and Dietetics*. 2017; 74(2):147–157.
- Huang JS, Chun S, Cheung C, Poon L, Terrones L. The nutritional value of food service meals ordered by hospitalized children. *Clinical Nutrition ESPEN*. 2016; 15:122–125.
- Ahmed M, Jones E, Redmond E, Hewedi M, Wingert A, El Rab MG. Food production and service in UK hospitals. *International Journal of Health Care Quality Assurance*. 2015; 28(1):40–54.
- Aminuddin NF, Vijayakumaran RK, Razak SA. Patient Satisfaction with hospital foodservice and its impact on plate waste in public hospitals in East Malaysia. *Hospital Practices and Research*. 2018; 3(3):90–97.
- Batra M, Taneja U. The role of services cape on emotions and satisfaction of patients in Indian Hospitals. *Journal of Health Management*. 2022; 24(3):433–439.
- Messina G, Fenucci R, Vencia F, Niccolini F, Quercioli C, Nante N. Patients' evaluation of hospital foodservice quality in Italy: what do patients really value? *Public Health Nutrition*. 2013; 16(4):730–737.
- Tamby Chik C, Zulkipli NA, Bachok S, Mohi Z, Mohd Shahril A. Plate Waste in Public Hospitals Foodservice Management in Selangor, Malaysia. *Malaysia Article in Indian Journal of Science and Technology*. 2019; 11(36): 1–5.
- Vijayakumaran RK, Eves A, Lumbers M. Patients Emotions during Meal Experience: Understanding through Critical Incident Technique. *International Journal of Hospital Research*. 2016; 5(4):113–121.
- Vijayakumaran RK, Eves A, Lumbers M. Understanding Patients' Meal Experiences through Staff's Role: Study on Malaysian Public Hospitals. *Hospital Practices and Research*. 2018; 3(2):50–58.
- Balasubramanian M. Total Quality Management [TQM] in the Healthcare Industry – Challenges, Barriers and Implementation Developing a Framework for TQM Implementation in a Healthcare Setup. *Science Journal of Public Health*. 2016;4(4): 271.
- Rapo S, Sydner YM, Kautto E, Hörnell A. Exploring patient satisfaction with hospital foodservice: A Swedish study using the Acute Care Hospital Foodservice Patient Satisfaction Questionnaire. *Nutrition and Dietetics*. 2021; 78(5):487–495.
- Jamaluddin R, Abd Manan, NA, Basri AM, Ab Karim MS. Patients' satisfaction with the bulk trolley system in a government hospital in Malaysia. *Leadership in Health Services*. 2010; 23(3):260–268.
- Lee JE. Importance-performance analysis of evaluation indicators in Hospital Nutrition Department. *J Korean Diet Assoc*. 2012; 18(4):326–343.
- Diez-Garcia RW, Japur CC, Tavares de Medeiros MA. Food and nutritional care quality indicators in hospital. *Journal of Hospital Administration*. 2013; 2(3): 132–141.
- Lee JE. Importance-performance analysis on foodservice management items of the dieticians at long-term care hospitals. *Indian Journal of Science and Technology*. 2016; 9(26):1–7.
- Park MS, Lyu ES. Importance and performance of dietitian's Task at long term care hospital foodservice in Busan · Kyungnam Area. *Korean Journal of Community Nutrition*. 2011; 16(5):602–612.
- Osman NS, Md Nor N, Md Sharif MS, Rahamat S, Abdul Hamid SB. Barriers and challenges of an outsource hospital foodservice operation: A narrative thematic analysis. *Malaysian Journal of Medicine and Health Sciences*. 2022; 18(SUPP15): 97-107.
- Yusoff, MSB. ABC of Content Validation and Content Validity Index Calculation. *Education in Medicine Journal*. 2019; 11(3), 49–54.
- Polit DF, Beck CT, Owen SV. Focus on research methods: Is the CVI an acceptable indicator of content

- validity? Appraisal and recommendations. *Research in Nursing and Health*. 2007; 30(4):459–467.
25. Polit, DF, Beck, CT. (2006). The content validity index: Are you sure you know what's being reported? Critique and recommendations. *Research in Nursing and Health*. 2006; 29(5), 489–497.
  26. Lynn, MR. Determination and quantification of content validity. *Nursing Research*. 1986; 35, 382–385.
  27. Yusoff MSB, Arifin WN, Hadie SNH. ABC of questionnaire development and validation for survey research. *Education in Medicine Journal*. 2021; 13(1):97–108.
  28. Sousa VD, Rojjanasrirat W. Translation, adaptation and validation of instruments or scales for use in cross-cultural health care research: A clear and user-friendly guideline. In *Journal of Evaluation in Clinical Practice*. 2011; 17(2):268–274.
  29. Yusoff MSB. ABC of response process validation and face validity index calculation. *Education in Medicine Journal*. 2019; 11(3):55–61.
  30. Marzuki MFM, Yaacob NA, Yaacob NM. Translation, cross-cultural adaptation, and avalidation of the Malay version of the System Usability Scale Questionnaire for the assessment of mobile apps. *JMIR Human Factors*. 2018; 5(2). <http://humanfactors.jmir.org/2018/2/e10308/>
  31. Bonett DG. Sample size requirements for testing and estimating coefficient alpha. *Journal of Educational and Behavioral Statistics*. 2002; 27(4):335–340.
  32. El Hajjar ST. Statistical analysis: internal-consistency reliability and construct Validity. *International Journal of Quantitative and Qualitative Research Methods*. 2018; 6(1):46–57. [www.eajournals.org](http://www.eajournals.org)
  33. Martilla JA, James JC. Importance-performance analysis. *Journal of Marketing*. 1977; 41(1):77–79.
  34. Halek M, Holle D, Bartholomeyczik S. Development and evaluation of the content validity, practicability and feasibility of the innovative dementia-oriented assessment system for challenging behaviour in residents with dementia. *BMC Health Services Research*. 2017; 17(554): 1–26.
  35. Zamanzadeh V, Ghahramanian A, Rassouli M, Abbaszadeh A, Alavi-Majd H, Nikanfar AR. Design and implementation content validity study: Development of an instrument for measuring Patient-Centered Communication. *Journal of CarignSceicnes*. 2015; 4(2):165 – 178.
  36. Davis LL. Instrument review: Getting the most from a panel of experts. *Applied Nursing Research*. 1992; 5:194–197.
  37. Wynd CA, Schaefer MA. The Osteoporosis Risk Assessment Tool: establisging content validity through a panel of Experts. *Applied Nursing Research*. 2002; 16(2):84–188.
  38. Barrett D, Heale R. What are Delphi studies? *Evid Nased Nurs*. 2020; 23(3):68–69.