

Manuscript Details

Manuscript number	APPETITE_2019_330_R1
Title	Exploration of associations between the FTO rs9939609 genotype, fasting and postprandial appetite-related hormones and perceived appetite in healthy men and women
Article type	Full Length Article

Abstract

Background: The fat mass and obesity-associated gene (FTO) rs9939609 A-allele has been associated with obesity risk. Although the exact mechanisms involved remain unknown, the FTO rs9939609 A-allele has been associated with an impaired postprandial suppression of appetite. Objectives: To explore the influence of FTO rs9939609 genotype on fasting and postprandial appetite-related hormones and perceived appetite in a heterogeneous sample of men and women. Design: 112 healthy men and women aged 18-50-years-old completed three laboratory visits for the assessment of FTO rs9939609 genotype, body composition, aerobic fitness, resting metabolic rate, visceral adipose tissue, liver fat, fasting leptin, and fasting and postprandial acylated ghrelin, total PYY, insulin, glucose and perceived appetite. Participants wore accelerometers for seven consecutive days for the assessment of physical activity and sedentary behaviour. Multivariable general linear models quantified differences between FTO rs9939609 groups for fasting and postprandial appetite outcomes, with and without the addition of a priori selected physiological and behavioural covariates. Sex-specific univariable Pearson's correlation coefficients were quantified between the appetite-related outcomes and individual characteristics. Results: 95% confidence intervals for mean differences between FTO rs9939609 groups overlapped zero in unadjusted and adjusted general linear models for all fasting ($P \geq 0.28$) and postprandial ($P \geq 0.19$) appetite-related outcomes. Eta² values for explained variance attributable to FTO rs9939609 were $< 5\%$ for all outcomes. An exploratory correlation matrix indicated that associations between fasting and postprandial acylated ghrelin, total PYY and general or abdominal adiposity were also small ($r = -0.23$ to 0.15 , $P \geq 0.09$). Fasting leptin, glucose and insulin and postprandial insulin concentrations were associated with adiposity outcomes ($r = 0.29$ to 0.81 , $P \leq 0.033$). Conclusions: Associations between the FTO rs9939609 genotype and fasting or postprandial appetite-related outcomes were weak in healthy men and women.

Keywords	FTO; appetite; ghrelin; PYY; hunger.
Taxonomy	Sex-based Differences on Appetite, Appetite Assessment
Manuscript category	Physiology and Metabolism
Corresponding Author	David Stensel
Corresponding Author's Institution	Loughborough University
Order of Authors	Fernanda R. Goltz, Alice Thackray, veronica varela mato, James King, James Dorling, Monika Dowejko, Sarabjit Mastana, Julie Thompson, Greg Atkinson, David Stensel
Suggested reviewers	Miriam Glegg, Andy Blannin, James Betts

Submission Files Included in this PDF

File Name [File Type]

Response to reviewers.docx [Response to Reviewers]

Abstract.docx [Abstract]

Revised manuscript.docx [Manuscript File]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given:
Data will be made available on request

Dear Dr. Appelhans,

RE: APPETITE_2019_330

18/05/2019

We would like to thank the reviewers for giving their time to carefully examine our manuscript. Our research team are delighted to be given the opportunity to revise our manuscript for additional consideration by *Appetite*. Please find below a list of point-by-point responses to the comments raised by the reviewers. For clarity, changes to the manuscript have been highlighted in yellow. We hope that we have interpreted these comments accurately and that our responses and manuscript modifications are satisfactory.

We look forward to hearing about our paper in due course.

Yours sincerely,



Professor David Stensel
Professor of Exercise Metabolism
School of Sport, Exercise and Health Sciences
Loughborough University
Leicestershire
LE11 3TU, UK
Phone: +44(0)1509 226344
Email: D.J.Stensel@lboro.ac.uk

Reviewer one:

Comment #1: The manuscript aims “to explore the influence of the FTO genotype on fasting and postprandial appetite-related hormones and perceived appetite in heterogeneous sample of men and women”. The study is innovative and current, but presents some important problems that should be reviewed.

Author response #1: We thank the reviewer for the kind comment on the novelty of our study and we hope our responses below and the modifications in the manuscript address the comments raised.

Comment #2: Line 2 - the authors need to put the "rs" of the FTO gene that was studied, considering that there are several "rs" in the scientific literature. Do not just put "risk AA genotype".

Author response #2: We have specified the “rs” of the FTO gene throughout the manuscript, including the title and the abstract.

Comment #3: Line 4 - The authors said that the study population was heterogeneous, but they were all adults. Therefore, the age difference of the research volunteers should be expected when it is proposed to evaluate adults without limiting the age group.

Author response #3: We recruited men and women aged between 18 and 50 years old. We have included this information in the abstract for clarity as follows:

Abstract, page 2, lines 32-35: 112 healthy men and women aged 18-50 years old completed three laboratory visits for the assessment of FTO rs9939609 genotype, body composition, aerobic fitness, resting metabolic rate, visceral adipose tissue, liver fat, fasting leptin, and fasting and postprandial acylated ghrelin, total PYY, insulin, glucose and perceived appetite.

Comment #4: Introduction: Paragraph 4 - the authors described ghrelin, adipose tissue, physical activity and specifically spoke of ghrelin in obese people. Were the other variables described in the paragraph observed in eutrophic or obese people? The behavior of several indicators described differ between eutrophic and obese. In addition, the study evaluated eutrophic.

Author response #4: Our study included participants with a wide range of adiposity, from normal weight to obesity (BMI range from 18.4 to 40.3 kg·m⁻², as described in Table 1). The wide range of adiposity enabled us to evaluate whether adiposity was associated with the appetite-related outcomes of interest. The evidence highlighted in the fourth paragraph of the introduction is an overview of potential factors that can influence appetite. We have specified for each study cited whether the study sample included individuals with normal weight, overweight or obesity, as follows:

Introduction, page 4, lines 90-106: Data from previous studies have indicated that women exhibit higher fasting concentrations of acylated ghrelin than men in those who were lean

(Alajmi et al. 2016; Douglas et al. 2017) and in those who were overweight/obese (Douglas et al. 2017). Furthermore, an inverse relationship between general adiposity levels and fasting ghrelin levels has been suggested in study samples including individuals who were lean and individuals who were obese, possibly because of elevated insulin or leptin levels (Tschöp et al. 2001; Shiiya et al. 2002; Sondergaard et al. 2009). Individuals who are obese also exhibit a reduced postprandial suppression of ghrelin (Le Roux et al. 2005) and blunted postprandial increases in PYY (Le Roux et al. 2006). Limited evidence has also suggested an inverse association between visceral adipose tissue and fasting ghrelin levels in women who were lean and women who were obese, likely caused by substances secreted by visceral adipocytes, such as TNF α and leptin (Sondergaard et al. 2009). Moreover, fat-free mass, as the largest contributor to resting metabolic rate, has been identified as a key driver of appetite and energy intake in individuals who were lean and in individuals who were obese (Blundell et al. 2015b). In a systematic review including studies in individuals with normal weight, overweight or obesity, physical activity has also been suggested to alter the sensitivity of the appetite control system by enhancing meal-induced satiety which may facilitate energy balance over the long term (Beaulieu et al. 2016).

Comment #5: Objective: The second objective proposed " to explore potential associations between fasting and postprandial appetite outcomes and physiological and behavioral characteristics" was not completely answered in the results and conclusion. The results of the first objective are in table 1, figure 1, table 2 and table 3. In table 4, the authors associate fasting insulin, glucose and leptin with anthropometrics, metabolic and physical active parameters.

Author response #5: The second objective of the study is answered in the results section in page 17, lines 326 to 357, where all sex-specific Pearson's correlation coefficients between appetite-related outcomes and individual characteristics are summarised. Table 4 highlights where significant correlations were observed, namely the correlations between the individual characteristics and fasting insulin, glucose and leptin. Additionally, this objective is also addressed in the discussion section on lines 374-376 and on lines 465-499. Nevertheless, we have included a sentence in the conclusion of the manuscript which answers the second objective directly, as follows:

Discussion, page 24, lines 530-532: The associations between fasting and postprandial acylated ghrelin, total PYY and general or abdominal adiposity were also small, while fasting leptin, glucose and insulin and postprandial insulin concentrations were consistently and positively associated with adiposity outcomes.

Comment #6: Participants: Why did the authors add 1% of blacks people in the study sample? It is well known that blacks people have different body composition and energy metabolism than White Europeans and Asians. Why did they not exclude blacks people? This sample is not representative of the race.

Author response #6: We did not recruit participants based on ethnicity as it was expected that the vast majority of the study sample would be white Europeans, considering the general

population where the study was conducted. Excluding participants of black or Asian ethnicity did not alter the interpretation of our findings and, therefore, it was preferred to maintain the original study sample in order to increase the statistical power of our analyses.

Comment #7: Preliminary testing: Why did you use three skinfolds to estimate body composition? It is a doubly indirect method for estimating body composition.

Author response #7: We appreciate the reviewer's comment and we agree that skinfolds is an indirect method to estimate body composition which presents inherent limitations. However, we did not have access to other more accurate methods of assessing total body fat in such a large sample (e.g. BOD POD, DEXA). It is known that, when performed by a trained and experienced examiner, skinfold measurements can provide a reliable estimation of body fat mass. Additionally, we used body fat mass estimated by skinfolds in conjunction with BMI and body fat distribution assessed with high-quality MRI scans (visceral adipose tissue, abdominal subcutaneous adipose tissue and liver fat). Our approach of using three skinfold sites was based on the equation which has been validated for the population we recruited for the study. We have included a sentence in the methods section of the manuscript to highlight the care taken for the consistency of skinfold measurements, as follows:

Methods, page 5, lines 137-138: All skinfold measurements were performed by the same experienced examiner throughout the study.

Comment #8: Blood sampling and biochemical analysis - paragraph 1 - lines 9 and 10 - the authors describe "haemoglobin concentrations and hematocrit", but did not show results of these analysis.

Author response #8: Haemoglobin concentration and haematocrit were assessed to ensure any changes in plasma volume did not affect the quantification of blood parameters. As no exercise was performed during the study visit where blood samples were collected, we did not expect to observe any significant plasma volume changes and these analyses were performed for reassurance only. We have clarified that "Correction of blood parameter concentrations for acute changes in plasma volume had a negligible influence on our findings and, therefore, the unadjusted plasma concentrations are displayed for simplicity" in the statistical analysis section (Methods, page 9, lines 247-249).

Comment #9: Statistical analysis: The Hardy-Weinberg equilibrium was calculated?

Author response #9: We have calculated the genetic variation of our population using the Hardy-Weinberg equation and can confirm there was no significant deviation from Hardy-Weinberg equilibrium. This information has been added to the methods as indicated below. Furthermore, the prevalence of the three FTO rs9939609 genotypes in our study sample was similar to the prevalence reported previously by Frayling et al. 2007 in 13 cohorts with 38,759 participants: 16% of the population as AA (19% in our study), 37% as TT (36% in our study) and 47% as AT (45% in our study).

Methods, page 8, lines 219-221: Genotype frequency of FTO rs9939609 was assessed using a goodness-of-fit chi-square test and did not deviate from Hardy-Weinberg equilibrium ($\chi^2 = 0.435, P = 0.509$).

Comment #10: Participants characteristics: lines 3-5 - results are expected and do not need be discussed in detail.

Author response #10: The sentence summarizing the differences observed between men and women was removed from the text, as requested by the reviewer.

Comment #11: Figure 1 - results are not innovative, but I recommend that you keep the figure. It would be important to add the p-value in the figures.

Author response #11: We have kept the figure and highlighted where the P-value was lower than 0.05 between males and females.

Comment #12: Sex-specific Pearson - We lacked discussing the result of the insulin ratio with VO₂ and glucose with VO₂. The authors could talk in the context of energy metabolism.

Author response #12: We have now highlighted the associations between insulin and glucose with V̇O₂ peak in the discussion section, as follows:

Discussion, pages 22-23, lines 484-488: Additionally, negative associations between V̇O₂ peak and fasting and postprandial insulin, fasting glucose and fasting leptin were observed. Acute and chronic exercise augments insulin sensitivity by increasing insulin-like growth factor 1, and individuals with higher cardiorespiratory fitness typically show higher insulin sensitivity (Borghouts and Keizer, 2000; Castro et al. 2016).

Comment #13: Table 4 is extensive, with many correlations already expected. In addition, it was not the objective of the study. I suggest a careful review of the results for table 4! Many correlations were already expected and need not be highlighted. I suggest highlighting the correlations necessary to respond to the objectives proposed in the study.

Author response #13: Table 4 was included in order to summarize the significant associations observed between fasting insulin, glucose and leptin and individual characteristics, which answers the second objective of the study i.e. to explore potential associations between fasting and postprandial appetite outcomes and physiological and behavioural characteristics. However, the table can be included as supplementary online material if deemed appropriate by the reviewer and/or editor.

Comment #14: Discussion:

Paragraph 1 – line 11 - The authors said that they evaluated "lifestyle characteristics", but only the physical activity practice was evaluated.

Author response #14: We used the term ‘lifestyle characteristics’ to summarize the measurements of both habitual physical activity levels and sitting time.

Comment #15: Paragraph 2 - line 11 - the authors refer to "heterogeneous samples" to justify the difference of the results found in the present study and in Karra et al (2013). Does age influence the relationship of ghrelin to appetite?

Author response #15: Our sample was heterogeneous not only in terms of age, but also in adiposity parameters (as shown in Table 1), as well as including both males and females. On the contrary, the study performed by Karra et al. only included healthy young lean males with an average age of ~23 years. These differences in study samples might explain differences in the observed results, as previous evidence indicates ghrelin levels can vary between males and females and also according to body adiposity (as indicated in the manuscript’s introduction). Additionally, although evidence is limited, it has been suggested that the loss of appetite and decline in energy intake in older adults may be related to the concomitant elevation in circulating leptin and insulin and a reduction in ghrelin concentrations (Landi et al. *Nutrients*, 2016;8(2):69). We have clarified that the study of Karra et al. included only lean young males in the discussion section, as follows:

Discussion, pages 19-20, lines 389-391: Differences between study samples can possibly explain discrepancies between findings, as Karra et al. (2013) recruited healthy young lean males, while our sample was composed of a heterogeneous group of males and females.

Comment #16: I would suggest adding also a result of a recent study published with obese women in which "Participants with the AA genotype had lower values than those with TT and TA in the postprandial period." (Magno et al. , 2018).

Author response #16: We appreciate the reviewer’s suggestion and the reference to the study performed by Magno is included in the discussion section (page 20, lines 393-397).

Comment #17: Paragraph 3 - line 10 - review use of numbers 3-36 subscript!

Author response #17: We have presented ‘3-36’ in subscript to indicate the form of PYY that was measured in the study by Karra et al. (2013). PYY₃₋₃₆ is commonly reported in the literature with 3-36 presented in subscript; therefore, we feel ‘PYY₃₋₃₆’ will be familiar to the reader.

Comment #18: Paragraph 6 - line 3 - review "women had significantly lower fat mass and fat free mass" because women had higher fat mass. See table 1!

Author response #18: The sentence highlighted by the reviewer reads “It should be noted that all participants received an identical standardised meal and, as women had significantly lower body mass and fat free mass, and consequently lower resting metabolic rate, it was expected that the postprandial suppression of appetite would be stronger in women.”. We have not

mentioned fat mass in this sentence but highlighted that both body mass and fat free mass were lower in women than men which is supported by the data presented in Table 1.

Comment #19: Conclusion: The conclusion does not address the second objective proposed by the authors (association between fasting and postprandial appetite with physiological and behavioral characteristics).

Author response #19: We have included a sentence in the conclusion of the manuscript which answers the second objective of the study, as follows:

Discussion, page 24, lines 530-532: The associations between fasting and postprandial acylated ghrelin, total PYY and general or abdominal adiposity were also small, while fasting leptin, glucose and insulin and postprandial insulin concentrations were consistently and positively associated with adiposity outcomes.

Comment #20: References: The references of Carvalho et al (2018) and Melhorn et al (2018) were not found. Please review all other references!

Author response #20: We thank the reviewer for bringing this to our attention. All references have been reviewed accordingly.

Reviewer two:

Comment #1: In the current manuscript the authors seek to understand the role of the obesity-associated gene FTO on behavioral feeding phenotype and associated physiologic and metabolic parameters. Specifically multiple indices of appetite, feeding peptide levels in plasma (in fasted and fed state), fitness and metabolic rate in healthy and FTO-identified patients were performed. This is achieved through a combination of laboratory visits and data obtained from an accelerometer that patients wore while away from the lab. The authors should be commended on this effort. This topic is relevant to the field of obesity research and associated feeding pathologies. I offer my constructive criticisms here.

Author response #1: We thank the reviewer for the positive comments on our manuscript and we hope that the helpful comments below have been addressed appropriately.

Comment #2: I appreciate the care taken to measure acylated ghrelin across fed and fasted states. However cephalic ghrelin secretion in anticipation of meals was not measured. I bring this up because normalizing each patient by fasting does not evaluate conditioned or pre-meal ghrelin responses associated with anticipation of food. I think this should be qualified in the discussion.

Author response #2: We presented the appetite and plasma concentrations of acylated ghrelin, total PYY, insulin and glucose relative to baseline values (i.e., delta) to minimise the potential influence of day-to-day biological variability in these outcomes. However, given that participants knew when the meal would be provided, we cannot rule out that a preprandial increase in ghrelin may reflect an anticipatory signal for food intake rather than initiating meal intake (e.g., Cummings et al. 2001 Diabetes, 50: 1714-1719; Frecka & Mattes 2008 Am J Physiol Gastrointest Liver Physiol, 294: G699-707). Therefore, we have included this in the discussion section as follows:

Discussion, page 24, lines 522-526: Furthermore, participants were aware of the meal timing so it is possible that the higher preprandial ghrelin concentrations reflected an anticipatory response to impending meal intake (Cummings et al. 2001). Future studies should consider isolating meal provision from time-related cues and/or examining the influence of cephalic phase ghrelin release during meal anticipation on postprandial appetite responses.

Comment #3: Is it possible that a laboratory setting is not appropriate to measure FTO X obeseogenic food environment interactions known to promote maladaptive physiologic responses that induce obesity? Given the lack of interactions it would seem suitable to mention this in the discussion inline with targeted weight loss for example.

Author response #3: We thank the reviewer for raising this point. The aim of our study was to determine the influence of the FTO rs9969309 genotype on fasting and postprandial appetite-related hormones and, therefore, it was important to study participants in a controlled environment and in response to a standardised meal to minimise the influence of any potential confounding factors. However, we agree that the laboratory setting may not be appropriate to

determine the effect of the FTO rs9939609 genotype on food choice and eating behavior and we have highlighted this as a limitation and potential future direction in the discussion as follows:

Discussion, pages 23-24, lines 518-521: Additionally, it is possible that a study design where individuals are exposed to an obesigenic food environment, such as an *ad libitum* buffet meal rather than a standardised meal stimulus, may be more appropriate to elucidate the effect of the FTO rs9939609 genotype on food choice and eating behaviour

Comment #4: Separate from physiologic responses, psychological process are also regulators of food intake. For example, Dang et al. 2018, recently reported that AA individuals have higher food craving than controls, supporting the contention that in some cases food reward mechanisms may contribute to body weight gain in FTO individuals. Although the authors did not set out to test this aspect of feeding behavior, the discussion of physiologic versus psychological mechanisms would strengthen the conclusion.

Author response #4: We thank the reviewer for the suggestion and we have included the findings from Dang et al. in the discussion section as well as highlighting the importance of assessing psychological factors in future studies in the conclusion, as follows:

Discussion, page 20, lines 418-420: Moreover, recent evidence suggests that AA individuals show higher total food cravings, compared to TT individuals, which correlated with BMI (Dang et al. 2018).

Discussion, page 24, lines 532-534: Further research is needed to clarify the precise role of the FTO rs9939609 genotype in moderating appetite control and energy intake, including both physiological and psychological factors that influence eating behaviour.

Exploration of associations between the FTO rs9939609 genotype, fasting and postprandial appetite-related hormones and perceived appetite in healthy men and women

Fernanda R. Goltz^{1,2}, Alice E. Thackray^{1,2}, Veronica Varela-Mato¹, James A. King^{1,2}, James L. Dorling³, Monika Dowejko¹, Sarabjit Mastana¹, Julie Thompson^{1,2}, Greg Atkinson⁴, David J. Stensel^{1,2}

ABSTRACT

Background: The fat mass and obesity-associated gene (FTO) rs9939609 A-allele has been associated with obesity risk. Although the exact mechanisms involved remain unknown, the FTO rs9939609 A-allele has been associated with an impaired postprandial suppression of appetite. **Objectives:** To explore the influence of FTO rs9939609 genotype on fasting and postprandial appetite-related hormones and perceived appetite in a heterogeneous sample of men and women. **Design:** 112 healthy men and women aged 18-50-years-old completed three laboratory visits for the assessment of FTO rs9939609 genotype, body composition, aerobic fitness, resting metabolic rate, visceral adipose tissue, liver fat, fasting leptin, and fasting and postprandial acylated ghrelin, total PYY, insulin, glucose and perceived appetite. Participants wore accelerometers for seven consecutive days for the assessment of physical activity and sedentary behaviour. Multivariable general linear models quantified differences between FTO rs9939609 groups for fasting and postprandial appetite outcomes, with and without the addition of *a priori* selected physiological and behavioural covariates. Sex-specific univariable Pearson's correlation coefficients were quantified between the appetite-related outcomes and individual characteristics. **Results:** 95% confidence intervals for mean differences between FTO rs9939609 groups overlapped zero in unadjusted and adjusted general linear models for all fasting ($P \geq 0.28$) and postprandial ($P \geq 0.19$) appetite-related outcomes. Eta² values for explained variance attributable to FTO rs9939609 were <5% for all outcomes. An exploratory correlation matrix indicated that associations between fasting and postprandial acylated ghrelin, total PYY and general or abdominal adiposity were also small ($r = -0.23$ to 0.15 , $P \geq 0.09$). Fasting leptin, glucose and insulin and postprandial insulin concentrations were associated with adiposity outcomes ($r = 0.29$ to 0.81 , $P \leq 0.033$). **Conclusions:** Associations between the FTO rs9939609 genotype and fasting or postprandial appetite-related outcomes were weak in healthy men and women.

Keywords: FTO, appetite, ghrelin, PYY, hunger.

1
2
3
4 1 **Exploration of associations between the FTO rs9939609 genotype, fasting and**
5 2 **postprandial appetite-related hormones and perceived appetite in healthy men and**
6 3 **women**

8 4 Fernanda R. Goltz ^{1,2}, Alice E. Thackray ^{1,2}, Veronica Varela-Mato ¹, James A. King ^{1,2},
9 5 James L. Dorling ³, Monika Dowejko ¹, Sarabjit Mastana ¹, Julie Thompson ^{1,2}, Greg
10 6 Atkinson ⁴, David J. Stensel ^{1,2}

13 7
14 8 ¹ National Centre for Sport and Exercise Medicine, School of Sport, Exercise and Health
15 9 Sciences, Loughborough University, Loughborough, United Kingdom.

16 10 ² University Hospitals of Leicester NHS Trust, Infirmary Square, Leicester, United Kingdom.

17 11 ³ Ingestive Behavior Laboratory, Pennington Biomedical Research Center, Baton Rouge,
18 12 United States.

19 13 ⁴ School of Health and Social Care, Teesside University, Middlesbrough, United Kingdom.
20 14

21 15 **Corresponding author:**

22 16 Professor David Stensel

23 17 School of Sport, Exercise and Health Sciences

24 18 Loughborough University

25 19 Leicestershire

26 20 LE11 3TU

27 21 United Kingdom

28 22 Phone: +44(0)1509 226344, Fax: +44(0)1509 226301, E-mail: D.J.Stensel@lboro.ac.uk
29 23

30 24 **Declarations of interest:** None.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

60
61
62
63 **ABSTRACT**
64
65

66 **Background:** The fat mass and obesity-associated gene (FTO) rs9939609 A-allele has been
67 associated with obesity risk. Although the exact mechanisms involved remain unknown, the FTO
68 rs9939609 A-allele has been associated with an impaired postprandial suppression of appetite.
69
70

71 **Objectives:** To explore the influence of FTO rs9939609 genotype on fasting and postprandial
72 appetite-related hormones and perceived appetite in a heterogeneous sample of men and women.
73

74 **Design:** 112 healthy men and women aged 18-50-years-old completed three laboratory visits for
75 the assessment of FTO rs9939609 genotype, body composition, aerobic fitness, resting
76 metabolic rate, visceral adipose tissue, liver fat, fasting leptin, and fasting and postprandial
77 acylated ghrelin, total PYY, insulin, glucose and perceived appetite. Participants wore
78 accelerometers for seven consecutive days for the assessment of physical activity and sedentary
79 behaviour. Multivariable general linear models quantified differences between FTO rs9939609
80 groups for fasting and postprandial appetite outcomes, with and without the addition of *a priori*
81 selected physiological and behavioural covariates. Sex-specific univariable Pearson's correlation
82 coefficients were quantified between the appetite-related outcomes and individual characteristics.
83

84 **Results:** 95% confidence intervals for mean differences between FTO rs9939609 groups
85 overlapped zero in unadjusted and adjusted general linear models for all fasting ($P \geq 0.28$) and
86 postprandial ($P \geq 0.19$) appetite-related outcomes. Eta² values for explained variance attributable
87 to FTO rs9939609 were <5% for all outcomes. An exploratory correlation matrix indicated that
88 associations between fasting and postprandial acylated ghrelin, total PYY and general or
89 abdominal adiposity were also small ($r = -0.23$ to 0.15 , $P \geq 0.09$). Fasting leptin, glucose and
90 insulin and postprandial insulin concentrations were associated with adiposity outcomes ($r =$
91 0.29 to 0.81 , $P \leq 0.033$). **Conclusions:** Associations between the FTO rs9939609 genotype and
92 fasting or postprandial appetite-related outcomes were weak in healthy men and women.
93
94
95
96
97
98
99

100
101
102
103
104 **Keywords:** FTO, appetite, ghrelin, PYY, hunger.
105
106
107
108
109
110
111
112
113
114
115
116
117
118

119
120
121
122 **52 INTRODUCTION**
123

124 53 The scientific understanding of appetite control has increased considerably in recent decades,
125
126 54 which has been helpful in elucidating the complex nature of energy balance and weight control.
127
128 55 Central components of the homeostatic control of appetite comprise signals from adipose tissue
129 56 and peptide hormones secreted from the digestive tract, which act acutely and/or chronically on
130
131 57 central neural pathways to influence hunger, satiety and subsequent energy intake (MacLean et
132 58 al. 2017). These signals and hormones include the tonic signals leptin and insulin that regulate
133
134 59 long-term changes in energy balance and adiposity status, as well as a variety of episodic gut
135 60 signals, which mediate hunger and satiety on a meal-by-meal basis (Blundell et al. 2008, 2015a;
136
137 61 MacLean et al. 2017). Notable among the episodic mediators of appetite and energy intake are
138
139 62 acylated ghrelin and peptide YY (PYY) which exert orexigenic and anorexigenic effects,
140 63 respectively, to facilitate meal initiation and termination (Neary and Batterham, 2009).
141

142 64 Over the last 16 years, our laboratory has measured circulating concentrations of appetite-related
143
144 65 hormones in response to meal ingestion in many studies. A consistent observation from this body
145 66 of work is the degree of variability in responses observed between participants studied under
146
147 67 identical conditions. Furthermore, using the “gold standard” replicated crossover study design
148
149 68 (Atkinson and Batterham, 2015; Senn, 2016), we have demonstrated recently the presence of
150 69 true interindividual heterogeneity in appetite perceptions and circulating concentrations of
151
152 70 acylated ghrelin, total PYY, insulin and glucose in response to a standardised meal, over and
153 71 above any random within-subject variability and measurement error (Goltz et al. 2019). Similar
154
155 72 findings were also observed in acylated ghrelin, total PYY and perceived appetite responses to
156
157 73 replicated single bouts of aerobic exercise (Goltz et al. 2018).

158 74 The factors responsible for interindividual variability in appetite-related hormone concentrations
159
160 75 are not fully understood, but it is plausible that differences in individual characteristics and
161
162 76 behaviours may contribute to the variability observed. In this regard, the fat mass and obesity-
163 77 associated gene (FTO) has been associated with obesity risk, with individuals homozygous for
164
165 78 the A allele (AA) of FTO rs9939609 having a 1.7-fold higher obesity risk than individuals
166
167 79 homozygous for the T allele (TT) (Frayling et al. 2007). Although the exact mechanisms through
168 80 which **FTO rs9939609** influences fat mass accumulation remain unknown, it has been suggested
169
170 81 that it exerts its effect on food intake rather than on energy expenditure (Speakman et al. 2008).
171
172 82 Furthermore, **rs9939609** AA individuals have been shown to exhibit an attenuated postprandial
173 83 suppression of hunger and acylated ghrelin compared with TT individuals, which may
174
175
176
177

178
179
180
181 84 predispose AA individuals to higher energy intake and, consequently, higher fat mass (Karra et
182 al. 2013). However, the study by Karra and colleagues was performed in young healthy weight
183 85 males and it is not known whether this influence of the FTO rs9939609 gene on postprandial
184 86 appetite regulation is observed in a heterogenous sample of men and women.
185
186 87

187
188 88 Beyond genetic influence, it has been speculated that other individual factors may affect appetite
189 regulation. Data from previous studies have indicated that women exhibit higher fasting
190 89 concentrations of acylated ghrelin than men in those who were lean (Alajmi et al. 2016; Douglas
191 90 et al. 2017) and in those who were overweight/obese (Douglas et al. 2017). Furthermore, an
192 inverse relationship between general adiposity levels and fasting ghrelin levels has been
193 91 suggested in study samples including individuals who were lean and individuals who were obese,
194 92 possibly because of elevated insulin or leptin levels (Tschöp et al. 2001; Shiiya et al. 2002;
195 93 Sondergaard et al. 2009). Individuals who are obese also exhibit a reduced postprandial
196 94 suppression of ghrelin (Le Roux et al. 2005) and blunted postprandial increases in PYY (Le
197 95 Roux et al. 2006). Limited evidence has also suggested an inverse association between visceral
198 96 adipose tissue and fasting ghrelin levels in women who were lean and women who were obese,
199 97 likely caused by substances secreted by visceral adipocytes, such as TNF α and leptin
200 98 (Sondergaard et al. 2009). Moreover, fat-free mass, as the largest contributor to resting metabolic
201 99 rate, has been identified as a key driver of appetite and energy intake in individuals who were
202 100 lean and in individuals who were obese (Blundell et al. 2015b). In a systematic review including
203 101 studies in individuals with normal weight, overweight or obesity, physical activity has also been
204 102 suggested to alter the sensitivity of the appetite control system by enhancing meal-induced
205 103 satiety which may facilitate energy balance over the long term (Beaulieu et al. 2016). Together,
206 104 these findings highlight the importance of investigating the effect of the FTO rs9939609 gene
207 105 on appetite parameters in a sample of males and females with a wide range of age, adiposity and
208 106 physical activity levels, including physiological and behavioural characteristics as covariates in
209 107 the analyses.
210
211

212 108 The primary aim of this study was to use objective assessment methods in order to explore the
213 109 influence of the FTO rs9939609 genotype on fasting and postprandial appetite-related hormones
214 110 and perceived appetite in a sample of healthy men and women. The secondary aim was to explore
215 111 potential associations between fasting and postprandial appetite outcomes and physiological and
216 112 behavioural characteristics.
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236

237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295

METHODS

Participants

With the approval of the University Ethics Advisory Sub-Committee, a total of 121 participants (57 men, 64 women) aged 18 to 50 years provided written informed consent before taking part in the study. All participants were deemed to be stable in their body mass (≤ 3 kg change in the previous 3 months), non-smokers, habitual breakfast eaters, had no history of cardiovascular or metabolic disease, and were not dieting or taking any medications known to influence the outcome measures. Female participants were premenopausal and postmenopausal and not pregnant. Nine participants withdrew from the study before completing all study measurements due to time constraints. Therefore, data are presented for 112 participants (56 men, 56 women) in this manuscript. The study sample self-reported ethnicity distribution was as follows: 93% white Europeans, 6% Asians and 1% black.

Visit 1: Preliminary testing

Participants attended the laboratory for a preliminary visit to confirm eligibility, and to undergo familiarisation, anthropometric measurements and determination of peak oxygen uptake ($\dot{V}\text{O}_2$ peak). The eligibility assessment included screening questionnaires to assess health status and food preferences and/or restrictions. Stature was measured to the nearest 0.1 cm and body mass to the nearest 0.1 kg using an electronic measuring station (Seca, Hamburg, Germany), and body mass index (BMI) was calculated. The sum of three skinfolds (chest, abdomen and thigh for men, and triceps, suprailiac and thigh for women) was used to estimate body density (Jackson and Pollock 1978, 1980) and body fat percentage (Siri, 1961). All skinfold measurements were performed by the same experienced examiner throughout the study. Waist circumference was measured as the narrowest point between the lower rib margin and the iliac crest.

Participants were familiarised with walking and running on the treadmill (Technogym Excite Med, Cesena, Italy) before completing an incremental uphill treadmill protocol to determine $\dot{V}\text{O}_2$ peak. The participants ran at a fixed individualised speed (4.5 to 14.0 $\text{km}\cdot\text{h}^{-1}$), with the initial gradient of the treadmill set to 0%. The treadmill gradient was increased by 1% every minute until volitional exhaustion. Heart rate was monitored continuously using short-range telemetry (Polar A3, Kempele, Finland), and ratings of perceived exertion (Borg, 1973) were recorded at the end of each minute. Expired air samples were monitored continuously using a breath-by-breath gas analysis system (Cortex Metalyser 3B, Leipzig, Germany). An average of

296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354

the breath-by-breath oxygen uptake data was taken every 10 s, and $\dot{V}O_2$ peak was defined as the highest 30 s rolling average.

Visit 2: Magnetic resonance imaging (MRI) scan

Each participant underwent an MRI scan in the supine position using a dual-echo Dixon fat and water sequence on a 3-T MRI scanner (MR750w, GE Healthcare, Chicago, USA). A detailed description of the protocol has been reported previously (Borga et al. 2015; West et al. 2016). Briefly seven overlapping image stacks were acquired from the neck to knee with stacks covering the abdomen (stacks 2 to 5) acquired during breath-hold. Additional abdominal slices were acquired with the IDEAL-IQ sequence to assess proton density fat fraction in the liver. Scans were analysed to quantify visceral adipose tissue, abdominal subcutaneous adipose tissue and liver fat fraction using the AMRA Profiler (AMRA Medical AB, Linköping, Sweden) (Borga et al. 2015; West et al. 2016).

Visit 3: Resting metabolic rate and test meal

All premenopausal female participants completed the main trial during the follicular phase of the menstrual cycle (days 6-12) to avoid potential hormonal influences on appetite parameters. Participants were asked to refrain from caffeine, alcohol, and strenuous exercise during the 24 h before the main trial. A standardised evening meal (3297 kJ, 40% fat, 39% carbohydrate, 21% protein) was consumed the evening before the main trial and only plain water was permitted after the meal until participants arrived at the laboratory the next day.

Participants reported to the laboratory at 08:00 after fasting overnight for 12 h. A cannula (Venflon; Becton Dickinson, Helsingborg, Sweden) was inserted into an antecubital vein for venous blood sampling, and participants rested for 60 min to eliminate any stress effects in response to the cannula (Chandarana et al. 2009). During this time, resting metabolic rate was measured using an open circuit indirect calorimetry system (GEM Nutrition Ltd., Cheshire, England). Participants were asked to lie in a comfortable supine position and were instructed not to talk or sleep, and to move as little as possible during the measurement. The clear hood canopy was placed over the head area, and plastic sheeting attached to the hood was placed around the body to form a seal between the air inside and outside the hood. Oxygen uptake, carbon dioxide production, respiratory exchange ratio and energy expenditure were determined at 30 s intervals over a 30 min period. The first 10 min of data was discarded to account for any initial short-term respiratory artefact.

355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413

A fasting venous blood sample and rating of perceived appetite were taken 60 min after the insertion of the cannula. Participants then consumed a standardised breakfast within 15 min marking the start of the postprandial assessment period (09:00; 0 h). Breakfast consisted of a ham and cheese sandwich, milkshake and chocolate biscuit which provided 4435 kJ of energy (41% carbohydrate, 18% protein, 41% fat). Subsequent venous blood samples and ratings of perceived appetite were taken at 0.5, 1 and 2 h after the start of the breakfast whilst the participants rested in a semi-supine position.

Appetite perceptions

Appetite perceptions (hunger, satisfaction, fullness, prospective food consumption) were assessed using 100 mm visual analogue scales (Flint et al. 2000). An overall appetite rating was calculated as the mean value of the four appetite ratings once satisfaction and fullness were reverse-scored (Stubbs et al. 2000).

Blood sampling and biochemical analysis

Venous blood samples were collected into pre-chilled EDTA monovettes (Sarstedt, Leicester, UK) for the determination of plasma acylated ghrelin, total PYY, leptin, insulin and glucose concentrations. Monovettes for acylated ghrelin also contained *p*-hydroxymercuribenzoic acid to prevent the degradation of acylated ghrelin by protease and were centrifuged at 2,383 g for 10 min at 4°C (Burkard, Hertfordshire, UK). The plasma supernatant was aliquoted into a storage tube and 100 µL of 1 M hydrochloric acid was added per millilitre of plasma. Samples were re-centrifuged at 2,383 g for 5 min at 4°C before being transferred into Eppendorf tubes and stored at -80°C for later analysis. Monovettes for total PYY, leptin, insulin and glucose were centrifuged immediately at 2,383 g for 10 min at 4°C prior to storage at -80°C. Haemoglobin concentration and haematocrit were quantified in duplicate at 0 and 2 h to estimate the acute change in plasma volume (Dill and Costill, 1974).

Commercially available enzyme-linked immunosorbent assays were used to determine the concentrations of plasma acylated ghrelin (Bertin Bioreagent, Montigney le Bretonneux, France), total PYY (Millipore, Billerica, MA, USA), leptin (R&D Systems, Minneapolis, MN, USA) and insulin (Mercodia, Uppsala, Sweden). Plasma glucose concentrations were determined by enzymatic, colorimetric methods using a benchtop analyser (Pentra 400, HORIBA Medical, Montpellier, France). The within-batch coefficient of variation for acylated ghrelin, total PYY, leptin, insulin and glucose concentrations were 4.3%, 5.1%, 8.3%, 4.7%, 0.4%, respectively.

414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472

An additional fasting venous blood sample was collected into a 2.7-mL EDTA monovette (Sarstedt, Leicester, UK) and the whole blood sample was stored at 4°C to undergo DNA extraction and genotyping. Genomic DNA was extracted from the whole blood samples using the QIAamp DNA Mini kit (QIAGEN, Hilden, Germany). The samples were genotyped for the rs9939609 allele within the FTO gene using the Applied Biosystems TaqMan® (Roche Molecular Systems, Pleasanton, California, USA) genotyping assay and real-time polymerase chain reaction system. Participants were assigned to one of three groups according to their genotype: homozygous major allele, TT (36%; males $n = 23$, females $n = 17$); heterozygous allele, AT (45%; males $n = 22$, females $n = 29$); or homozygous minor allele, AA (19%; males $n = 11$, females $n = 10$). Genotype frequency of FTO rs9939609 was assessed using a goodness-of-fit chi-square test and did not deviate from Hardy-Weinberg equilibrium ($\chi^2 = 0.435$, $P = 0.509$).

Habitual physical activity and sedentary time

Participants wore an ActiGraph GT3X+ accelerometer (ActiGraph, Pensacola, USA) on an elasticated belt on the waist above the mid-line of the thigh on their non-dominant side of the body. The device was initialised at a frequency of 100HZ and downloaded using ActiLife software v6.11.8 and firmware version 2.0.0. ActiGraph data were downloaded in 60-seconds epochs and physical activity was classified as low, light and moderate-to-vigorous. Participants also wore an activPAL3 accelerometer (PAL Technologies Ltd., Glasgow, UK), attached directly to the skin on the midline of the anterior aspect of the thigh in line with the ActiGraph GT3X+ accelerometer. The activPAL3 determines posture using information derived from accelerations of the thigh, including the gravitational component, using a triaxial accelerometer (Atkin et al. 2012). The activPAL3 is a valid measure of time spent sitting/lying, standing, and walking in adults (Kozey-Keadle et al. 2011). ActivPAL3 sitting time data were retrieved and clustered into 60-seconds epochs using a customized spreadsheet. Participants were advised to wear both devices concurrently and continuously over a 7-day period. Non-wear time and sleep time were removed from the analysis and moderate-to-vigorous physical activity (MVPA) and sitting time data were averaged over the seven-day period.

Statistical analyses

We estimated the effect size detection sensitivity given our sample size using NQuery (version 3, Statistical Solutions, Cork, Ireland). For a total sample size of 110 and three study groups, we

473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531

estimated that a “medium” (Cohen, 1998) η^2 value of 0.18 would be detected in a univariable model as statistically significant ($P < 0.050$) with power of 90%.

Postprandial overall appetite and plasma concentrations of acylated ghrelin, total PYY, insulin and glucose are presented relative to baseline values (delta) to minimise the potential influence of day-to-day biological variability (Deighton et al. 2013, 2014). Total area under the curve (AUC) values were calculated using the trapezoidal method. Correction of blood parameter concentrations for acute changes in plasma volume had a negligible influence on our findings and, therefore, the unadjusted plasma concentrations are displayed for simplicity.

Multivariable general linear models were used to quantify the mean differences (and 95% confidence intervals) between FTO rs9939609 genotype groups for each fasting and postprandial appetite outcome. The eta-squared statistic (with associated 90% confidence interval) was also estimated for each model and each outcome (Kline, 2004; Steiger, 2004). This statistic is interpreted in a similar way as the coefficient of determination, where $100 \times \eta^2$ gives the explained variance attributable to the FTO groups. A 90% rather than a 95% confidence interval is reported because the eta-squared statistic can only be positive in sign. The model residuals of the appetite outcome variables were explored for parity to a Gaussian distribution using histograms. The model residuals for fasting acylated ghrelin and insulin concentrations were observed to show a positively skewed distribution so these data were logarithmically-transformed prior to analysis (Bland and Altman, 1996). Three models were used for each of the fasting and postprandial appetite outcomes, as follows:

1. Model I: Univariable models with FTO rs9939609 genotype as single fixed effect;
2. Model II: A multivariable model based on the selection of matched covariates studied by Karra et al. (2013), i.e., age, fat mass and visceral adipose tissue. FTO rs9939609 genotype was entered as a fixed effect and sex, age, fat mass and visceral adipose tissue were entered as covariates;
3. Model III: A multivariable model, where FTO rs9939609 genotype was entered as a fixed effect and sex, age, BMI, $\dot{V}O_2$ peak, resting metabolic rate, visceral adipose tissue, abdominal subcutaneous adipose tissue, liver fat, sitting time and MVPA were entered as covariates. Rather than the now discouraged use of stepwise selection procedures, these covariates were included based on their hypothesised influence on the outcome variables, while considering the potential that some predictors were mathematically coupled (Flom and Cassell, 2007; Whittingham et al. 2006). For

532
533
534
535 273 example, total fat mass was excluded from this model because multiple specific
536 adiposity parameters were considered.
537 274
538

539 275 The covariates in models II and III were each standardised prior to analysis by dividing each
540 datum by twice the respective SD (Gelman and Pardoe, 2007). In sensitivity analyses, model III
541 276 was also run with (i) waist circumference replacing BMI; (ii) percentage body fat replacing BMI;
542 277 and (iii) with a sex-by-genotype interaction term.
543
544 278
545

546 279 Univariable general linear models with FTO rs9939609 genotype as a single fixed effect were
547 used to quantify differences between genotype groups for body mass, BMI and fat mass.
548 280 Between-sex differences in participant characteristics and appetite-related outcomes in the
549 281 fasting and postprandial states were assessed using univariable general linear models with sex
550 as a single fixed effect. Sex-specific univariable Pearson's correlation coefficients were
551 282 quantified between appetite-related outcomes and individual characteristics, and between
552 283 appetite-related blood parameters and perceived appetite.
553
554 284
555 285

556 95% confidence intervals (95% CI) were quantified for correlation coefficients. P-values are
557 286 expressed in exact terms apart from very low values, which are expressed as $P < 0.001$. A
558 threshold of statistical significance was accepted as $P < 0.050$, although we deemed a P value of
559 287 < 0.005 as a stronger indication of potentially more reproducible results in line with recent advice
560 288 (Benjamin et al. 2017). All statistical analyses were performed in SPSS (v.23, IBM Corporation,
561
562 289 New York, USA).
563
564 290
565 291
566
567 292
568

569 293 **RESULTS**

570 571 294 **Missing data**

572
573 295 Due to technical issues with the equipment, resting metabolic rate is presented for 107
574 participants (53 males), sitting time for 96 participants (47 males) and MVPA for 100
575 296 participants (49 males). Eleven participants were unable to undertake the MRI scan for safety
576 297 reasons and, therefore, visceral adipose tissue and abdominal subcutaneous adipose tissue are
577 presented for 101 participants (50 males). Liver fat could not be quantified from some images
578 298 due to motion artefacts and, therefore, data is presented for 97 participants (48 males).
579
580 299
581 300
582

583 301 **Participant characteristics and appetite-related outcomes**

584
585 302 Participant characteristics, perceived appetite and appetite-related blood parameters in the
586 fasting and postprandial states are presented in Table 1. Postprandial delta values for acylated
587 303
588
589
590

591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649

304 ghrelin, total PYY, insulin and glucose concentrations and perceived overall appetite are
305 presented in Figure 1.

650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708

Table 1. Participant characteristics and appetite outcomes in the fasting and postprandial states.

	All (n = 112)	Range (min to max)	Men (n = 56)	Women (n = 56)	P	Mean difference 95% CI
Age (years)	34 (9)	18 to 50	35.3 (9.7)	33.5 (9.1)	0.303	-5.4 to 1.7
Stature (cm)	171.0 (9.2)	149.1 to 200.4	178.5 (6.6)	165.3 (6.2)	< 0.001	-15.6 to -10.8
Body mass (kg)	74.9 (14.7)	48.5 to 140.4	83.3 (12.9)	66.5 (11.1)	< 0.001	-21.2 to -12.2
Body mass index (kg·m ⁻²)	25.2 (3.9)	18.4 to 40.3	26.1 (3.7)	24.4 (4.0)	0.016	-3.2 to -0.3
Waist circumference (cm)	82.7 (10.8)	62.4 to 125.0	88.4 (9.8)	77.0 (8.7)	< 0.001	-14.9 to -8.0
Fat mass (kg)	16.9 (8.4)	3.5 to 47.8	15.5 (9.1)	18.2 (7.4)	0.078	-0.3 to 5.9
Fat free mass (kg)	58.1 (12.2)	36.8 to 92.6	67.8 (8.8)	48.3 (5.5)	< 0.001	-22.2 to -16.8
V̇O ₂ peak (mL·kg ⁻¹ ·min ⁻¹)	44.0 (9.3)	21.0 to 81.0	49.0 (9.3)	39.0 (6.1)	< 0.001	-13.0 to -7.1
Resting metabolic rate (kcal)*	1617 (322)	889 to 2567	1808 (290)	1430 (232)	< 0.001	-478 to -277
Visceral adipose tissue (L)*	1.70 (1.26)	0.11 to 6.22	2.27 (1.41)	1.14 (0.75)	< 0.001	-1.58 to -0.69
Abdominal subcutaneous adipose tissue (L)*	5.39 (3.02)	1.45 to 16.86	4.49 (2.39)	6.27 (3.33)	0.003	0.64 to 2.93
Liver fat (%)*	2.12 (1.81)	0.46 to 10.45	2.62 (2.19)	1.63 (1.16)	0.006	-1.69 to -0.28
Sitting time (min·day ⁻¹)*	509 (85)	256 to 737	513 (73)	504 (95)	0.630	-43 to 26
MVPA (min·day ⁻¹)*	55 (31)	11 to 163	57 (30)	54 (33)	0.706	-15 to 10
Fasting leptin (ng·mL ⁻¹)	8.62 (8.63)	1.34 to 43.85	4.07 (3.08)	13.16 (9.95)	< 0.001	6.33 to 11.84
Fasting acylated ghrelin (pg·mL ⁻¹)	173.6 (491.8)	12.0 to 4410.6	103.3 (108.8)	243.8 (682.9)	0.131	-42.6 to 323.6
Fasting total PYY (pg·mL ⁻¹)	117.5 (50.5)	13.6 to 270.0	121.9 (47.9)	113.0 (53.1)	0.353	-27.8 to 10.0
Fasting insulin (pmol·L ⁻¹)	23.3 (15.0)	2.9 to 97.1	22.9 (14.3)	23.6 (15.8)	0.825	-5.0 to 6.3
Fasting glucose (mmol·L ⁻¹)	5.24 (0.43)	4.29 to 6.56	5.37 (0.43)	5.12 (0.39)	0.001	-0.41 to -0.10
Fasting overall appetite (mm)	70.8 (15.3)	19 to 95	71.2 (13.4)	70.4 (17.1)	0.787	-6.5 to 5.0
Acylated ghrelin delta AUC (2 h, pg·mL ⁻¹)	-87.9 (126.6)	-1183.5 to 165.8	- 51.3 (56.3)	- 124.6 (162.6)	0.002	-118.9 to -27.8
Total PYY delta AUC (2 h, pg·mL ⁻¹)	101.6 (61.0)	-26.4 to 340.7	99.0 (62.4)	104.2 (59.9)	0.653	-17.7 to 28.1
Insulin delta AUC (2 h, pg·mL ⁻¹)	420.6 (236.8)	121.3 to 1485.8	403.9 (256.6)	437.3 (216.3)	0.458	-55.5 to 122.2
Glucose delta AUC (2 h, pg·mL ⁻¹)	0.77 (1.59)	-2.20 to 5.79	0.54 (1.37)	1.00 (1.77)	0.125	-0.13 to 1.05
Overall appetite delta AUC (2 h, pg·mL ⁻¹)	-77.4 (34.4)	-150.0 to -14.0	-65.7 (30.9)	-89.1 (34.0)	< 0.001	-35.5 to -11.1

Values are mean (SD). P values and 95% CI are from univariable general linear models with sex as a single fixed effect.
* n = 107 (53 males) for resting metabolic rate, 96 (47 males) for sitting time, 100 (49 males) for MVPA, 101 (50 males) for visceral adipose tissue and abdominal subcutaneous adipose tissue, and 97 (48 males) for liver fat.
AUC, area under the curve; CI, confidence interval; MVPA, moderate-to-vigorous physical activity, PYY, peptide YY; V̇O₂ peak, peak oxygen uptake.

709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749




Figure 1. Delta postprandial values for acylated ghrelin (A), total peptide YY (PYY) (B), insulin (C), glucose (D) and overall perceived appetite (E) in 56 males and 56 females. Grey rectangles indicate meal consumed within 15 min. Values are presented as mean (SD). * indicates P < 0.05 between males and females.

Univariable and multivariable general linear models

No statistically significant influence of the FTO rs9939609 genotype was identified for body mass ($\text{Eta}^2 = 0.027$, $P = 0.234$), BMI ($\text{Eta}^2 = 0.003$, $P = 0.688$) or fat mass ($\text{Eta}^2 = 0.025$, $P = 0.259$).

Fasting appetite-related outcomes

Separate univariate modelling (model I) did not reveal any statistically significant influence of the FTO rs9939609 genotype on fasting acylated ghrelin, total PYY, insulin, glucose, leptin or overall appetite ($P \geq 0.501$) (Table 2). Similarly, no significant effect of the FTO rs9939609 genotype was detected on fasting appetite-related outcomes in model II ($P \geq 0.098$) or model III ($P \geq 0.453$) (Table 2). All eta-squared values were very low (< 0.05). Replacing BMI with waist circumference, replacing BMI with body fat percentage, and including a sex-by-genotype interaction term in the sensitivity analyses did not result in a significant effect of the FTO rs9939609 genotype on any of the fasting appetite-related outcomes ($P \geq 0.470$, $P \geq 0.437$, $P \geq 0.455$, respectively).

Postprandial appetite-related outcomes

Separate univariate modelling (model I) did not reveal any statistically significant influence of the FTO rs9939609 genotype on delta AUC for acylated ghrelin, total PYY, insulin, glucose, leptin or overall appetite ($P \geq 0.322$) (Table 3). Similarly, no significant effect of the FTO rs9939609 genotype was detected on delta AUC for any of the appetite-related outcomes in model II ($P \geq 0.271$) or model III ($P \geq 0.186$) (Table 3). Again, all eta-squared values were very low (< 0.05). Replacing BMI with waist circumference, replacing BMI with body fat percentage, and including a sex-by-genotype interaction term in the sensitivity analyses did not result in a significant effect of the FTO rs9939609 genotype on any of the postprandial appetite-related outcomes ($P \geq 0.133$, $P \geq 0.102$, $P \geq 0.206$, respectively). A sensitivity analysis was undertaken on all the postprandial outcomes AUC by adding the respective fasting measurement as a covariate to the model. Again, no statistically significant differences between FTO groups could be detected ($P > 0.200$) and mean differences were small.

Table 2. Estimated marginal means from the multivariable general linear models used to quantify the differences between FTO rs9939609 genotype groups in each fasting appetite outcome.

	Model I			Model II			Model III		
	AT (n = 49)	AA (n = 21)	TT (n = 40)	AT (n = 45)	AA (n = 18)	TT (n = 37)	AT (n = 34)	AA (n = 17)	TT (n = 28)
Fasting acylated ghrelin (log pg·mL ⁻¹)	4.47 (4.25 to 4.69) Eta ² = 0.003 (90% CI: 0.000-0.023), <i>P</i> = 0.835	4.59 (4.26 to 4.92)	4.51 (4.27 to 4.75)	4.42 (4.18 to 4.65) Eta ² = 0.009 (90% CI: 0.000-0.047), <i>P</i> = 0.660	4.57 (4.20 to 4.94)	4.57 (4.30 to 4.83)	4.42 (4.20 to 4.64) Eta ² = 0.024 (90% CI: 0.000-0.091), <i>P</i> = 0.453	4.56 (4.23 to 4.88)	4.29 (4.03 to 4.54)
Fasting total PYY (pg·mL ⁻¹)	110.3 (96.1 to 124.5) Eta ² = 0.013 (90% CI: 0.000-0.055), <i>P</i> = 0.501	123.5 (101.8 to 145.2)	120.4 (104.7 to 136.2)	109.2 (94.0 to 124.4) Eta ² = 0.018 (90% CI: 0.000-0.069), <i>P</i> = 0.434	123.6 (100.2 to 147.0)	122.4 (105.7 to 139.1)	114.3 (97.6 to 130.9) Eta ² = 0.001 (90% CI: 0.000-0.014), <i>P</i> = 0.977	117.2 (93.3 to 141.0)	114.1 (95.0 to 133.2)
Fasting insulin (log pmol·L ⁻¹)	3.00 (2.83 to 3.16) Eta ² = 0.007 (90% CI: 0.000-0.038), <i>P</i> = 0.699	2.87 (2.61 to 3.12)	2.97 (2.79 to 3.16)	3.03 (2.88 to 3.19) Eta ² = 0.007 (90% CI: 0.000-0.041), <i>P</i> = 0.716	2.93 (2.70 to 3.17)	2.96 (2.79 to 3.13)	3.01 (2.81 to 3.20) Eta ² = 0.002 (90% CI: 0.000-0.028), <i>P</i> = 0.935	2.98 (2.70 to 3.27)	2.95 (2.72 to 3.18)
Fasting glucose (mmol·L ⁻¹)	5.23 (5.11 to 5.36) Eta ² = 0.002 (90% CI: 0.000-0.016), <i>P</i> = 0.882	5.28 (5.09 to 5.47)	5.22 (5.09 to 5.36)	5.27 (5.15 to 5.38) Eta ² = 0.027 (90% CI: 0.000-0.087), <i>P</i> = 0.278	5.28 (5.11 to 5.46)	5.14 (5.02 to 5.27)	5.24 (5.10 to 5.38) Eta ² = 0.018 (90% CI: 0.000-0.078), <i>P</i> = 0.553	5.30 (5.10 to 5.51)	5.16 (5.00 to 5.32)
Fasting leptin (ng·mL ⁻¹)	9.17 (6.70 to 11.65) Eta ² = 0.005 (90% CI: 0.000-0.030), <i>P</i> = 0.779	8.06 (4.27 to 11.84)	7.95 (5.21 to 10.69)	9.77 (8.15 to 11.39) Eta ² = 0.049 (90% CI: 0.000-0.122), <i>P</i> = 0.098	6.67 (4.17 to 9.17)	7.93 (6.15 to 9.71)	9.76 (7.91 to 11.62) Eta ² = 0.010 (90% CI: 0.000-0.057), <i>P</i> = 0.713	8.71 (6.05 to 11.37)	8.72 (6.59 to 10.85)
Fasting overall appetite (mm)	70.0 (65.7 to 74.4) Eta ² = 0.005 (90% CI: 0.000-0.033), <i>P</i> = 0.748	69.6 (63.0 to 76.2)	72.2 (67.4 to 77.0)	67.6 (63.0 to 72.3) Eta ² = 0.019 (90% CI: 0.000-0.072), <i>P</i> = 0.402	70.2 (63.0 to 77.4)	72.4 (67.3 to 77.6)	66.8 (60.9 to 72.7) Eta ² = 0.005 (90% CI: 0.000-0.034), <i>P</i> = 0.850	68.9 (60.4 to 77.3)	69.3 (62.5 to 76.0)

Model I: Univariable model with FTO rs9939609 genotype as single fixed effect. Model II: Multivariable model with FTO rs9939609 genotype as a fixed effect and sex, age, fat mass and visceral adipose tissue as covariates. Model III: Multivariable model with FTO rs9939609 genotype as a fixed effect and sex, age, body mass index, peak oxygen uptake, resting metabolic rate, visceral adipose tissue, abdominal subcutaneous adipose tissue, liver fat, sitting time and moderate-to-vigorous physical activity as covariates.

Values are mean (95% confidence interval (CI)). Eta², 90% CI and *P*-values are from the fixed effect of the FTO rs9939609 genotype group.

850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890

Table 3. Estimated marginal means from the multivariable general linear models used to quantify the differences between FTO rs9939609 genotype groups in each postprandial appetite outcome.

	Model I			Model II			Model III		
	AT (n = 49)	AA (n = 21)	TT (n = 40)	AT (n = 45)	AA (n = 18)	TT (n = 37)	AT (n = 34)	AA (n = 17)	TT (n = 28)
Acylated ghrelin delta AUC (2 h pg·mL ⁻¹)	-76.0 (-110.8 to -41.2)	-86.3 (-139.5 to -33.1)	-96.3 (-134.9 to -57.8)	-69.5 (-107.1 to -32.0)	-93.1 (-151.1 to -35.0)	-103.2 (-144.5 to -61.8)	-87.4 (-106.9 to -67.9)	-87.0 (-114.9 to -59.0)	-67.8 (-90.2 to -45.4)
	Eta ² = 0.006 (90% CI: 0.000-0.034), <i>P</i> = 0.740			Eta ² = 0.015 (90% CI: 0.000-0.063), <i>P</i> = 0.494			Eta ² = 0.026 (90% CI: 0.000-0.097), <i>P</i> = 0.414		
Total PYY delta AUC (2 h pg·mL ⁻¹)	101.1 (84.2 to 118.1)	89.7 (63.8 to 115.6)	113.4 (94.7 to 132.2)	98.5 (80.2 to 116.8)	86.5 (58.2 to 114.8)	113.7 (93.5 to 133.8)	103.5 (81.2 to 125.8)	80.4 (48.4 to 112.4)	120.1 (94.4 to 145.7)
	Eta ² = 0.021 (90% CI: 0.000-0.072), <i>P</i> = 0.322			Eta ² = 0.028 (90% CI: 0.000-0.088), <i>P</i> = 0.271			Eta ² = 0.050 (90% CI: 0.000-0.137), <i>P</i> = 0.186		
Insulin delta AUC (2 h pmol·L ⁻¹)	411 (345 to 476)	404 (303 to 503)	432 (359 to 504)	409 (342 to 477)	415 (311 to 519)	430 (356 to 504)	411 (330 to 492)	429 (313 to 545)	463 (370 to 556)
	Eta ² = 0.002 (90% CI: 0.000-0.017), <i>P</i> = 0.875			Eta ² = 0.002 (90% CI: 0.000-0.022), <i>P</i> = 0.921			Eta ² = 0.010 (90% CI: 0.000-0.055), <i>P</i> = 0.728		
Glucose delta AUC (2 h mmol·L ⁻¹)	0.66 (0.21 to 1.12)	0.60 (-0.10 to 1.30)	1.01 (0.51 to 1.52)	0.60 (0.19 to 1.02)	0.54 (-0.09 to 1.18)	0.79 (0.34 to 1.25)	0.68 (0.19 to 1.17)	0.44 (-0.26 to 1.14)	0.88 (0.32 to 1.44)
	Eta ² = 0.012 (90% CI: 0.000-0.054), <i>P</i> = 0.511			Eta ² = 0.006 (90% CI: 0.000-0.036), <i>P</i> = 0.766			Eta ² = 0.013 (90% CI: 0.000-0.066), <i>P</i> = 0.642		
Overall appetite delta AUC (2 h mm)	-79.3 (-89.1 to -69.5)	-72.4 (-87.4 to -57.5)	-79.2 (-90.1 to -68.4)	-75.3 (-85.2 to -65.4)	-73.6 (-88.8 to -58.3)	-82.1 (-93.0 to -71.2)	-73.4 (-85.4 to -61.4)	-75.6 (-92.7 to -58.4)	-75.6 (-89.3 to -61.8)
	Eta ² = 0.006 (90% CI: 0.000-0.036), <i>P</i> = 0.718			Eta ² = 0.012 (90% CI: 0.000-0.056), <i>P</i> = 0.568			Eta ² = 0.001 (90% CI: 0.000-0.021), <i>P</i> = 0.965		

Model I: Univariable model with FTO rs9939609 genotype as single fixed effect. Model II: Multivariable model with FTO rs9939609 genotype as a fixed effect and sex, age, fat mass and visceral adipose tissue as covariates. Model III: Multivariable model with FTO rs9939609 genotype as a fixed effect and sex, age, body mass index, peak oxygen uptake, resting metabolic rate, visceral adipose tissue, abdominal subcutaneous adipose tissue, liver fat, sitting time and moderate-to-vigorous physical activity as covariates.

Values are mean (95% confidence interval (CI)). Eta², 90% CI and *P*-values are from the fixed effect of the FTO rs9939609 genotype group.

Sex-specific Pearson's correlation coefficients

Appetite-related outcomes and individual characteristics

No significant correlations were observed between fasting acylated ghrelin and age, BMI, fat mass, $V\dot{O}_2$ peak, resting metabolic rate, visceral fat, abdominal subcutaneous adipose tissue, liver fat, average sitting or average MVPA in men ($r = -0.18$ to 0.07 , $P \geq 0.185$) or women ($r = -0.19$ to 0.06 , $P \geq 0.175$). Similarly, no significant correlations were observed between fasting total PYY and any of the individual characteristics in men ($r = -0.13$ to 0.14 , $P \geq 0.330$) or women ($r = -0.14$ to 0.10 , $P \geq 0.323$). Pearson's correlation coefficients between individual characteristics and fasting insulin, glucose and leptin are presented in Table 4. In summary, fasting insulin was positively correlated with general and abdominal adiposity parameters in both sexes and with liver fat in men ($r = 0.32$ to 0.53 , $P \leq 0.010$). Fasting insulin was negatively correlated with $V\dot{O}_2$ peak in both sexes and with MVPA in men ($r = -0.35$ to -0.47 , $P \leq 0.004$). Fasting glucose was positively correlated with total and abdominal adiposity parameters in both sexes, with age and liver fat in men, and with resting metabolic rate in women ($r = 0.28$ to 0.44 , $P \leq 0.017$). Fasting glucose was negatively correlated with $V\dot{O}_2$ peak in both sexes ($r = -0.29$ to -0.28 , $P \leq 0.020$). Fasting leptin was positively correlated with general and abdominal adiposity parameters in both sexes, and with age and liver fat in men ($r = 0.24$ to 0.83 , $P \leq 0.040$). Fasting leptin was negatively correlated with $V\dot{O}_2$ peak in both sexes and with MVPA in men ($r = -0.35$ to -0.64 , $P \leq 0.006$). In men, fasting overall appetite was negatively associated with fat mass ($r = -0.31$, $P = 0.022$, 95% CI = -0.53 to -0.05) and abdominal subcutaneous adipose tissue ($r = -0.30$, $P = 0.032$, 95% CI = -0.53 to -0.02). No significant correlations between fasting overall appetite and individual characteristics were observed in women ($r = -0.12$ to 0.09 , $P \geq 0.391$).

Delta AUC for acylated ghrelin was positively associated with sitting time ($r = 0.29$, $P = 0.048$, 95% CI = 0.00 to 0.53) and negatively associated with age ($r = -0.32$, $P = 0.017$, 95% CI = -0.54 to -0.06) in men. Insulin AUC was positively associated with visceral adipose tissue in men ($r = 0.38$, $P = 0.007$, 95% CI = 0.11 to 0.59) and women ($r = 0.32$, $P = 0.021$, 95% CI = 0.05 to 0.55), and with fat mass ($r = 0.39$, $P = 0.003$, 95% CI = 0.14 to 0.59), abdominal subcutaneous adipose tissue ($r = 0.31$, $P = 0.026$, 95% CI = 0.03 to 0.54) and liver fat ($r = 0.47$, $P = 0.001$, 95% CI = 0.21 to 0.66) in men. Insulin AUC was negatively associated with $V\dot{O}_2$ peak ($r = -0.44$, $P = 0.001$, 95% CI = -0.63 to -0.20) and MVPA ($r = -0.38$, $P = 0.007$, 95% CI = -0.60 to -0.11) in

950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008

364 men. None of the correlations between AUC for total PYY, glucose and overall appetite and
365 individual characteristics were statistically significant ($r = -0.23$ to 0.24 , $P \geq 0.061$).

Table 4. Sex-specific Pearson's correlation coefficients between fasting appetite-related blood markers and individual characteristics.

	Fasting insulin (pmol·L ⁻¹)	Fasting glucose (mmol·L ⁻¹)	Fasting leptin (ng·mL ⁻¹)
Age (years)	Men: $r = -0.01$, $P = 0.457$, 95% CI = -0.27 to 0.25 Women: $r = -0.16$, $P = 0.123$, 95% CI = -0.40 to 0.11	Men: $r = 0.34$, $P = 0.005$, 95% CI = 0.08 to 0.55 Women: $r = 0.08$, $P = 0.270$, 95% CI = -0.19 to 0.33	Men: $r = 0.24$, $P = 0.040$, 95% CI = -0.02 to 0.47 Women: $r = -0.07$, $P = 0.298$, 95% CI = -0.33 to 0.20
Body mass index (kg·m ⁻²)	Men: $r = 0.39$, $P = 0.003$, 95% CI = 0.14 to 0.59 Women: $r = 0.53$, $P < 0.001$, 95% CI = 0.31 to 0.69	Men: $r = 0.33$, $P = 0.013$, 95% CI = 0.07 to 0.54 Women: $r = 0.35$, $P = 0.004$, 95% CI = 0.10 to 0.56	Men: $r = 0.62$, $P < 0.001$, 95% CI = 0.43 to 0.76 Women: $r = 0.77$, $P < 0.001$, 95% CI = 0.64 to 0.86
Fat mass (kg)	Men: $r = 0.49$, $P < 0.001$, 95% CI = 0.26 to 0.67 Women: $r = 0.32$, $P = 0.008$, 95% CI = 0.06 to 0.54	Men: $r = 0.44$, $P < 0.001$, 95% CI = 0.20 to 0.63 Women: $r = 0.28$, $P = 0.017$, 95% CI = 0.02 to 0.50	Men: $r = 0.83$, $P < 0.001$, 95% CI = 0.73 to 0.90 Women: $r = 0.75$, $P < 0.001$, 95% CI = 0.61 to 0.85
V̇O ₂ peak (mL·kg ⁻¹ ·min ⁻¹)	Men: $r = -0.47$, $P < 0.001$, 95% CI = -0.65 to -0.24 Women: $r = -0.35$, $P = 0.004$, 95% CI = -0.56 to -0.10	Men: $r = -0.29$, $P = 0.015$, 95% CI = -0.51 to -0.03 Women: $r = -0.28$, $P = 0.020$, 95% CI = -0.50 to -0.02	Men: $r = -0.64$, $P < 0.001$, 95% CI = -0.77 to -0.45 Women: $r = -0.58$, $P < 0.001$, 95% CI = -0.73 to -0.37
Resting metabolic rate (kcal)	Men: $r = -0.04$, $P = 0.381$, 95% CI = -0.31 to 0.23 Women: $r = 0.03$, $P = 0.402$, 95% CI = -0.24 to 0.29	Men: $r = -0.12$, $P = 0.205$, 95% CI = -0.38 to 0.15 Women: $r = 0.35$, $P = 0.005$, 95% CI = 0.09 to 0.56	Men: $r = 0.05$, $P = 0.369$, 95% CI = -0.22 to 0.32 Women: $r = 0.05$, $P = 0.359$, 95% CI = -0.22 to 0.31
Visceral adipose tissue (L)	Men: $r = 0.41$, $P = 0.002$, 95% CI = 0.15 to 0.62 Women: $r = 0.33$, $P = 0.010$, 95% CI = 0.06 to 0.55	Men: $r = 0.42$, $P = 0.001$, 95% CI = 0.15 to 0.63 Women: $r = 0.36$, $P = 0.005$, 95% CI = 0.09 to 0.58	Men: $r = 0.65$, $P < 0.001$, 95% CI = 0.45 to 0.79 Women: $r = 0.62$, $P < 0.001$, 95% CI = 0.42 to 0.76
Abdominal subcutaneous adipose tissue (L)	Men: $r = 0.43$, $P = 0.002$, 95% CI = 0.17 to 0.63 Women: $r = 0.44$, $P = 0.001$, 95% CI = 0.19 to 0.64	Men: $r = 0.39$, $P = 0.005$, 95% CI = 0.13 to 0.60 Women: $r = 0.34$, $P = 0.013$, 95% CI = 0.07 to 0.56	Men: $r = 0.79$, $P < 0.001$, 95% CI = 0.66 to 0.87 Women: $r = 0.79$, $P < 0.001$, 95% CI = 0.66 to 0.87
Liver fat (%)	Men: $r = 0.49$, $P < 0.001$, 95% CI = 0.24 to 0.68 Women: $r = 0.06$, $P = 0.338$, 95% CI = -0.22 to 0.33	Men: $r = 0.33$, $P = 0.010$, 95% CI = 0.05 to 0.56 Women: $r = 0.07$, $P = 0.305$, 95% CI = -0.21 to 0.34	Men: $r = 0.44$, $P = 0.001$, 95% CI = 0.18 to 0.64 Women: $r = 0.18$, $P = 0.112$, 95% CI = -0.11 to 0.44
Average sitting time (min·day ⁻¹)	Men: $r = -0.06$, $P = 0.340$, 95% CI = -0.34 to 0.23 Women: $r = 0.12$, $P = 0.196$, 95% CI = -0.17 to 0.39	Men: $r = -0.12$, $P = 0.210$, 95% CI = -0.39 to 0.17 Women: $r = 0.13$, $P = 0.190$, 95% CI = -0.16 to 0.40	Men: $r = -0.12$, $P = 0.207$, 95% CI = -0.39 to 0.17 Women: $r = 0.05$, $P = 0.353$, 95% CI = -0.23 to 0.33
Average MVPA time (min·day ⁻¹)	Men: $r = -0.44$, $P = 0.001$, 95% CI = -0.64 to -0.18 Women: $r = -0.01$, $P = 0.493$, 95% CI = -0.28 to 0.27	Men: $r = -0.03$, $P = 0.420$, 95% CI = -0.31 to 0.25 Women: $r = 0.09$, $P = 0.274$, 95% CI = -0.19 to 0.36	Men: $r = -0.35$, $P = 0.006$, 95% CI = -0.57 to -0.08 Women: $r = -0.10$, $P = 0.241$, 95% CI = -0.36 to 0.18

AUC, area under the curve; FTO, fat mass and obesity associated gene; MVPA, moderate-to-vigorous physical activity, PYY, peptide YY; V̇O₂ peak, peak oxygen uptake.

1050
1051
1052
1053 *Perceived appetite and appetite-related blood parameters*
1054

1055
1056 Fasting overall appetite was negatively associated with fasting insulin ($r = -0.32$, $P = 0.015$, 95%
1057 CI = -0.54 to -0.06) and fasting leptin ($r = -0.35$, $P = 0.008$, 95% CI = -0.56 to -0.10) in men.
1058
1059 Delta AUC for overall appetite was positively associated with insulin AUC ($r = 0.35$, $P = 0.009$,
1060 95% CI = 0.10 to 0.56) in women. No other significant correlations between overall appetite and
1061
1062 appetite-related blood parameters were evident in the fasted or postprandial state ($r = -0.20$ to
1063
1064 0.26 , $P \geq 0.052$).

1065
1066
1067 **DISCUSSION**
1068

1069 The primary finding of this study is that very little influence of the FTO **rs9939609** genotype
1070
1071 was identified for fasting and postprandial perceived appetite and appetite-related blood
1072
1073 outcomes in healthy men and women. Explained variance for FTO group on all outcomes was
1074
1075 small ($< 5\%$) according to the thresholds suggested by Cohen (1998). Even the upper 90%
1076
1077 confidence limits of the explained variance were low for each outcome ($< 15\%$). In the context
1078
1079 of precision medicine, we maintain that explained variance would need to be much larger than
1080
1081 our observed values for the FTO **rs9939609** gene to be a useful predictor of appetite-related
1082
1083 outcomes. We also found that fasting and postprandial acylated ghrelin and total PYY were not
1084
1085 associated with general or abdominal adiposity, while leptin, glucose and insulin concentrations
1086
1087 were consistently associated with adiposity variables. Our study is the first to employ an
1088
1089 integrative approach to investigate associations between a variety of genetic, physiological and
1090
1091 lifestyle characteristics with appetite-related outcomes. Previous research has provided limited
1092
1093 evidence on the influence of specific individual characteristics on appetite-related blood
1094
1095 parameters and appetite perceptions.

1096 The FTO gene represents the most extensively-studied gene that has been associated with a
1097
1098 higher risk of obesity (Frayling et al. 2007), yet evidence on the physiological mechanisms
1099
1100 involved is limited. The study undertaken by Karra et al. (2013) supported the hypothesis that
1101
1102 satiety control differs between FTO **rs9939609** genotype groups. Specifically, the group with
1103
1104 higher obesity risk (AA) presented attenuated suppression of acylated ghrelin and perceived
1105
1106 hunger after consumption of a meal, which can naturally lead to higher energy intake and,
1107
1108 consequently, higher body mass (Karra et al. 2013). However, our results do not support this
1109
1110 hypothesis as we found very little influence of genotype group on acylated ghrelin concentrations
1111
1112 or perceived appetite ratings. Differences between study samples can possibly explain

1109
1110
1111
1112 398 discrepancies between findings, as Karra et al. (2013) recruited healthy young lean males, while
1113
1114 399 our sample was composed of a heterogeneous group of males and females. Additionally, Karra
1115
1116 400 et al. (2013) selectively sampled their participants in order to match groups for certain variables,
1117
1118 401 whereas we adopted a multivariate-adjusted approach to our data analysis. Interestingly, recent
1119
1120 402 studies have reported lower postprandial total ghrelin concentrations in AA compared to AT and
1121
1122 403 TT individuals (Magno et al. 2018; Melhorn et al. 2018), and postprandial hunger ratings were
1123
1124 404 either similar between genotype groups (Melhorn et al. 2018) or were lower in AA individuals
1125
1126 405 (Magno et al. 2018). These findings were observed despite the AA individuals exhibiting higher
1127
1128 406 energy intake during an *ad libitum* buffet (Melhorn et al. 2018). Of note, the active part of ghrelin
1129
1130 407 (acylated ghrelin) only represents approximately 5 to 10% of total ghrelin (Hosoda et al, 2000;
1131
1132 408 Yoshimoto et al. 2002) and, therefore, the assessment of total ghrelin in these studies could
1133
1134 409 potentially explain the variability in findings.

1135
1136 410 Our research group has recently conducted a replicated crossover study to examine individual
1137
1138 411 appetite responses to meal intake in healthy men recruited according to their FTO rs9939609
1139
1140 412 genotype (AA or TT) (Goltz et al. 2019). The findings from this study highlighted the existence
1141
1142 413 of interindividual variability in perceived appetite and acylated ghrelin, total PYY, insulin and
1143
1144 414 glucose responses to a standardised meal over and above any measurement errors and/or natural
1145
1146 415 variance of the outcomes. However, the magnitude of postprandial appetite parameter responses
1147
1148 416 after meal intake was not influenced by the FTO rs9939609 gene (Goltz et al. 2019). In line with
1149
1150 417 our findings, previous studies have reported no differences between FTO rs9939609 genotype
1151
1152 418 groups for fasting glucose and insulin (Speakman et al. 2008), fasting leptin (Speakman et al.
1153
1154 419 2008; Karra et al. 2013; Melhorn et al. 2018), fasting and postprandial PYY₃₋₃₆ (Karra et al. 2013)
1155
1156 420 and fasting and postprandial GLP-1 (Melhorn et al. 2018). Beyond the subjective appetite and
1157
1158 421 appetite-related blood outcomes assessed in this study, AA and TT individuals have been shown
1159
1160 422 to exhibit divergent neural responsiveness to food cues within homeostatic and reward brain
1161
1162 423 regions in both fasted and postprandial states (Karra et al. 2013). Specifically, AA individuals
1163
1164 424 rated high-energy food images as more appealing than TT individuals, and positive associations
1165
1166 425 between circulating acylated ghrelin and central neural system responsiveness to food cues were
1167
1168 426 observed only in TT individuals (Karra et al. 2013). Moreover, recent evidence suggests that AA
1169
1170 427 individuals show higher total food cravings, compared to TT individuals, which correlated with
1171
1172 428 BMI (Dang et al. 2018). Additional studies are needed to elucidate the precise role that FTO
1173
1174 429 rs9939609 plays in moderating appetite control and energy intake which include both central and
1175
1176 430 peripheral factors implicated in appetite regulation.

1168
1169
1170
1171 431 Although evidence to date suggests a negligible impact of FTO rs9939609 genotype on energy
1172
1173 432 expenditure, higher levels of physical activity seem to exert a protective effect on the obesity risk
1174
1175 433 associated with FTO (Sonestedt et al. 2009; Speakman, 2015). On the contrary, diets with higher
1176
1177 434 fat content can exacerbate the susceptibility to obesity linked to the FTO rs9939609 high-risk
1178
1179 435 genotype (Sonestedt et al. 2009; Speakman, 2015). Our study included objectively assessed
1180
1181 436 sitting time, MVPA and cardiorespiratory fitness as covariates in the statistical analyses.
1182
1183 437 However, only 20% of our participants accumulated, on average, less than 30 min of MVPA per
1184
1185 438 day, indicating that most participants in our sample had relatively high levels of physical activity.
1186
1187 439 Therefore, we cannot rule out the possibility of this hindering our ability to detect differences in
1188
1189 440 appetite-related outcomes between the genotype groups (Speakman et al. 2008). Our study did
1190
1191 441 not include any assessment of habitual dietary intake and, therefore, fat intake was not taken into
1192
1193 442 consideration in our analyses. Nevertheless, it is well known that the currently available dietary
1194
1195 443 intake assessment tools do not provide reliable data, and this currently represents a major
1196
1197 444 challenge for those involved in nutrition-related research, clinical practice or policy development
1198
1199 445 (Dhurandhar et al. 2015; Archer et al. 2018).

1195 446 In contrast to previous studies (Alajmi et al. 2016; Douglas et al. 2017), we did not observe a
1196
1197 447 statistically significant difference in fasting concentrations of acylated ghrelin between men and
1198
1199 448 women. The reason for this disparity is unclear but it is worth noting that two female participants
1200
1201 449 were identified as clear outliers within our sample, with fasting acylated ghrelin concentrations
1202
1203 450 of 2,899 and 4,411 pg·mL⁻¹. These extremely high concentrations of acylated ghrelin were
1204
1205 451 observed consistently in all four samples collected for each participant, indicating these values
1206
1207 452 represented physiological characteristics of these two individuals rather than merely one-off
1208
1209 453 measurement errors. Further studies are needed to investigate potential causes and consequences
1210
1211 454 of such extreme concentrations of acylated ghrelin, and care should be taken when interpreting
1212
1213 455 group mean results, as group means can be greatly impacted by such outliers. Nevertheless,
1214
1215 456 exclusion of the outliers did not influence any of the statistical models in this study and, therefore,
1216
1217 457 data are presented with the outliers included. Higher concentrations of fasting glucose were
1218
1219 458 observed in men than women in the current study, which may be indicative of a greater degree
1220
1221 459 of insulin resistance resulting from the higher visceral adipose tissue and liver fat levels observed
1222
1223 460 in men (Marchesini et al, 2001; Ibrahim, 2010). Higher levels of fasting leptin were observed in
1224
1225 461 women, likely because of the higher fat mass values in relation to total body mass in women,
1226
1227 462 compared to men (Marshall et al. 2000; Rosenbaum and Leibel, 2014).

1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285

463 After meal consumption, greater changes in acylated ghrelin and overall appetite were observed
464 in women than men. It should be noted that all participants received an identical standardised
465 meal and, as women had significantly lower body mass and fat free mass, and consequently lower
466 resting metabolic rate, it was expected that the postprandial suppression of appetite would be
467 stronger in women. However, it is interesting to observe that, apart from acylated ghrelin, no
468 other statistically significant differences were observed between men and women in any of the
469 remaining postprandial appetite-related blood parameters. Previous evidence has demonstrated a
470 stronger suppression of acylated ghrelin in women than men after acute exercise and standardised
471 meals (Douglas et al. 2017), but not after the consumption of a standardised liquid meal (Carroll
472 et al. 2007).

473 Our exploratory analyses did not identify any statistically significant or meaningful association
474 between adiposity parameters and fasting or postprandial concentrations of acylated ghrelin and
475 total PYY. This is in contrast with findings from previous studies which demonstrated a lower
476 postprandial suppression of total and acylated ghrelin (Le Roux et al. 2005; Carrol et al. 2007)
477 and a blunted postprandial elevation in PYY (Le Roux et al. 2006) in individuals with obesity.
478 However, as expected, fasting insulin, glucose and leptin and postprandial insulin were all
479 positively associated with general and visceral adiposity, demonstrated by moderate to very large
480 correlation coefficients, which is consistent with the well-established role of leptin in signalling
481 adiposity levels (Rosenbaum and Leibel, 2014) and the impact of adiposity on insulin resistance
482 (Ibrahim, 2010). Additionally, fat free mass, which represents the largest determinant of resting
483 metabolic rate, has been identified as a primary determinant of appetite and energy intake
484 (Blundell et al. 2015b). However, our findings did not reveal any significant associations of
485 appetite-related hormones or perceived appetite with resting metabolic rate.

486 While acute bouts of exercise have been shown consistently to transiently suppress appetite (King
487 et al. 2017), chronic exercise and high levels of physical activity have been suggested to increase
488 the overall drive to eat and, concomitantly, to increase the satiating effect of a standardised meal
489 (King et al. 2009; Beaulieu et al. 2016). We did not identify any significant associations between
490 habitual physical activity levels and fasting or postprandial acylated ghrelin, total PYY, glucose
491 or perceived appetite. However, a negative association was observed between MVPA and fasting
492 leptin and insulin, and postprandial insulin in men. **Additionally, negative associations between**
493 **V̇O₂ peak and fasting and postprandial insulin, fasting glucose and fasting leptin were observed.**
494 **Acute and chronic exercise augments insulin sensitivity by increasing insulin-like growth factor**

1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344

1, and individuals with higher cardiorespiratory fitness typically show higher insulin sensitivity (Borghouts and Keizer, 2000; Castro et al. 2016). Furthermore, a recent meta-analysis showed that leptin concentrations can be reduced by exercise in individuals who are overweight even in the absence of dietary interventions or major weight loss (BMI reduction of > 2.5%) (Rostás et al. 2017). Postprandial acylated ghrelin was positively associated with sitting time in men, but this correlation was small in magnitude and would not be considered significant if the stricter threshold of $P < 0.005$ was applied in line with recent recommendations (Benjamin et al. 2017). Perceived fasting overall appetite was negatively associated with total fat mass in men supporting previous evidence suggesting the existence of negative feedback signals originating from fat mass in order to regulate appetite and maintain body weight (Weise et al. 2014; Blundell et al. 2015a). However, no association was observed between postprandial perceived appetite and any adiposity parameter in our study. Interestingly, no statistically significant associations between fasting or postprandial perceived overall appetite and acylated ghrelin or total PYY were identified. Even though circulating concentrations of acylated ghrelin and PYY vary on a meal-to-meal basis, concomitantly with perceived appetite, the magnitude and direction of the changes in hormone concentrations are not always mirrored by changes in perceived appetite (Goltz et al. 2018). In contrast, postprandial overall appetite AUC was positively associated with postprandial insulin AUC in women, which is consistent with previous findings showing that postprandial insulin concentrations are positively associated with postprandial satiety and negatively associated with postprandial hunger (Flint et al. 2007). The strengths of our study include the use of an integrative approach and objective assessment methods to explore the associations of the FTO rs9939609 genotype with fasting and postprandial appetite-related hormones and perceived appetite, taking into consideration a variety of individual characteristics that have been previously suggested to influence appetite parameters. Furthermore, the recruitment of a highly heterogeneous sample for parameters such as age, adiposity and cardiorespiratory fitness levels adds strength to our analyses. Finally, the careful standardisation of diet and physical activity in the 24 h preceding the laboratory visit, as well as the inclusion of a cannula acclimatisation period, also contributed to the quality of the study outcome measurements obtained. However, it should be highlighted that our study employed an exploratory approach and the cross-sectional design makes it impossible to imply any causation in our results. Our results may have been compromised by the reduced sample size and by the loss of power in some of the statistical models due to missing data. Additionally, it is possible

1345
1346
1347
1348 527 that a study design where individuals are exposed to an obesigenic food environment, such as an
1349
1350 528 *ad libitum* buffet meal rather than a standardised meal stimulus, may be more appropriate to
1351
1352 529 elucidate the effect of FTO rs9939609 genotype on food choice and eating behaviour.
1353
1354 530 Furthermore, participants were aware of the meal timing so it is possible that the higher
1355
1356 531 preprandial ghrelin concentrations reflected an anticipatory response to impending meal intake
1357
1358 532 (Cummings et al. 2001). Future studies should consider isolating meal provision from time-
1359
1360 533 related cues and/or examining the influence of cephalic phase ghrelin release during meal
1361
1362 534 anticipation on postprandial appetite responses.

1363 535 In conclusion, the FTO rs9939609 genotype did not have any significant influence on fasting or
1364
1365 536 postprandial perceived appetite or appetite-related blood parameters in healthy men and women.
1366
1367 537 The associations between fasting and postprandial acylated ghrelin, total PYY and general or
1368
1369 538 abdominal adiposity were also small, while fasting leptin, glucose and insulin and postprandial
1370
1371 539 insulin concentrations were consistently and positively associated with adiposity outcomes.
1372
1373 540 Further research is needed to clarify the precise role of the FTO rs9939609 genotype in
1374
1375 541 moderating appetite control and energy intake, including both physiological and psychological
1376
1377 542 factors that influence eating behaviour. Specifically, well-controlled long-term studies are
1378
1379 543 needed to improve understanding of the effect of the FTO rs9939609 genotype on appetite and
1380
1381 544 energy intake during and after interventions targeting weight loss and/or prevention of weight
1382
1383 545 gain. Understanding the complex interaction between genetics and other individual
1384
1385 546 characteristics, physiological appetite parameters and perceived appetite is of crucial importance
1386
1387 547 for planning targeted strategies for weight control.

1388 548 1389 549 **ACKNOWLEDGEMENTS**

1390 550 This research was funded by the NIHR Leicester Biomedical Research Centre. The views
1391
1392 551 expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the
1393
1394 552 Department of Health and Social Care.

1395 553 1396 554 **REFERENCES**

1397 555 Alajmi N, Deighton K, King JA, Reischak-Oliveira A, Wasse LK, Jones J, et al. Appetite and
1398
1399 556 energy intake responses to acute energy deficits in females versus males. *Med Sci Sports
1400
1401
1402
1403* Exerc. 2016;48(3):412-20.

1404
1405
1406
1407 558 Archer E, Lavie CJ, Hill JO. The failure to measure dietary intake engendered a fictional
1408
1409 559 discourse on diet-disease relations. *Front Nutr.* 2018;5:105.
1410
1411 560 Atkin AJ, Gorely T, Clemes SA, Yates T, Edwardson C, Brage S, et al. Methods of
1412
1413 561 measurement in epidemiology: sedentary behaviour. *Int J Epidemiol.* 2012;41(5):1460-71.
1414
1415 562 Atkinson G, Batterham AM. True and false interindividual differences in the physiological
1416
1417 563 response to an intervention. *Exp Physiol* 2015;100(6):577-88.
1418
1419 564 Beaulieu K, Hopkins M, Blundell J, Finlayson G. Does habitual physical activity increase the
1420
1421 565 sensitivity of the appetite control system? A systematic review. *Sports Med.* 2016;46(12):1897-
1422
1423 566 919.
1424
1425 567 Benjamin DJ, Berger J, Johannesson M, Nosek BA, Wagenmakers E, Berk R, et al. Redefine
1426
1427 568 statistical significance. *PsyArXiv* (online). 2017. Available at:
1428 569 <https://doi.org/10.31234/osf.io/mky9j> (Accessed on: January 11, 2019).
1429
1430 570 Bland JM, Altman DG. Transforming data. *BMJ.* 1996;23;312(7033):770.
1431
1432 571 Blundell JE, Finlayson G, Gibbons C, Caudwell P, Hopkins M. The biology of appetite control:
1433
1434 572 Do resting metabolic rate and fat-free mass drive energy intake? *Physiol Behav.* 2015;152(Pt
1435
1436 573 B):473-8.
1437
1438 574 Blundell JE, Gibbons C, Caudwell P, Finlayson G, Hopkins M. Appetite control and energy
1439
1440 575 balance: impact of exercise. *Obes Rev.* 2015;16 Suppl 1:67-76.
1441
1442 576 Blundell JE, Levin F, King NA, Barkeling B, Gustafsson T, Hellstrom PM, et al.
1443
1444 577 Overconsumption and obesity: peptides and susceptibility to weight gain. *Regul*
1445
1446 578 *Pept.* 2008;149(1-3):32-8.
1447
1448 579 Borg GA. Perceived exertion. A note on “history” and methods. *Med Sci Sports.* 1973;5(2):90–
1449
1450 580 3.
1451
1452 581 Borga M, Thomas EL, Romu T, Rosander J, Fitzpatrick J, Dahlqvist Leinhard O, et al.
1453
1454 582 Validation of a fast method for quantification of intra-abdominal and subcutaneous adipose
1455
1456 583 tissue for large-scale human studies. *NMR Biomed.* 2015;28(12):1747-53.
1457
1458 584 Borghouts LB, Keizer HA. Exercise and insulin sensitivity: a review. *Int J Sports*
1459
1460 585 *Med.* 2000;21(1):1-12.

1463
1464
1465
1466 586 Carroll JF, Kaiser KA, Franks SF, Deere C, Caffrey JL. Influence of BMI and gender
1467 on postprandial hormone responses. *Obesity*. 2007;15(12):2974-83.
1468 587
1469
1470 588 Castro MG, Venutolo C, Yau PL, Convit A. Fitness, insulin sensitivity, and frontal lobe integrity
1471 in adults with overweight and obesity. *Obesity*. 2016;24(6):1283-9.
1472 589
1473
1474 590 Chandarana K, Drew ME, Emmanuel J, Karra E, Gelegen C, Chan P, et al. Subject
1475 standardization, acclimatization, and sample processing affect gut hormone levels and appetite
1476 591 in humans. *Gastroenterology*. 2009;136(7):2115-26.
1477 592
1478
1479 593 Cohen J. *Statistical power analysis for the behavioral sciences*. 1988. 2nd ed. Hillsdale, NJ:
1480 Lawrence Erlbaum Associates.
1481 594
1482
1483 595 Cummings DE, Purnell JQ, Frayo RS, Schmidova K, Wisse BE, Weigle DS. A preprandial rise
1484 in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes*. 2001;50(8):1714-
1485 596 9.
1486 597
1487
1488
1489 598 Dang LC, Samanez-Larkin GR, Smith CT, Castrellon JJ, Perkins SF, Cowan RL, et al.
1490 FTO affects food cravings and interacts with age to influence age-related decline in food
1491 599 cravings. *Physiol Behav*. 2018;192:188-93.
1492 600
1493
1494 601 Deighton K, Batterham RL, Stensel DJ. Appetite and gut peptide responses to exercise and
1495 calorie restriction. The effect of modest energy deficits. *Appetite*. 2014;81:52-9.
1496 602
1497
1498 603 Deighton K, Karra E, Batterham RL, Stensel DJ. Appetite, energy intake, and PYY₃₋₃₆
1499 responses to energy-matched continuous exercise and submaximal high-intensity exercise.
1500 *Appl Physiol Nutr Metab*. 2013;38(9):947-52
1501 604
1502
1503 605
1504 606 Dill DB, Costill DL. Calculation of percentage changes in volumes of blood, plasma, and red
1505 cells in dehydration. *J Appl Physiol*. 1974;37:247-8.
1506 607
1507
1508 608 Douglas JA, King JA, Clayton DJ, Jackson AP, Sargeant JA, Thackray AE, et al. Acute effects
1509 of exercise on appetite, ad libitum energy intake and appetite-regulatory hormones in lean and
1510 609 overweight/obese men and women. *Int J Obes*. 2017;41(12):1737-44.
1511 610
1512
1513 611 Dhurandhar NV, Schoeller D, Brown AW, Heymsfield SB, Thomas D, Sørensen TI, et al.
1514 Energy balance measurement: when something is not better than nothing. *Int J Obes*.
1515 612 2015;39(7):1109-13.
1516 613
1517
1518
1519
1520
1521

1522
1523
1524
1525 614 Flint A, Gregersen NT, Gluud LL, Møller BK, Raben A, Tetens I, et al. Associations between
1526
1527 615 postprandial insulin and blood glucose responses, appetite sensations and energy intake in
1528
1529 616 normal weight and overweight individuals: a meta-analysis of test meal studies. *Br J*
1530
1531 617 *Nutr* 2007;98(1):17-25.

1532
1533 618 Flint A, Raben A, Blundell JE, Astrup A. Reproducibility, power and validity of visual analogue
1534
1535 619 scales in assessment of appetite sensations in single test meal studies. *Int J Obes*. 2000;24(1):38–
1536
1537 620 48.

1538
1539 621 Flom P, Cassell D. Stopping stepwise: Why stepwise and similar selection methods are bad, and
1540
1541 622 what you should use. *NESUG 2007 Proceedings: Statistics and data analysis*.

1542
1543 623 Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, et al. A
1544
1545 624 common variant in the FTO gene is associated with body mass index and predisposes to
1546
1547 625 childhood and adult obesity. *Science*. 2007;316(5826):889-94.

1548
1549 626 Gelman A, Pardoe I. Average predictive comparisons for models with nonlinearity, interactions,
1550
1551 627 and variance components. *Sociol Methodol*. 2007;37(1):23-51.

1552
1553 628 Goltz FR, Thackray AE, Atkinson G, Lolli L, King JA, Dorling JL, et al. True interindividual
1554
1555 629 variability exists in postprandial appetite responses in healthy men but is not moderated by the
1556
1557 630 FTO genotype. *J Nutr*. 2019 [in press].

1558
1559 631 Goltz FR, Thackray AE, King JA, Dorling JL, Atkinson G, Stensel DJ. Interindividual
1560
1561 632 responses of appetite to acute exercise: a replicated crossover study. *Med Sci Sports*
1562
1563 633 *Exerc*. 2018;50(4):758-68.

1564
1565 634 Hosoda H, Kojima M, Matsuo H, Kangawa K. Ghrelin and des-acyl ghrelin: two major forms of
1566
1567 635 rat ghrelin peptide in gastrointestinal tissue. *Biochem Biophys Res Commun* 2000;279(3):909-
1568
1569 636 13.

1570
1571 637 Ibrahim MM. Subcutaneous and visceral adipose tissue: structural and functional differences.
1572
1573 638 *Obes Rev*. 2010;11(1):11-8.

1574
1575 639 Jackson AS, Pollock ML, Ward A. Generalized equations for predicting body density of women.
1576
1577 640 *Med Sci Sports Exerc*.1980;12(3):175-81.

1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639

641 Jackson AS, Pollock ML. Generalized equations for predicting body density of men. *Br J Nutr.* 1978;40(3):497-504.

642

643 Karra E, O'Daly OG, Choudhury AI, Yousseif A, Millership S, Neary MT, et al. A link between FTO, ghrelin, and impaired brain food-cue responsivity. *J Clin Invest.* 2013;123(8):3539-51.

644

645 King JA, Deighton K, Broom DR, Wasse LK, Douglas JA, Burns SF, et al. Individual variation in hunger, energy intake, and ghrelin responses to acute exercise. *Med Sci Sports Exerc.* 2017;49(6):1219-28.

646

647

648 King NA, Caudwell PP, Hopkins M, Stubbs JR, Naslund E, Blundell JE. Dual-process action of exercise on appetite control: increase in orexigenic drive but improvement in meal-induced satiety. *Am J Clin Nutr.* 2009;90(4):921-7.

649

650

651 Kline RB. *Beyond significance testing: Reforming data analysis methods in behavioral research.* 2004. Washington, DC: American Psychological Association.

652

653 Kozey-Keadle S, Libertine A, Lyden K, Staudenmayer, J, Freedson, PS. Validation of wearable monitors for assessing sedentary behavior. *Med Sci Sports Exerc.* 2011;43(8):1561-7.

654

655 Le Roux CW, Batterham RL, Aylwin SJ, Patterson M, Borg CM, Wynne KJ, et al. Attenuated peptide YY release in obese subjects is associated with reduced satiety. *Endocrinology.* 2006;147(1):3-8.

656

657

658 Le Roux CW, Patterson M, Vincent RP, Hunt C, Ghatei MA, Bloom SR. Postprandial plasma ghrelin is suppressed proportional to meal calorie content in normal-weight but not obese subjects. *J Clin Endocrinol Metab.* 2005;90(2):1068-71.

659

660

661 MacLean PS, Blundell JE, Mennella JA, Batterham RL. Biological control of appetite: A daunting complexity. *Obesity (Silver Spring).* 2017;25;Suppl 1:S8-S16.

662

663 Magno FCCM, Guaraná HC, Fonseca ACP, Cabello GMK, Carneiro JRI, Pedrosa AP, et al. Influence of FTO rs9939609 polymorphism on appetite, ghrelin, leptin, IL6, TNF α levels, and food intake of women with morbid obesity. *Diabetes Metab Syndr Obes.* 2018;11:199-207.

664

665

666 Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M, et al. Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes.* 2001;50(8):1844-50.

667

1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698

668 Marshall JA, Grunwald GK, Donahoo WT, Scarbro S, Shetterly SM. Percent body fat and lean mass explain the gender difference in leptin: analysis and interpretation of leptin in hispanic and non-hispanic white adults. *Obes Res.* 2000;8(8):543-52.

671 Melhorn SJ, Askren MK, Chung WK, Kratz M, Bosch TA, Tyagi V, et al. FTO genotype impacts food intake and corticolimbic activation. *Am J Clin Nutr.* 2018;107(2):145-54.

673 Neary MT, Batterham RL. Gut hormones: implications for the treatment of obesity. *Pharmacol Ther.* 2009;124(1):44-56.

675 Rosenbaum M, Leibel RL. Role of leptin in energy homeostasis in humans. *J Endocrinol.* 2014;223(1):T83-96.

677 Rostás I, Pótó L, Mátrai P, Hegyi P, Tenk J, Garami A, et al. In middle-aged and old obese patients, training intervention reduces leptin level: A meta-analysis. *PLoS One.* 2017;12(8):e0182801.

680 Senn S. Mastering variation: variance components and personalised medicine. *Stat Med* 2016;35:966–77.

682 Shiiya T, Nakazato M, Mizuta M, Date Y, Mondal MS, Tanaka M, et al. Plasma ghrelin levels in lean and obese humans and the effect of glucose on ghrelin secretion. *J Clin Endocrinol Metab.* 2002;87(1):240–4.

685 Siri WE. Body composition from fluid space and density. In Brozek J & Hanschel A (Eds.), *Techniques for measuring body composition.* 1961;223-44. Washington, DC: National Academy of Science.

688 Sonestedt E, Roos C, Gullberg B, Ericson U, Wirfält E, Orho-Melander M. Fat and carbohydrate intake modify the association between genetic variation in the FTO genotype and obesity. *Am J Clin Nutr.* 2009;90(5):1418-25.

691 Sondergaard E, Gormsen LC, Nellemann B, Vestergaard ET, Christiansen JS, Nielsen S. Visceral fat mass is a strong predictor of circulating ghrelin levels in premenopausal women. *Eur J Endocrinol.* 2009;160(3):375-9.

694 Speakman JR. The 'fat mass and obesity related' (FTO) gene: mechanisms of impact on obesity and energy balance. *Curr Obes Rep.* 2015;4(1):73-91.

1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757

696 Speakman JR, Rance KA, Johnstone AM. Polymorphisms of the FTO gene are associated with
697 variation in energy intake, but not energy expenditure. *Obesity*. 2008;16(8):1961-5.

698 Steiger JH. Beyond the F test: Effect size confidence intervals and tests of close fit in the analysis
699 of variance and contrast analysis. *Psychol Methods*. 2004;9(2):164-82

700 Stubbs RJ, Hughes DA, Johnstone AM, Rowley E, Reid C, Elia M, et al. The use of visual
701 analogue scales to assess motivation to eat in human subjects: a review of their reliability and
702 validity with an evaluation of new hand-held computerized systems for temporal tracking of
703 appetite ratings. *Br J Nutr*. 2000;84: 405–15.

704 Tschöp M, Weyer C, Tataranni PA, Devanarayan V, Ravussin E, Heiman ML. Circulating
705 ghrelin levels are decreased in human obesity. *Diabetes*. 2001;50(4):707-9.

706 Yoshimoto A, Mori K, Sugawara A, Mukoyama M, Yahata K, Suganami T, et al. Plasma ghrelin
707 and desacyl ghrelin concentrations in renal failure. *J Am Soc Nephrol* 2002;13(11):2748-52.

708 Weise CM, Hohenadel MG, Krakoff J, Votruba SB. Body composition and energy expenditure
709 predict ad-libitum food and macronutrient intake in humans. *Int J Obes*. 2014;38(2):243-51.

710 West J, Dahlqvist Leinhard O, Romu T, Collins R, Garratt S, Bell JD, et al. Feasibility of MR-
711 based body composition analysis in large scale population studies. *PLoS*
712 *One*. 2016;11(9):e0163332.

713 Whittingham MJ, Stephens PA, Bradbury RB, Freckleton RP. Why do we still use stepwise
714 modelling in ecology and behaviour? *J Anim Ecol*. 2006;75(5):1182-9.